

# ANNUAL STOCK ASSESSMENT AND FISHERY EVALUATION REPORT: MARIANA ARCHIPELAGO FISHERY ECOSYSTEM PLAN 2018



Western Pacific Regional Fishery Management Council  
1164 Bishop St., Suite 1400  
Honolulu, HI 96813

PHONE: (808) 522-8220  
FAX: (808) 522-8226

[www.wpcouncil.org](http://www.wpcouncil.org)

*The ANNUAL STOCK ASSESSMENT AND FISHERY EVALUATION REPORT for the MARIANA ARCHIPELAGO FISHERY ECOSYSTEM 2018 was drafted by the Fishery Ecosystem Plan Team. This is a collaborative effort primarily between the Western Pacific Regional Fishery Management Council (WPRFMC), National Marine Fisheries Service (NMFS)-Pacific Island Fisheries Science Center (PIFSC), Pacific Islands Regional Office (PIRO), Division of Aquatic Resources (HI) Department of Marine and Wildlife Resources (American Samoa), Division of Aquatic and Wildlife Resources (Guam), and Division of Fish and Wildlife (CNMI).*

*This report attempts to summarize annual fishery performance looking at trends in catch, effort and catch rates as well as provide a source document describing various projects and activities being undertaken on a local and federal level. The report also describes several ecosystem considerations, including fish biomass estimates, biological indicators, protected species, habitat, climate change, and human dimensions. Information like marine spatial planning and best scientific information available for each fishery are described. This report provides a summary of annual catches relative to the Annual Catch Limits established by the Council in collaboration with the local fishery management agencies.*

**Edited By:** Thomas Remington, Marlowe Sabater, Asuka Ishizaki, and Sylvia Spalding, WPRFMC.

**This document can be cited as follows:**

*WPRFMC, 2019. Annual Stock Assessment and Fishery Evaluation Report for the Mariana Archipelago Fishery Ecosystem Plan 2018. Remington, T., Sabater, M., Ishizaki, A., Spalding, S. (Eds.) Western Pacific Regional Fishery Management Council. Honolulu, Hawaii 96813 USA. 276 pp. + Appendices.*

The **Western Pacific Regional Fishery Management Council** acknowledges the valuable contributions of the following Plan Team members for drafting sections of this report:

**Guam Division of Aquatic and Wildlife Resources:** Brent Tibbatts.

**CNMI Division of Fish and Wildlife:** Michael Tenorio, Keena Leon Guerrero, and William Dunn.

**NMFS Pacific Islands Fisheries Science Center:** Justin Hospital, Ivor Williams, Joe O'Malley, Michael Parke, Tom Oliver, Hannah Barkley, Frank Parrish, T. Todd Jones, Minling Pan, and Kirsten Leong.

**NMFS Pacific Islands Regional Office:** Brett Schumacher and Rebecca Walker.

The Council also acknowledges the staff of the **NMFS PIFSC Western Pacific Fisheries Information Network (WPacFIN)** for providing the technical support to generate the data summaries.

The Council would like to thank the following individuals for their contributions to the report: Samuel Kahng and Alexander Filardo.

This page was intentionally left blank.

## **EXECUTIVE SUMMARY**

As part of its five-year fishery ecosystem plan (FEP) review, the Council identified the annual reports as a priority for improvement. The former annual reports have been revised to meet National Standard regulatory requirements for Stock Assessment and Fishery Evaluation (SAFE) reports. The purpose of the reports is twofold: to monitor the performance of the fishery and ecosystem to assess the effectiveness of the FEP in meeting its management objectives; and to maintain the structure of the FEP living document. The reports are comprised of three chapters: fishery performance, ecosystem considerations, and data integration. The Council will iteratively improve the annual SAFE report as resources allow.

The Fishery Performance chapter of this report first presents a general description of the local fishery within Commonwealth of Northern Mariana Islands (CNMI) and Guam, including both the bottomfish and coral reef management unit species (MUS). The fishery data collection system is then explained, encompassing shore-based and boat-based creel surveys, commercial receipt books, and boat inventories. Fishery meta-statistics for each MUS are organized into a summary dashboard table showcasing the values for the most recent fishing year and a comparison to short-term (10-year) and long-term (20-year) averages. Time series for catch and effort statistics are also provided alongside annual catch limit determinations. For 2018 catch in CNMI, only the slipper lobster exceeded their overfishing limit (OFL), allowable biological catch (ABC), or annual catch limit (ACL). Slipper lobsters had not appeared in the catch record until 2016, and now have exceeded the ACL for three consecutive years. This can likely be attributed to the implementation of the Territory Science Initiative (TSI) project that aimed to improve the reporting and compliance to the commercial receipt book data collection program by the Saipan fish vendors. For 2018 catch in Guam, no MUS were identified that had a recent average catch above the thresholds stipulated by the OFL, ABC, and ACL. ACLs were not specified by NMFS for the coral reef ecosystem MUS in 2017 because NMFS had recently acquired new information that require additional environmental analyses to support the Council's ACL recommendations for these MUS (50 CFR Part 665). Similarly in 2018, no ACLs were implemented for the same reason, plus the passing of the ecosystem components amendment that re-classified nearly all reef-associated species as ecosystem components that do not require ACLs.

For the Mariana Archipelago, the main fisheries monitored are the bottomfish, crustacean, and coral reef fisheries. In CNMI, catch with the bottomfishing gear notably decreased in 2018 considering both all species as well as BMUS only. Bottomfishing CPUE had a significant decrease of 85% from the recent 10-year average. Fishing effort, fishery participation, and fishery bycatch continued to decrease in the last year relative recent decline trends. For the coral reef fisheries, statistics for shore-based and boat-based fisheries are shown separately. The estimated CPUEs in 2018 for both shore- and boat-based reef fisheries in CNMI generally showed increasing recent trends despite several being below their 10-year averages. The fishery participation (number of gear hours) in both shore- and boat-based coral reef fisheries decreased for the most part, but there was a noted increase in the number of shore-based spearfishers in 2018 compared to recent averages. The number of fishing participants in 2018 generally decreased for both the shore- and boat-based fisheries, but there was a significant increase in the number of boat-based spearfishers. Coral reef bycatch in CNMI has been decreasing in both fisheries over time.

For Guam in 2018, the bottomfish fishery catch was roughly 32,750 lbs. and exhibited a 6% increase in all species catch alongside an 11% increase in BMUS catch relative to the recent short-term trend. No commercial catch trends were given in this report due to issues with data confidentiality (i.e., less than three vendors that reported data). There were general decreases in 2018 bottomfishing CPUE relative to both of the recent 10- and 20-year averages. The total estimated number of fishing trips for bottomfish decreased by 36% compared both short- and long-term averages to 697, though the number of fishers in 2018 showed a slight increase (+6%) relative to recent averages. Bottomfish bycatch statistics decreased in 2018 relative to short- and long-term trends. The Guam coral reef shore- and boat-based fisheries had large declines in catch and CPUE in 2018 relative to both short- and long-term trends despite showing increases in catch from 2017 levels. CPUE for the boat-based coral reef fishery declined relative to its recent averages, whereas pounds caught per gear hour in the shore-based coral reef fishery significantly increased. The fishing effort estimates in 2018 were generally down relative to 10- and 20-year averages except for the boat-based trolling, which showed an increase to nearly 7 million gear hours. Participation was mixed across fisheries and gear types, though the most notable changes included a large increase in gear hours for boat-based gill netting and a large decrease in participants for boat-based gill netting. Coral reef fishery bycatch was down roughly 10-15% compared to short- and long-term averages, but was part of a gradual increasing trend in Guam over the past decade.

An Ecosystem Considerations chapter was added to the annual SAFE report following the Council's review of its fishery ecosystem plans and revised management objectives. Fishery independent ecosystem survey data, human dimensions, protected species, climate and oceanographic, essential fish habitat, and marine planning information are included in the ecosystem considerations section. Fishery independent ecosystem survey data was acquired through visual surveys conducted in CNMI, Pacific Remote Island Area, American Samoa, Guam, Main Hawaiian Islands, and Northwestern Hawaiian Islands. This report illustrates the mean fish biomass for the reef areas within these locations. Additionally, the mean reef fish biomass and mean size of fishes (>10 cm) for CNMI and Guam are presented by sampling year and reef area. Finally, the reef fish population estimates for each study site within CNMI and Guam are provided for hard bottom habitat (0-30 m).

For CNMI, life history parameters including maximum age, asymptotic length, growth coefficient, hypothetical age at length zero, natural mortality, age at 50% maturity, age at sex switching, length at which 50% of a fish species are capable of spawning, and length of sex switching are provided for 10 species of reef fish and 11 species of bottomfish. The same nine life history parameters are provided for 12 reef species and 11 bottomfish in Guam.

Summarized length derived parameters for coral reef fish and bottomfish in CNMI and Guam include: maximum fish length, mean length, sample size, sample size for L-W regression, and length-weight coefficients. Values for 25 coral reef fish species and 10 bottomfish species are presented for CNMI. Values for 22 coral reef fish species and three bottomfish species are presented for Guam.

The socioeconomics section outlines the pertinent economic, social, and community information available for assessing the successes and impacts of management measures or the achievements of the Fishery Ecosystem Plan for the Mariana Archipelago. It meets the objective "Support

Fishing Communities” adopted at the 165<sup>th</sup> Council meeting; specifically, it identifies the various social and economic groups within the region’s fishing communities and their interconnections. The section begins with an overview of the socioeconomic context for the region, and then provides a summary of relevant studies and data for Mariana Islands, followed by summaries of relevant studies and data for each fishery within the Mariana Archipelago. Socioeconomics data will be included in later versions of this report as resources allow.

There were no new data reported for the crustacean or precious coral fisheries in the CNMI or Guam. Considering the CNMI bottomfish fishery, there was an estimated total of 4,612 pounds sold for \$21,994. The average cost of a bottomfish trip in CNMI was not available in 2018; the cost per trip in 2017 was \$38 down from \$65 in 2016. Considering the CNMI reef fishery, there was an estimated 29,006 pounds sold for \$82,547. Note that data on the cost per coral reef trips in the CNMI was considered confidential in 2018. Considering Guam’s bottomfish fishery in 2018, there was an estimated 3,557 pounds sold for \$17,022. The average cost of a bottomfish trip decreased from 2017 to \$57 compared to \$72; this is likely closely related to fuel cost per trip, which dropped from \$35 to \$27 from 2017 to 2018. Considering Guam’s reef fishery, there was 133,941 pounds sold in 2018 for \$392,548. Note that data on the cost per coral reef trips in the Guam was also considered confidential in 2018.

The protected species section of this report summarizes information and monitors protected species interactions in fisheries managed under the Mariana Archipelago FEP. These fisheries generally have limited impacts to protected species, and do not have federal observer coverage. Consequently, this report tracks fishing effort and other characteristics to detect potential changes to the level of impacts to protected species. Fishery performance data contained in this report indicate that there have been no notable changes in the fisheries that would affect the potential for interactions with protected species, and there is no other information to indicate that impacts to protected species have changed in recent years in the Mariana Archipelago.

The climate change section of this report includes indicators of current and changing climate and related oceanic conditions in the geographic areas for which the Western Pacific Regional Fishery Management Council has responsibility. In developing this section, the Council relied on a number of recent reports conducted in the context of the U.S. National Climate Assessment including, most notably, the 2012 Pacific Islands Regional Climate Assessment and the Ocean and Coasts chapter of the 2014 report on a Pilot Indicator System prepared by the National Climate Assessment and Development Advisory Committee. The primary goal for selecting the indicators used in this report is to provide fisheries-related communities, resource managers, and businesses with climate-related situational awareness. In this context, indicators were selected to be fisheries relevant and informative, build intuition about current conditions in light of changing climate, provide historical context and recognize patterns and trends.

The atmospheric concentration of carbon dioxide (CO<sub>2</sub>) trend is increasing exponentially with the time series maximum at 409 ppm. The oceanic pH at Station Aloha, in Hawaii has shown a significant linear decrease of -0.0386 pH units, or roughly a 9.4% increase in acidity ([H<sup>+</sup>]) since 1989. Annual mean sea surface temperature (SST) was 28.26°C in 2018, and over the period of record, annual SST has increased at a rate of 0.022°C/year. The annual anomaly was 0.188 °C hotter than average, with intensification in the northern islands. Tropical cyclone activity was just above average in the Western North Pacific in 2018, with 29 storms, 13 typhoons, and 7

super typhoons. Super Typhoon Yutu notably made landfall on the islands of Tinian and Saipan as a Category 5 equivalent typhoon with estimated winds of 180 mph and a central minimum pressure of 905 mb, marking the second strongest tropical cyclone to impact any U.S. territory on record. The local trend in sea level rise is 3.68 millimeters/year based on monthly mean sea level data from 1993 to 2018 which is equivalent to a change of 1.21 feet in 100 years.

The Mariana Archipelago FEP and National Standard 2 guidelines require that this report include a report on the review of essential fish habitat (EFH) information. The 2018 annual report includes cumulative impacts on EFH as well as a review of relevant life history and habitat information for four common coral reef crustaceans. In 2017, descriptions of relevant life history and habitat information for four common coral reef crustaceans were provided. The guidelines also require a report on the condition of the habitat. In the 2018 annual report, mapping progress and benthic cover are included as indicators, pending development of habitat condition indicators for the Mariana Archipelago not otherwise represented in other sections of this report. The annual report addresses any Council directives toward its plan team, and there were no directives in 2018.

The marine planning section of this report records activities with multi-year planning horizons and begins to track the cumulative impact of established facilities. Development of the report in the future will focus on identifying appropriate data streams. Military activities in the Marianas continue to impact fisheries and their access. With the Records of Decision on the Mariana Islands Testing and Training and Guam and CNMI Military Relocation SEIS, access to fishing grounds will be impacted at Ritidian Point on Guam and at Farallon de Medinilla in CNMI during live-fire exercises. Nearshore water quality will be impacted in Northern Guam until the Northern District Wastewater Treatment Plant is upgraded. A re-release of the draft CNMI Joint Military Training Environmental Impact Statement (EIS) is expected sometime in early 2019. The 2019 Mariana Islands Training and Testing Final Supplemental EIS is expected in spring 2020. CNMI representatives have requested a military liaison to be present at their advisory panel committee meetings, especially to discuss the expired Memorandum of Understanding (MOU) for the Garapan Anchorage, and the Department of Defense will establish a coordinating council to discuss issues associated with increased military activity in the CNMI. A new MOU was not signed in 2018.

The data integration chapter of this report is still under development. The archipelagic data integration chapter explores the potential association between fishery parameters and ecologically-associated variables that may be able to explain a portion of the variance in fishery-dependent data. A contractor completed preliminary evaluations in 2017, and results of exploratory analyses were included for the first time in the 2017 Annual SAFE Report; however, suggested revisions during review by the Archipelagic Plan Team delayed updates to be implemented for the data integration section for the Mariana Archipelago. Results presented from the 2017 analyses showed that the coral reef fisheries in Guam and CNMI generally had a negative association with local precipitation levels. The reef fishery in Guam was shown to have a negative relationship with chlorophyll-*a* concentration. There was no relationship uncovered between akule and precipitation despite a general understanding of the connection between the two factors. All evaluated coral reef taxa in Guam showed a negative relationship with sea surface temperature in the region, while species in the Acanthurids (i.e., surgeonfish/unicornfish) had no notable relationship.



Multivariate analyses were employed in the data integration chapter to more completely evaluate the aggregate effects of ecological parameters on the selected fisheries. A non-metric multidimensional scaling analysis showed that, while presented evaluations were not able to identify any significant levels of association between expanded creel catch data and a range of environmental parameters, the first axis, responsible for explaining 91% of the variance, illustrated the strongest relationships with salinity (negative) and rainfall (positive).

Going forward with the data integration analyses and presentation of results for Chapter 3 of the Annual SAFE Reports, the Plan Team suggested several improvements to implement in the coming year: standardizing and correcting values in CPUE time series, incorporating longer stretches of phase lag, completing comparisons on the species-level and by dominant gear types, incorporating local knowledge on shifts in fishing dynamics over the course of the time series, and utilizing the exact environmental data sets presented in the ecosystem consideration chapter of the annual report. Many of these recommendations were applied to datasets from Hawaii in 2018, and will similarly be done for Mariana Archipelago data integration analyses in the upcoming report cycles. Implementation of these suggestions will allow for the preparation and publication of a more finalized version of the data integration chapter in coming years.

Regarding this 2018 Annual SAFE Report, the 2019 Archipelagic Fishery Ecosystem Plan Team recommends the Council:

- Direct staff to work with NMFS to convene the Plan Team working group for American Samoa, Guam, CNMI and Hawaii to define the ecosystem component species that will be monitored as species that comprise the functional groups (e.g., ‘parrotfish’, ‘browsing surgeon’, ‘mid-size targeted surgeon’, ‘medium large snappers’, ‘non-planktivorous butterflyfishes’), and those that comprise key species in the fisheries (i.e., top 5 consistently monitored important species and the 10 annual catch landings)
- Direct staff to work with NMFS and AS-DMWR, CNMI-DFW, Guam-DAWR, Hawaii-DAR on the revisions to the fisheries modules of the Archipelagic SAFE Reports due to the changes in the Management Unit Species brought about by the Ecosystem Component designation;
- Direct staff to work with NMFS-PIFSC-Ecosystem Science Division and Division of Aquatic Resources on applying the GLM framework to the survey data in order to validate the modeling results.

Additional work item recommendations included:

- Council staff and the Archipelagic Plan Team Chair to work with NMFS and AS-DMWR, CNMI-DFW, Guam-DAWR, Hawaii-DAR on determining which table(s) to remove from the Annual SAFE Report due to the ecosystem component amendment, etc.;
- WPacFIN to follow-up on the status of the creel survey method documentation;
- The report to incorporate more nuance in the narratives of the fishery performance sections; include the issue on pounds sold greater than pounds caught;

- The report to identify presence and absence of hi-liners in the data sets as well as define the criteria of what a hi-liner is;
- Regarding effort and participation metrics for the Annual SAFE Report, Council staff and PIFSC employees to calculate the average fishermen per trip and ensure interview has number of fishermen and average numbers of gear per trip.
- Add the abstracts for relevant data integration studies to Chapter 3; and
- “Cross-walk” tables with the information regularly needed to complete Environmental Assessments (EAs).

This page was intentionally left blank.

## TABLE OF CONTENT

|   |     |
|---|-----|
| Executive Summary .....   | v   |
| Acronyms and Abbreviations .....  | xxi |
| 1 Fishery Performance .....   | 25  |
| 1.1 CNMI Fishery Descriptions .....   | 25  |
| 1.1.1 Background .....  | 25  |
| 1.1.2 Fishery Data Collection System .....  | 28  |
| 1.1.3 Meta-Data Dashboard Statistics .....  | 32  |
| 1.1.4 Fishery Summary Dashboard Statistics .....                                    | 34  |
| 1.1.5 Catch Statistics .....  | 37  |
| 1.1.6 Catch-per-Unit-Effort (CPUE) Statistics .....                                 | 45  |
| 1.1.7 Effort Statistics .....   | 46  |
| 1.1.8 Participants .....  | 47  |
| 1.1.9 Bycatch Estimates .....   | 50  |
| 1.1.10 Number of Federal Permit Holders .....                                       | 52  |
| 1.1.11 Status Determination Criteria .....  | 53  |
| 1.1.12 Overfishing Limit, Acceptable Biological Catch, and Annual Catch Limits .... | 57  |
| 1.1.13 Best Scientific Information Available .....                                  | 59  |
| 1.1.14 Harvest Capacity and Extent .....  | 61  |
| 1.1.15 Administrative and Regulatory Actions .....                                  | 62  |
| 1.1.16 References .....   | 62  |
| 1.2 Guam Fishery Descriptions .....   | 64  |
| 1.2.1 Bottomfish Fishery .....  | 64  |
| 1.2.2 Coral Reef Fishery .....  | 66  |
| 1.2.3 Fishery Data Collection System .....  | 66  |
| 1.2.4 Meta-Data Dashboard Statistics .....  | 67  |
| 1.2.5 Creel Survey Meta-Data Statistics .....                                       | 68  |
| 1.2.6 Fishery Summary Dashboard Statistics .....                                    | 71  |
| 1.2.7 Catch statistics .....  | 74  |
| 1.2.8 Catch-per-Unit-Effort (CPUE) Statistics .....                                 | 85  |
| 1.2.9 Effort Statistics .....   | 87  |
| 1.2.10 Participants .....   | 90  |
| 1.2.11 Bycatch Estimates .....  | 95  |
| 1.2.12 Number of Federal Permit Holders .....                                       | 99  |
| 1.2.13 Status Determination Criteria .....  | 100 |
| 1.2.14 Overfishing Limit, Acceptable Biological Catch, and Annual Catch Limits ..   | 104 |
| 1.2.15 Best Scientific Information Available .....                                  | 106 |
| 1.2.16 Harvest Capacity and Extent .....  | 108 |
| 1.2.17 Other Relevant Ocean-Uses and Fishery-Related Information .....              | 109 |
| 1.2.18 Administrative and Regulatory Actions .....                                  | 109 |
| 1.2.19 References .....   | 109 |
| 2 Ecosystem Considerations .....  | 113 |
| 2.1 Coral Reef Ecosystem Parameters .....   | 113 |
| 2.1.1 Regional Reef Fish Biomass .....  | 113 |
| 2.1.2 CNMI Reef Fish Biomass .....  | 116 |
| 2.1.3 CNMI Archipelagic Mean Fish Size .....  | 119 |

|       |   |     |
|-------|---|-----|
| 2.1.4 | CNMI Reef Fish Population Estimates.....                                    | 122 |
| 2.1.5 | Guam Reef Fish Biomass .....  | 125 |
| 2.1.6 | Guam Archipelagic Mean Size.....  | 127 |
| 2.1.7 | Guam Reef Fish Population Estimates .....                                   | 129 |
| 2.2   | Life History and Length-Derived Parameters.....                             | 130 |
| 2.2.1 | CNMI Coral Reef Ecosystem – Reef Fish Life History .....                    | 130 |
| 2.2.2 | CNMI Bottomfish Ecosystem – Bottomfish Life History.....                    | 138 |
| 2.2.3 | Guam Coral Reef Ecosystem – Reef Fish Life History.....                     | 145 |
| 2.2.4 | Guam Bottomfish Ecosystem – Bottomfish Life History.....                    | 153 |
| 2.2.5 | References.....   | 159 |
| 2.3   | Socioeconomics .....  | 161 |
| 2.3.1 | Response to Previous Council Recommendations.....                           | 162 |
| 2.3.2 | CNMI.....   | 163 |
| 2.3.3 | Guam.....   | 174 |
| 2.3.4 | Ongoing Research and Information Collection .....                           | 187 |
| 2.3.5 | Relevant PIFSC Economics and Human Dimensions Publications: 2018 .....      | 189 |
| 2.3.6 | References.....   | 189 |
| 2.4   | Protected Species .....   | 192 |
| 2.4.1 | Indicators for Monitoring Protected Species Interaction.....                | 192 |
| 2.4.2 | FEP Conservation Measures.....  | 192 |
| 2.4.3 | Status of Protected Species Interactions in the Marianas FEP Fisheries..... | 195 |
| 2.4.4 | Identification of Emerging Issues .....                                     | 195 |
| 2.4.5 | Identification of Research, Data, and Assessment Needs.....                 | 197 |
| 2.5   | Climate and Oceanic Indicators.....   | 198 |
| 2.5.1 | Introduction.....   | 198 |
| 2.5.2 | Response to Previous Plan Team and Council Recommendations.....             | 199 |
| 2.5.3 | Conceptual Model.....   | 199 |
| 2.5.4 | Selected Indicators.....  | 201 |
| 2.5.5 | References.....   | 222 |
| 2.6   | Essential Fish Habitat .....  | 225 |
| 2.6.1 | Introduction.....   | 225 |
| 2.6.2 | Habitat Use by MUS and Trends in Habitat Condition .....                    | 227 |
| 2.6.3 | Report on Review of EFH Information .....                                   | 233 |
| 2.6.4 | EFH Levels .....  | 233 |
| 2.6.5 | Research and Information Needs.....   | 235 |
| 2.6.6 | References.....   | 236 |
| 2.7   | Marine Planning.....  | 238 |
| 2.7.1 | Introduction.....   | 238 |
| 2.7.2 | Response to Previous Council Recommendations.....                           | 238 |
| 2.7.3 | Marine Managed Areas Established under FEPs.....                            | 238 |
| 2.7.4 | Fishing Activities and Facilities .....                                     | 241 |
| 2.7.5 | Non-Fishing Activities and Facilities .....                                 | 241 |
| 2.7.6 | Pacific Islands Regional Planning Body Report.....                          | 241 |
| 2.7.7 | References.....   | 244 |
| 3     | Data Integration .....  | 246 |
| 3.1   | Introduction.....   | 246 |

3.1.1 Potential Indicators for Nearshore Fisheries..... 246

3.1.2 2018 Recommendations for Chapter Development..... 247

3.1.3 Past Work..... 248

3.2 Precipitation..... 248

3.2.1 Guam..... 248

3.3 Sea Surface Temperature..... 252

3.3.1 CNMI..... 254

3.3.2 Guam..... 258

3.4 Primary Productivity..... 261

3.4.1 CNMI..... 261

3.4.2 Guam..... 267

3.5 Multivariate Assessments of Other Ecosystem Variables ..... 271

3.5.1 Non-metric Multidimensional Scaling..... 271

3.6 References..... 274

Appendix A: Guam and CNMI FEP management unit species list.....A-1

Appendix B: List of Protected Species and Designated Critical Habitat.....B-1

TABLE OF TABLES

Table 1. Summary of CNMI creel survey meta-data ..... 33

Table 2. Summary of commercial receipt book meta-data ..... 34

Table 3. 2018 annual indicators for the coral reef and bottomfish fisheries describing performance relative to short-term (10-year) and long-term (20-year) averages ..... 35

Table 4. Summary of catch (lbs.) for all species caught by bottomfishing gear from 1983-2018 as recorded by creel surveys and the commercial purchase system..... 38

Table 5. Summary of BMUS catch (lbs.) for from 1983-2018 for creel surveys and the commercial purchase system ..... 39

Table 6. Summary of CREMUS catch (lbs.) from 1983-2018 for creel surveys and the commercial purchase system ..... 40

Table 7. Expanded CNMI creel survey catch estimates (lbs.) from 2005-2018..... 41

Table 8. Catch for top species complexes in CNMI boat-based creel data from 2000-2018 ..... 43

Table 9. Catch for top species complexes in CNMI shore-based creel data from 2000-2018 ..... 44

Table 10. CPUE for top fishing methods in CNMI shore-based fisheries from 2005-2018 ..... 45

Table 11. CPUE for top fishing methods in CNMI boat-based fisheries from 2000-2018 ..... 45

Table 12. Effort estimates in CNMI coral reef and bottomfish fisheries from 2000-2018 ..... 46

Table 13. Number of fishermen participating in the CNMI bottomfish fishery and number of gears in the CNMI coral reef fishery from 2000-2018 ..... 48

Table 14. Bycatch estimates in CNMI non-bottomfishing boat-based fisheries from 2000-2018 50

Table 15. Bycatch estimates in the CNMI bottomfish fishery from 2000-2018 ..... 51

Table 16. Bycatch estimates in the CNMI shore-based fishery from 2000-2018..... 51

Table 17. Number of federal permits holders for the CNMI crustacean and bottomfish fisheries from 2009-2018 ..... 53

Table 18. Overfishing threshold specifications for the BMUS in CNMI..... 53

Table 19. Rebuilding control rules for the BMUS in CNMI ..... 54

Table 20. Status determination criteria for CNMI CREMUS using CPUE based proxies..... 55

Table 21. Stock assessment parameters for the CNMI BMUS complex (Yau et al. 2015). ..... 56

Table 22. Best available MSY estimates for the coral reef MUS in CNMI ..... 57

Table 23. CNMI ACL table with three-year average catch in pounds ..... 58

Table 24. CNMI proportion of harvest capacity and extent for 2018..... 61

Table 25. Summary of Guam creel survey meta-data describing survey performance parameters with potential influence on the creel survey expansion from 1982-2018..... 68

Table 26. Summary of Guam commercial receipt book meta-data describing reporting performance parameters with potential influence on total commercial landing estimates from 1980-2018 ..... 70

Table 27. 2018 annual indicators for the coral reef and bottomfish fishery describing fishery performance comparing current estimates with short-term (10 year) and long-term (20 year) average values..... 71

Table 28. Summary of time series of catch (lbs.) for all species caught using the bottomfishing gear in Guam from 1980-2018..... 74

Table 29. Summary of time series of BMUS catch (lbs.) in Guam from 1980-2018..... 75

Table 30. Summary of time series of CREMUS catch (lbs.) in Guam from 1980-2018..... 77

Table 31. Summary of expanded creel survey time series of catch (lbs.) by gear type in Guam for 1982-2018 ..... 79

|   |     |
|---|-----|
| Table 32. Catch time series of 11 top CREMUS from Guam boat-based creel survey expansion data from 2000-2018.....   | 82  |
| Table 33. Catch time series of 11 top CREMUS from Guam shore-based creel survey expansion data from 1980-2018.....  | 83  |
| Table 34. CPUE time series for dominant fishing methods in Guam shore-based fisheries CPUE from 1984-2018 .....   | 85  |
| Table 35. CPUE time series for dominant fishing methods in Guam boat-based fisheries CPUE from 1982-2018 .....  | 86  |
| Table 36. Time series of effort estimates (gear hours or no. trips for bottomfish) from Guam coral reef and bottomfish fisheries for 1982-2018.....                   | 88  |
| Table 37. Number of boats participating in the Guam bottomfish fishery and number of gears in the Guam boat- and shore-based coral reef fisheries from 1982-2018..... | 91  |
| Table 38. Time series of bycatch estimates in Guam boat-based non-bottomfishing fisheries from 1982-2018 .....  | 96  |
| Table 39. Time series of bycatch estimates in the Guam bottomfish fishery from 1982-2018....  | 97  |
| Table 40. Time series of bycatch estimates in the Guam shore-based fishery from 1984-2018 ..  | 98  |
| Table 41. Number of federal permits holders for the crustacean and bottomfish fisheries of Guam from 2009-2018 .....  | 100 |
| Table 42. Overfishing threshold specifications for Guam BMUS.....   | 100 |
| Table 43. Rebuilding control rules for Guam BMUS.....   | 101 |
| Table 44. Status determination criteria for the coral reef management unit species using CPUE based proxies.....  | 102 |
| Table 45. Stock assessment parameters for the Guam BMUS complex.....  | 103 |
| Table 46. Best available MSY estimates for the coral reef MUS in Guam .....   | 104 |
| Table 47. Guam ACL table with three-year average catch (lbs.) .....   | 105 |
| Table 48. Guam 2018 ACL proportion of harvest extent and the harvest capacity.....  | 108 |
| Table 49. Reef fish estimates for CNMI CREMUS in 0-30 m hard bottom habitat.....  | 122 |
| Table 50. Reef fish population estimates for Guam CREMUS .....  | 129 |
| Table 51. Available age, growth, and reproductive maturity information for coral reef species targeted for otolith and gonad sampling in CNMI.....                    | 133 |
| Table 52. Available length-derived information for various coral reef species in CNMI.....  | 135 |
| Table 53. Available age, growth, and reproductive maturity information for bottomfish species targeted for otoliths and gonads sampling in CNMI.....                  | 141 |
| Table 54. Available length-derived information for various bottomfish species in CNMI.....  | 143 |
| Table 55. Available age, growth, and reproductive maturity information for coral reef species targeted for otoliths and gonads sampling in Guam .....                 | 147 |
| Table 56. Available length derived information for various coral reef species in Guam.....  | 150 |
| Table 57. Available age, growth, and reproductive maturity information for bottomfish species targeted for life history sampling (otoliths and gonads) in Guam .....  | 156 |
| Table 58. Available length derived information for various bottomfish species in Guam.....  | 158 |
| Table 59. Commercial landings and revenue information of CNMI bottomfish fishery, 2009-2018.....  | 168 |
| Table 60. Average trip costs for CNMI bottomfish trips, 2009–2017.....  | 170 |
| Table 61. Commercial landings and revenue information of CNMI coral reef fishery, 2009-2018 .....   | 172 |
| Table 62. Average costs for CNMI bottomfish trips, 2009–2017 .....  | 174 |



|  |     |
|--|-----|
| Table 63. Commercial landings, revenue, and price information of Guam bottomfish fishery, 2009-2018 .....  | 182 |
| Table 64. Average trip costs for Guam bottomfish fishing trips from 2009–2018.....   | 183 |
| Table 65. Commercial landings and revenue information of Guam coral reef fish fishery, 2009-2018.....  | 186 |
| Table 66. Average trip costs for Guam spear/snorkel fish trips from 2011–2018 (adjusted to 2018 dollars) .....   | 187 |
| Table 67. Pacific Islands Region 2018 Commercial Fishing Economic Assessment Index .....   | 188 |
| Table 68. Pacific Islands Region 2019 Commercial Fishing Economic Assessment Index .....   | 188 |
| Table 69. Summary of ESA consultations for Mariana Archipelago FEP Fisheries.....  | 192 |
| Table 70. Candidate ESA species, and ESA-listed species being evaluated for critical habitat designation .....   | 195 |
| Table 71. Summary of habitat mapping in CNMI .....   | 229 |
| Table 72. Mean percent cover of live coral from RAMP sites collected from towed-diver surveys in the Mariana Archipelago .....   | 231 |
| Table 73. Mean percent cover of macroalgae from RAMP sites collected from towed-diver surveys in the Mariana Archipelago .....   | 231 |
| Table 74. Mean percent cover of crustose coralline algae from RAMP sites collected from towed-diver surveys in the Mariana Archipelago .....   | 232 |
| Table 75. Level of EFH information available for the Western Pacific PCMUS.....  | 233 |
| Table 76. Level of EFH information available for Western Pacific BMUS and seamount groundfish MUS complexes .....  | 234 |
| Table 77. Level of EFH information available for the Western Pacific crustacean MUS.....   | 235 |
| Table 78. MMAs established under FEPs from 50 CFR § 665 .....  | 240 |
| Table 79. Department of Defense major planning activities .....  | 242 |
| Table 80. Families in creel surveys from the U.S. Western Pacific analyzed in this report.....   | 255 |
| Table 81. Correlation coefficients ( <i>r</i> ) between recreational coral reef fishery catch (kg) and SST (°C) in the CNMI for 12 top taxa harvested from 2000-2016 .....   | 255 |
| Table 82. Correlation coefficients ( <i>r</i> ) between recreational coral reef fishery catch (in kg) and SST (°C) in Guam for 12 top taxa harvested from 1985-2016.....   | 259 |
| Table 83. Correlation coefficients ( <i>r</i> ) from comparisons of time series of the CNMI recreational coral reef fishery annual catch (kg) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) from 2000-2014 .....  | 264 |
| Table 84. Correlation coefficients ( <i>r</i> ) from comparisons of time series of for shore-and boat-based creel survey records in Guam (kg) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) for 12 top taxa harvested from 1998 - 2014. Significant correlations are indicated in bold ( $\alpha=0.05$ )..... | 268 |

## TABLE OF FIGURES

|  |     |
|--|-----|
| Figure 1. Mean fish biomass ( $\text{g}/\text{m}^2 \pm$ standard error) of CREMUS from 2010-2018 by latitude; figure continued on next page.....   | 114 |
| Figure 2. Mean fish biomass ( $\text{g}/\text{m}^2 \pm$ standard error) of CREMUS from 2010-2018 by latitude with mean estimates overlaid in red; figure continued on next page.....   | 117 |
| Figure 3. Mean fish size (cm, TL $\pm$ standard error) of CNMI CREMUS from 2010-2018 by latitude with mean estimates overlaid in red; figure continued on next page.....   | 120 |
| Figure 4. Mean fish biomass ( $\text{g}/\text{m}^2 \pm$ standard error) of Guam CREMUS from the years 2010-2018 with mean estimates overlaid in red.....   | 126 |
| Figure 5. Mean fish size (cm, TL $\pm$ standard error) of Guam CREMUS from 2010-2018 with mean estimates overlaid in red.....  | 128 |
| Figure 6. Average length over time of representative CNMI CREMUS from the BioSampling Program; figure continues for various species onto the next two pages.....   | 136 |
| Figure 7. Average length over time of CNMI BMUS derived from the BioSampling Program; continued from previous page.....  | 145 |
| Figure 8. Average length over time of representative Guam CREMUS derived from the BioSampling Program; continued from previous two pages.....  | 153 |
| Figure 9. Average length over time of representative Guam BMUS derived from the BioSampling Program.....   | 159 |
| Figure 10. Settlement of the Pacific Islands, courtesy of Wikimedia Commons, <a href="https://commons.wikimedia.org/wiki/File:Polynesian_Migration.svg">https://commons.wikimedia.org/wiki/File:Polynesian_Migration.svg</a> ..... | 161 |
| Figure 11. The commercial landings and revenues of BMUS, for the CNMI bottomfish fishery, 2009-2018 (Adjusted to 2018 dollars).....  | 167 |
| Figure 12. The prices of BMUS for the CNMI bottomfish fishery, 2009-2018.....  | 168 |
| Figure 13. Average costs for CNMI bottomfish trips, 2009–2017 (adjusted to 2017 dollars)...  | 169 |
| Figure 14. The pounds sold and revenues for the CNMI reef fish fishery, 2009-2018 (adjusted to 2018 dollars).....  | 171 |
| Figure 15. The fish prices of CREMUS for the CNMI reef fishery, 2009-2018.....   | 172 |
| Figure 16. Average costs for CNMI spearfishing trips, 2009–2017 (adjusted to 2017 dollars). 173  |     |
| Figure 17. The pounds sold and revenues for the Guam bottomfish fishery, 2009-2018 (adjusted to 2018 dollars).....   | 181 |
| Figure 18. The prices of BMUS for the Guam bottomfish fishery, 2009-2018.....  | 181 |
| Figure 19. Average trip costs for Guam bottomfish fishing trips from 2009–2018 (adjusted to 2018 dollars).....   | 183 |
| Figure 20. The commercial landings and revenues, for the Guam reef fish fishery, 2009-2018   | 185 |
| Figure 21. The prices of CREMUS for the Guam reef fish fishery, 2009-2018.....   | 185 |
| Figure 22. Average trip costs for Guam spear/snorkel trips from 2011–2018 (adjusted to 2018 dollars).....  | 186 |
| Figure 23. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability.....  | 200 |
| Figure 24. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability.....  | 201 |
| Figure 25 Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of anthropogenic climate change.....  | 202 |
| Figure 26. Regional spatial grids representing the scale of the climate change indicators being monitored.....   | 202 |

|  |     |
|--|-----|
| Figure 27. Monthly mean (red) and seasonally-corrected (black) atmospheric carbon dioxide at Mauna Loa Observatory, Hawaii .....   | 204 |
| Figure 28. Oceanic pH (black) and its trend (red) at Station ALOHA from 1989 – 2016 .....  | 205 |
| Figure 29. Oceanic Niño Index from 1950-2018 (top) and 2000–2018 (bottom) with El Niño periods in red and La Niña periods in blue .....  | 206 |
| Figure 30. Pacific Decadal Oscillation from 1950–2018 (top) and 2000–2018 (bottom) with positive warm periods in red and negative cool periods in blue.....  | 207 |
| Figure 31. 2018 Pacific basin tropical cyclone tracks .....  | 209 |
| Figure 32. 2018 tropical storm totals by region.....   | 210 |
| Figure 33. 2018 Accumulated Cyclone Energy (ACE) Index by region.....  | 210 |
| Figure 34. Sea surface temperature climatology and anomalies from 1982-2018 .....  | 212 |
| Figure 35. Coral Thermal Stress Exposure measured at CNMI Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks) .....  | 214 |
| Figure 36. Chlorophyll-A (Chl-A) and Chl-A Anomaly from 1982-2018 .....  | 216 |
| Figure 37. CMAP precipitation across the Marianas Grid with 2018 values in red.....  | 218 |
| Figure 38a. Sea surface height and anomaly .....   | 219 |
| Figure 39. Monthly mean sea level without regular seasonal variability due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents.....  | 221 |
| Figure 40. Substrate EFH Limit of 700 m isobath around the Mariana Archipelago (from Ryan et al. 2009) .....   | 228 |
| Figure 41. CNMI Land and Seafloor Area and Primary Data Coverage .....   | 230 |
| Figure 42. Regulated fishing areas of the Mariana Archipelago .....  | 239 |
| Figure 43. Total annual precipitation (in.) in Guam from 1980-2016.....  | 249 |
| Figure 44. Time series of total annual estimated (i.e. expanded) landings of atulai in kilograms from Guam shore-and boat-based creel survey records from 1982-2016.....   | 250 |
| Figure 45. Time series of total sampled (left) and expanded (right) effort for top gear types in shore-based (top) and boat-based (bottom) creel surveys in Guam from 1982-2016.....   | 251 |
| Figure 46. Linear regression between total atulai catch (kg) in the Guam shore-based and boat-based creel survey records and total annual rainfall (in.) from 1982-2016 .....  | 251 |
| Figure 47. Time series of total annual expanded landings of <i>Decapterus macarellus</i> (kg) in Guam shore-and boat-based creel survey records from 1982-2016 .....   | 252 |
| Figure 48. Time series of SST (°C) in the CNMI from 1985-2016 (CV = 0.55) .....  | 253 |
| Figure 49. Time series of SST (°C) from 1985-2016 in Guam (CV = 1.38).....   | 254 |
| Figure 50. Time series of total annual catch (kg; blue) for the CNMI recreational coral reef fishery plotted alongside average annual SST (°C; black) from 2000-2016 .....   | 255 |
| Figure 51. Linear regression showing the correlation between total annual catch (kg) in creel survey records and average annual SST (°C) in the CNMI from 2000-2016.....   | 256 |
| Figure 52. Linear regressions showing the three top correlations between total annual catch (kg) from creel survey records and average annual SST (°C) in the CNMI from for (a) Mullids, (b) Carangids, and (c) Lutjanids from 2000–2016 ..... | 257 |
| Figure 53. Time series of total annual catch (kg; blue) in the Guam shore-and boat-based creel survey records plotted with average annual SST (°C; black) from 1985-2016.....  | 258 |
| Figure 54. Linear regression between total annual catch (kg) for shore- and boat-based creel survey records and average annual SST (°C) in Guam from 1985-2016 .....   | 259 |

|   |     |
|---|-----|
| Figure 55. Linear regressions showing three top correlations between total annual catch (kg) for shore-and boat-based creel survey records and average annual SST (°C) in Guam for (a) Lutjanids, (b) Siganids, and (c) Mugilids from 1985–2016.....  | 260 |
| Figure 56. Time series of fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) around the CNMI from 1998-2016 (CV=6.28).....  | 262 |
| Figure 57. Time series of fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) around Guam from 1998-2016 (CV=7.03).....  | 263 |
| Figure 58. Comparison of the CNMI recreational reef fish catch (kg; black) from creel survey records with two years of time lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ; blue) from 2000-2014 ( $r = 0.32$ ) .....  | 264 |
| Figure 59. Linear regression between total annual catch (kg) phase lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) in CNMI from 2000-2014.....   | 265 |
| Figure 60. Linear regressions showing the three top correlations between total annual catch (kg) for the CNMI from creel survey records with phase lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) for (a) Holocentrids, (b) Serranids, and (c) Acanthurids from 2000–2014 .....           | 266 |
| Figure 61. Comparison of Guam recreational reef fish catch for shore-and boat-based creel survey records (kg; black) with two years of time lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ; blue) from 1998-2014.....  | 267 |
| Figure 62. Linear regression between total annual catch (kg) for Guam shore-and boat-based creel survey records with phase lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) from 1998-2014.....   | 268 |
| Figure 63. Linear regressions showing the three top correlations between total annual catch (kg) for Guam for shore-and boat-based creel survey records with phase lag (t+2 years) and fluorometric chlorophyll- <i>a</i> concentrations (mg/m <sup>3</sup> ) for (a) Balistidae, (b) Mugilidae, and (c) Mullidae from 1998–2014..... | 270 |
| Figure 64. NMS scree plot showing the stress test to determine dimensionality for the final solution for the CNMI multivariate analysis .....   | 272 |
| Figure 65. One-dimensional scatterplot overlaid with a joint biplot depicting ordination scores resulting from an NMS analysis on creel survey expanded catch data and prominent environmental parameters in the CNMI from 2000-2014.....   | 273 |
| Figure 66. NMS scree plot showing the stress test to determine dimensionality for the final solution for the Guam multivariate analysis; two-axis solution was recommended.....   | 273 |
| Figure 67. Two-dimensional scatterplot overlaid with a joint biplot depicting ordination scores resulting from an NMS analysis on creel survey expanded catch data and prominent environmental parameters in Guam from 1989-2014.....   | 274 |

## ACRONYMS AND ABBREVIATIONS

| Acronym           | Meaning  |
|-------------------|--|
| ABC               | Acceptable Biological Catch  |
| ACE               | Accumulated Cyclone Energy   |
| ACL               | Annual Catch Limits  |
| ACT               | Annual Catch Target  |
| AM                | Accountability Measures  |
| AVHRR             | Advanced Very High Resolution Radiometer                               |
| BAC-MSY           | Biomass Augmented Catch MSY  |
| B <sub>FLAG</sub> | warning reference point for biomass                                    |
| BiOp              | Biological Opinion   |
| BMUS              | Bottomfish Management Unit Species                                     |
| BOEM              | Bureau of Ocean Energy Management                                      |
| BSIA              | Best Scientific Information Available                                  |
| CFR               | Code of Federal Regulations  |
| CMLS              | Commercial Marine License System                                       |
| CMS               | coastal and marine spatial   |
| CMUS              | Crustacean Management Unit Species                                     |
| CNMI              | Commonwealth of the Northern Mariana Islands                           |
| CPUE              | Catch per Unit Effort  |
| CRED              | Coral Reef Ecosystem Division  |
| CREMUS            | Coral Reef Ecosystem Management Unit Species                           |
| DAWR              | Division of Aquatic and Wildlife Resources                             |
| DGI               | Daily Growth Increments  |
| DLNR-DAR          | Department of Land and Natural Resources-Division of Aquatic Resources |
| DLNR-DFW          | Department of Land and Natural Resources-Division of Fish and Wildlife |
| DPS               | Distinct Population Segment  |
| EEZ               | Exclusive Economic Zone  |
| EFH               | Essential Fish Habitat   |
| EIS               | Environmental Impact Statement   |
| EKE               | Eddy kinetic energy  |
| ENSO              | El Niño Southern Oscillation   |
| EO                | Executive Order  |
| ESA               | Endangered Species Act   |
| FEP               | Fishery Ecosystem Plan   |
| FMP               | Fishery Management Plan  |
| FRS               | Fishing Report System  |
| GAC               | Global Area Coverage   |
| GFS               | global forecast system   |
| HAPC              | Habitat Area of Particular Concern                                     |
| HDAR              | Hawaii Division of Aquatic Resources                                   |
| IBTrACS           | International Best Track Archive for Climate Stewardship               |

|             |   |
|-------------|---|
| LOF         | List of Fisheries   |
| LVPA        | Large Vessel Prohibited Area                                    |
| MFMT        | Maximum Fishing Mortality Threshold                             |
| MHI         | Main Hawaiian Island  |
| MMA         | marine managed area   |
| MMPA        | Marine Mammal Protection Act                                    |
| MPA         | marine protected area   |
| MPCC        | Marine Planning and Climate Change                              |
| MPCCC       | Council's MPCC Committee  |
| MSA         | Magnuson-Stevens Fishery Conservation and Management Act        |
| MSFCMA      | Magnuson-Stevens Fishery Conservation and Management Act        |
| MSST        | Minimum Stock Size Threshold                                    |
| MSY         | Maximum Sustainable Yield                                       |
| MUS         | management unit species   |
| NCADAC      | National Climate Assessment & Development Advisory<br>Committee |
| NCDC        | National Climatic Data Center                                   |
| NEPA        | National Environmental and Policy Act                           |
| NESDIS      | National Environmental Satellite, Data, and Information Service |
| NMFS        | National Marine Fisheries Service                               |
| NMFS        | National Marine Fisheries Service                               |
| NOAA        | National Oceanic and Atmospheric Administration                 |
| NWHI        | Northwestern Hawaiian Islands                                   |
| OFL         | Overfishing Limits  |
| OFR         | Online Fishing Report   |
| ONI         | Ocean Niño Index  |
| OR&R        | Office of Response and Restoration                              |
| OY          | Optimum Yield   |
| PacIOOS     | Pacific Integrated Ocean Observing System                       |
| PCMUS       | Precious Coral Management Unit Species                          |
| Pelagic FEP | Fishery Ecosystem Plan for the Pacific Pelagic Fisheries        |
| PI          | Pacific Islands   |
| PIBHMC      | Pacific Island Benthic Habitat Mapping Center                   |
| PIFSC       | Pacific Island Fisheries Science Center                         |
| PIRCA       | Pacific Islands Regional Climate Assessment                     |
| PIRO        | NOAA NMFS Pacific Islands Regional Office                       |
| PMUS        | pelagic management unit species                                 |
| POES        | Polar Operational Environmental Satellite                       |
| PRIA        | Pacific Remote Island Areas                                     |
| RAMP        | Reef Assessment and Monitoring Program                          |
| RPB         | Regional Planning Body  |
| SAFE        | Stock Assessment and Fishery Evaluation                         |
| SBRM        | Standardized Bycatch Reporting Methodologies                    |
| SDC         | Status Determination Criteria                                   |
| SEEM        | Social, Economic, Ecological, Management uncertainties          |
| SPC         | Stationary Point Count  |

|         |   |
|---------|---|
| SST     | Sea Surface Temperature                             |
| TAC     | Total Allowable Catch                               |
| USACE   | United States Army Corps of Engineers               |
| WPacFIN | Western Pacific Fishery Information Network         |
| WPRFMC  | Western Pacific Regional Fishery Management Council |
| WPSAR   | Western Pacific Stock Assessment Review             |
| WW3     | Wave Watch 3  |

This page was intentionally left blank.



# 1 FISHERY PERFORMANCE

## 1.1 CNMI FISHERY DESCRIPTIONS

### 1.1.1 Background

The Commonwealth of the Northern Mariana Islands (CNMI) is a chain of islands in the Western Pacific Ocean. Along with the island of Guam, the chain is historically known as the Mariana Islands. The CNMI consists of 14 small islands situated in a north-south direction, stretching a distance of about 500 km. The surrounding waters of the CNMI play an integral role in the everyday lives of its citizens. The ocean is a major source of food and leisure activities for residents and tourists alike. Archeological research has also revealed evidence of fishing activities in the CNMI dating back 3,000 years. Although the composition of fishing activities in the Marianas has changed significantly since then, a common view of its importance remains.

#### **Fisheries during the German occupation**

During the German occupational period (1899-1914) a majority of the economic focus in the Northern Marianas was on the copra industry. Few commercial fisheries were noted during this period of time, as the German administration focused efforts on crop production and feral cattle trade (Russell 1999). Chamorro and Carolinians utilized the protected lagoon and open waters with several fishing methods: talaya (cast net), chinchulu (surround net), gigao (fish weir), tokcha (spear), tupak (hook and line), and Carolinians additionally gleaned sea cucumbers for the Asian Markets. Most of these activities were for subsistence purposes, with the catch being distributed and bartered among relatives and acquaintances.

#### **Fisheries during the Japanese occupation**

Fisheries development prospered during the Japanese administration (1914-1945), becoming the nation's second largest industry. Small pelagic fishing operations were established and the Garapan port became the main area for drying fish. Large scale fishing activities occurred during the 1930s, shown as Saipan produced 11 percent of total tuna landed in Micronesia (Bowers 2001). However, efforts to develop the tuna fishery shifted to Palau and Federated States of Micronesia (FSM) due to the availability of bait fish in the region. Subsistence fishing still persisted within the lagoon and fringing reefs, and was mainly conducted by the natives though a large extraction of sea cucumbers did occur. There were several main fishing methods used during this period: cast net, spear, gill net, surround net, hook and line, and gleaning. During this period, the topshell (*Trochus niloticus*) was also introduced into the Marianas.

#### **Fisheries during the U.S. military occupation**

The fishing industry was destroyed during World War 2, but quickly rebuilt afterwards with support from the U.S. military. Okinawans who operated the fishery prior to the war were hired to operate and train locals to fish commercially, targeting pelagic species. A company called Saipan Fishing Company operated during this time, and contributed to the early re-development of post-war commercial fisheries in the CNMI (Bowers 2001). Most of the fishing activities were for *Katsuwonus pelamis* (bonito) and other tuna species. However, other resources, such as big-eye scad, reef fish, and lobster, were also harvested during calm weather. The Chamorro and Carolinians continued subsistence fishing in the lagoon after the war. Although limited quantities of monofilament nets were available during this period, they were used to capture lagoon fish

and along the reef lines. The use of modern fishing gear such as masks, rubber fins, and flash lights made it much easier to harvest coral reef resources during this time.

### **Fisheries activities within the past two decades**

The CNMI has had numerous changes in its fisheries over the past twenty years. In the mid-1990s, commercial fishing activities increased significantly. Commercial SCUBA fishing became a common method, not only to support local demand for reef fish, but to bolster exports to Guam as well. Large-scale commercial bottomfishing in the Northern Islands of the CNMI peaked starting in the mid-1990s through 2002, with landings being both sold locally and exported to Japan. Troll fishing continued to be dominant during this period. An exploratory, deepwater shrimp fishery also developed, but did not last due to internal company issues and gear losses. Around this time, a sea cucumber fishery also began on Rota before migrating to Saipan; ultimately, however, this fishery was found to be unstable and was subsequently halted.

Several fishing companies entered the fisheries only to close down a few years later. The CNMI reached its highest population during the last two decades, most of whom have been migrant workers from Asia. The tourism industry has also been increasing, which contributes to high demand for fresh fish. Subsistence fishing within the nearshore waters of Saipan, Tinian, and Rota has also increased.

In the 2000s, small-scale troll, bottom and reef fish fisheries persisted, with landings sold locally. Federal and state support was provided multiple times to further develop fisheries in the CNMI with intermittent success. An exploratory longline fishery was funded and operated in the CNMI in the mid-2000 for about two years, but eventually closed down due to low productivity of high-value, pelagic fish, among other issues within the business. A few larger (40-80') bottomfishing vessels were also operational during this period, with a majority of them fishing the northern islands and offshore banks. A few of these vessels were recipients of financial assistance to improve their fishing capacities.

Fisheries in the CNMI have generally been relatively small and fluid, with 16-20' boats fishing within 20 miles from Saipan. Many of these small vessels conduct multiple fishing activities during a single trip. For example, a company that is supported mainly by troll fishing may also conduct bottomfishing and spearfishing to supplement their income. Fishing businesses tend to enter and exit the fishery when it is economically beneficial to do so, as they are highly sensitive to changes in the economy, development, population, and regulations. Subsistence fishing continues; however, fishing methods and target species have shifted in step with population demographics and fishery restrictions. Nearshore hook and line, cast net, and spear fishing are common activities, but fishing methods such as gill net, surround net, drag net, and SCUBA-spear have been restricted or outright banned in the CNMI since the early 2000s.

#### **1.1.1.1 Bottomfish Fishery**

The bottomfish fishery has not changed much from its early years in certain aspects. Relatively small (<25ft) fishing vessels are still being used to access bottom fishing grounds around Saipan and Tinian, while the larger (>25ft) vessels are used to access bottomfish resources in the Northern Islands. Only a handful of these larger bottom fishing vessels are operating within the CNMI. Most of the small bottomfishing vessels are owned by vendors; there are, however, a few subsistence bottomfishers that participate in the fishery intermittently. More recently, improved

technologies, such as sophisticated electronics to locate fish and various types of reels replacing handlines, have entered the CNMI bottomfish fishery.

Two distinct types of bottomfish fisheries are identified in the CNMI: shallow-water bottom fishing, which targets fish at depths down to 150 m, and deepwater bottom fishing, which targets fish at depths greater than 150 m. Species targeted by the shallow-water fishery consist of the Redgill Emperor (*Lethrinus rubrioperculatus*), Black Jack (*Caranx lugubris*), Matai (*Epinephelus fasciatus*), Sas (*Lutjanus kasmira*), and Lunartail Grouper (*Variola louti*), among other fish residing at similar depths. Species targeted by the deepwater bottom fishing depths (>150m) include onaga (*Etelis corsucans*), ehu (*E. carbunculus*), yellowtail kalekale (*Pristipomiodes auricilla*), amberjack (*Seriola dumerili*), blueline gindai (*P. argyrogrammicus*), gindai (*P. zonatus*), opakapaka (*P. filamentosus*), and eightbanded grouper (*Hyporthodus octofasciatus*), among other fish residing at similar depths.

Bottomfish Management Unit Species (BMUS) are not the only species caught in the shallow-bottom fishery. Coral Reef Ecosystem Management Unit Species (CREMUS) are also caught in the shallow-bottom fishery because of their close proximity to reefs. These fish are caught with various hook and line gears, including homemade handlining gear, rod and reel, and electric reels. Deep-water bottomfishing requires more efficient fishing gears, such as hydraulic reels. Bottomfishing trips generally occur during the day, but fishing trips to the Northern Islands can take two to four days depending on vessel size and refrigeration capacity. These trips are most productive during calm weather months. Successful fishermen targeting deep-water bottomfish tend to fish for one to four years before leaving the fishery, whereas the majority of fishermen targeting shallow-water bottomfish tend to leave the fishery after the first year.

The overall participation of fishermen in the bottomfish fishery tends to occur on a relatively short-term basis (i.e., less than four years). The slight difference between shallow-water and deep-water fishermen likely reflects the greater skill and investment required to participate in the deepwater bottomfish fishery. In addition, deepwater bottomfishing tends to include larger ventures that are more buffered from the impulses of individual choice, and are usually dependent on a skilled captain and fishermen. Overall, the long-term commitment to hard work, maintenance and repairs, and staff retention appear to be challenging for CNMI bottomfish fishermen to sustain their efforts for more than a few years. A full list of BMUS species is provided in Appendix A.

### 1.1.1.2 Coral Reef Fishery

Coral reef fisheries have been generally steady in recent years relative to previous decades. Small-scale nearshore fisheries in the CNMI continue to be important socially, culturally, recreationally, financially, and for subsistence. Most fishermen are subsistence fishers, with a number of them selling a portion of their catch to roadside vendors and some of these vendors employing the fishermen to maintain a constant supply of reef fish. Most of the fishing for coral reef species occurs within the Saipan lagoon and fringing reefs around the islands, targeting mainly finfish and invertebrates. All reef fish catches are sold to local markets or used for personal consumption with a minimal portion exported for off-island residents. Shoreline access is the most common way to harvest coral reef resources. Vessels are generally used during calm weather to fish areas not as accessible other times of the year, with fishing trips to other islands

being made when the weather is favorable. Fishing methods have not changed significantly compared to previous years; hook and line, cast netting, spear fishing, and gleaning are methods still being used today. Some of the common families found in the CNMI reef fish markets are Acanthuridae (surgeonfish), scoridae (parrotfish), multiday (goatfish), serenade (grouper), abridge (wrasse), holocentridae (soldier/squirrelfish), carangidae (jacks), scombridae (scad), haemulidae (sweetlips), gerridae (mojarra), kyphosidae (rudderfish), and mugilidae (mullet), as well as other non-fish families. A full list of CREMUS species is provided in Appendix A.

### **1.1.2 Fishery Data Collection System**

A majority of the information collected by the CNMI Division of Fish and Wildlife (DFW) is fishery-dependent. Since the early-1980s, attempts were made to establish a data collection program for the nearshore fisheries, but failed due to intergovernmental issues. Over recent decades, significant time and effort has been made to further develop nearshore fishery data collection. This effort has resulted in the re-establishment of the shore-based creel survey program by DFW in collaboration with other local and federal agencies.

#### **1.1.2.1 Creel Surveys**

Currently the CNMI maintains both a boat- and shore-based creel survey for the island of Saipan, with plans for expansion to the populated neighboring islands. The programs were established in 2000 and 2005, respectively, in order to strengthen the capacity of DFW in providing sufficient information to the public regarding local fisheries. Other programs, such as the invoicing system and importation monitoring, provide supplemental information on harvest and demand for the fishery.

Effective management of Saipan's marine fishery resources requires the collection of fishing effort, methods used, and harvest. The CNMI boat- and shore-based creel surveys are some of the major data collection systems used by DFW to estimate the total annual boat-based participation, effort, and harvest while surveying nearshore fishery resources. These surveys were formerly known as the "CNMI offshore and inshore creel survey", but are now referred to as "boat- or shore-based" because they cover all fishing done from a boat or from shore. This is an important distinction because where the fishing activity is initiated (i.e. boat or shore) determines how that type of activity will be accounted for in the survey systems. For instance, very small boats launched from non-standard launching areas (e.g. from the back of a pickup truck on a beach) are not included in the boat-based creel survey.

The objective of the boat-based creel survey program is to quantify fishing participation, effort, and catch done from on a vessel in CNMI's waters. DFW had an early creel survey data collection program in 1984, and 1990 to 1994, however since the methods were not standardized, the data collected with that early program is not currently being used. The early program was eventually terminated due to a lack of resources. On April 2, 2000, the DFW fishery staff reinitiated the boat-based creel survey program on the island's boat-based fishery following a three year hiatus. The fishery survey collects data on the island's boating activities and interviews returning commercial and noncommercial fishermen at the three most active launching ramps/docks on the island: Smiling Cove, Sugar Dock, and Fishing Base. Essential fishery information is collected and processed from both commercial and noncommercial vessels to help

better inform management decisions. The two types of data collection programs utilized by Saipan's boat-based creel survey program include: boat-based participation count to collect participation data, and a boat-based access point survey to collect catch and effort data (through survey maps, boat logs, and interviews) at the three major boat ramp areas listed above. The data collected are then expanded at a stratum level (quarterly vs. annually, charter vs. non-charter, weekday vs. weekend, etc.) to create estimated landings by gear type for CNMI's boat-based fishery. The shore-based survey currently covers the Western Lagoon of Saipan. Some pilot surveys are being conducted on Saipan's Eastern beaches such as Laolao Bay, Obyan Beach, and Ladder Beach. Other accessible areas are not covered at this time due to existing limited resource availability and logistical constraints. With the assistance of the WPacFIN program at PIFSC, data processing software and a database were developed to process these survey data.

In May 2005, DFW fishery staff reinitiated the creel survey program for the island's shore-based fishery following a hiatus of 11 years. The Western Lagoon starts from the northwest (Wing Beach) and extends to the southwest (Agingan Point) of Saipan. This encompasses over twenty accessible and highly active shoreline access points. Saipan's shore-based creel survey is also a stratified randomized data collection program. This program collects two types of data to estimate catch and effort information in the shore-based fishery: participation count and interview. The participation count involves counting the number of people fishing on randomly selected days and their method of fishing along the shoreline. The interview involves dialoging with fishermen to determine catch, method used, length and weights of fish, species composition, catch disposition, and if any fish were not kept (i.e., bycatch). The data collected from this program have been used to expand and create annual estimated landings for the shore-based fishery in the CNMI.

In October 2018, the islands of Saipan and Tinian were directly hit by Super Typhoon Yutu. The damage inflicted by the typhoon delayed both creel surveys and collection of commercial receipt invoices. About a month after the typhoon, creel surveys were regularly conducted again, and boat-based surveys followed soon thereafter. Vendors prioritized repairing typhoon-related damages to their businesses, and the number of invoices collected decreased as a result.

From January to June 2018, there were 36 boat-based surveys scheduled. A total of 63 interviews were completed with an expanded catch estimate of 243,259 lbs. landed. The vessel/trailer participation survey is ongoing and includes all launching areas on the west coast of Saipan where all boat-based fishing occurs. For this reporting period, a total of 122 boat vessels/trailers were registered as "out fishing". During this progress period, the most common fishing methods encountered were trolling, bottomfishing, and hook and line fishing. The expanded harvest estimate for trolling was 151,270 lbs. Estimated catch for bottomfishing and hook and line gear types were 83,246 lbs. and 8,743 lbs., respectively.

In the second half of the year, from July to December 2018, there were 27 boat-based surveys scheduled. A total of 30 interviews were completed with an expanded catch estimate of 245,397 lbs. landed. The vessel/trailer participation survey is ongoing and includes all launching areas on the west coast of Saipan, where all boat-based fishing occurs. For this reporting period, a total of 104 boat vessels/trailers were registered as "out fishing". Because the same vessel may be out fishing on more than one day, this count should not be used to estimate the total number of unique fishing vessels. During this progress period, the most common fishing methods

encountered were trolling, bottomfishing and hook and line fishing. The expanded harvest estimate was 237,585 lbs. for trolling, while the estimated catch was 1,985 lbs. for bottomfishing and 2,767 lbs. for hook and line.

Consistent collection and entry of offshore data have continued. Issues with vehicle maintenance and repair are currently the biggest problems facing conducting offshore surveys. In November, a new data technician was hired to help with collection efforts.

#### **1.1.2.2 Vendor Invoice**

The DFW has been collecting fishery statistics on Saipan's commercial fishing fleet since the mid-1970s. With the assistance of the NMFS WPacFIN program, the DFW also expanded its fisheries monitoring programs to include the other two major inhabited islands in the CNMI, Rota and Tinian. The DFW's principal method of collecting domestic commercial fisheries data is a dealer invoicing system, sometimes referred to as a "trip ticket" system. The DFW provides numbered two-part invoices to all purchasers of fresh fishery products (including hotels, restaurants, stores, fish markets, and roadside vendors). Dealers then complete an invoice each time they purchase fish directly from fishers; one copy goes to the DFW and one copy goes to their records. Some advantages of this data collection method are that it is relatively inexpensive to implement and maintain, and it is fairly easy to completely cover the commercial fisheries. The DFW can also provide feedback to dealers and fishers to ensure data accuracy and continued cooperation over time.

There are some disadvantages to the trip ticket system, including: (1) dependency on non-DFW personnel to identify the catch and record the data, (2) restrictions on the types of data that can be collected, (3) required education and cooperation of all fish purchasers, and (4) limited recordings of fish actually sold to dealers. Therefore, a potentially important portion of the total landings typically goes unrecorded. Since 1982, the DFW has tried to minimize these disadvantages in several ways by (1) maintaining a close working relationship with dealers, (2) adding new dealers to their list and educating them, and (3) implementing a creel survey to help estimate total catch (including recreational and subsistence portion). The current system collects data from dealers in Saipan, where the DFW estimates more than 90 percent of all CNMI commercial landings are made. The DFW also estimates that the proportion of total commercial landings that have been recorded in the Saipan database since 1983 is about 90 percent; however, coverage has been relatively mottled over the years. Previous volumes of FSWP reported only recorded landings, but in recent volumes, the data have been adjusted to represent 100 percent coverage and are referenced as "estimated commercial landings" in the tables and figures.

These data elements are collected for all purchases of fishery products; however, species identification is frequently identified only to a group level, especially for reef fish.

For the period of January 1, 2018 to June 30, 2018, there were 1,618 invoices collected from 27 vendors around Saipan. A total of 58,215.49 lbs. of fish valued at \$175,166.22 were recorded from the sales receipt program. For the second half of the year over the period of July 1, 2018 to December 31, 2018, there were 1,174 invoices collected from 22 vendors around the island. A total of 46,876.00 lbs. of fish valued at \$129,757.08 were recorded from the sales receipt program. Consistent, scheduled visits to collect purchase data helped increase vendor

participation. Invoice collection in the months following Super Typhoon Yutu decreased due to the community's focus on recovery.

### 1.1.2.3 Bio-Sampling

The bio-sampling database contains general and specific bio-data obtained from individual commercial spearfish catches landed on Saipan from six different vendors over the course of 2011. The following data was captured for each fishing trip sampled: date, fishing gear type, time/hours fished, location fished, number/names of fishers, lengths/weights of individual fish, number/weight of octopus and squid, number/carapace size/weight/sex of lobster, and whether it was boat- or shore-based fishing trip.

Although sampling effort was intended to be spread evenly among all participating vendors, smaller vendors were inherently much more difficult to sample within the time constraints allowed. Therefore, a regular sampling schedule was implemented for the island's two largest vendors that included two weekdays and one weekend day each week starting in January/February 2011. Problems encountered in sampling the smaller vendors included: more days in any given month where no fish were purchased, the work area wasn't conducive for sampling, and communication problems. The bio-sampling database focuses on nighttime (non-SCUBA) spearfishing activities. Due to vendor-imposed limitations, other gear types that typically land their catch during normal business hours were not sampled.

### 1.1.2.4 Exemption Netting

In 2003, the use of gill nets was prohibited in the CNMI. In 2005, the DFW decided to allow gill netting under special circumstances. Gill netting is now allowed under strict conditions provided by the DFW with their permission such that all gill netting activities are to be monitored and recorded by DFW personnel.

In 2010, a law was passed allowing for the use of gill nets for the purpose of subsistence on the island of Rota. The following year, a regulation allowing subsistence net fishing was passed for the island of Tinian.

For a majority of the permitted gillnet activities, length and weight measurements were taken at the fishing site. Fork lengths were measured in millimeters and weights were measured in grams. If time did not permit for individual measurements, then length measurements were taken for each fish and total weight was taken for each species. Length/weight ratios were used to estimate weights of sampled fish. Information has been collected for activities conducted on the island of Saipan, but no official collection of information has been collected for Rota or Tinian.

### 1.1.2.5 Life History

The CNMI DFW life history program began in 1996 with the redgill emperors (*Lethrinus rubrioperculatus*). Since then, sampling has been conducted on other species, including *A. lineatus*, Myripristinae (*Myripristis violacea*, *M. kuntee*, *M. pralineae*, *M. bernti*, *M. murdjan*), *L. harak*, *Naso lituratus*, *Chlorurus sordidus*, and *C. undulatus*. Other life history programs have also developed over the past years. In collaboration with NMFS, DFW personnel collect life history information on *Scarus rubroviolaceus*, *Lethrinus atkinsoni*, and *Parupeneus barbarinus*

through funding provided by NOAA-NMFS. The life history survey captures biological information, including reproductive cycle, age at length, and age at maturity. The DFW is continually working to improve the understanding of reef fish life history in the CNMI through these types of programs.

#### **1.1.2.6 Monitoring of Imported Fish**

The DFW Fisheries Data Sections collect fisheries-related importation invoices from the Department of Commerce at the end of every month. The data is then entered into a ticket receipt system and reviewed prior to being sent out for compilation by PIFSC. A majority of the information entered into the system can only be identified to the family taxa.

#### **1.1.2.7 Vessel Inventory**

Little progress has been made towards the vessel inventory project, as staff were generally focused on improving inshore, offshore, and receipt data collection programs. This work is also impacted by policies of the CNMI Department of Public Safety, which manages vessel licensing. Going forward, additional emphasis will be put on improving the vessel inventory project, especially once the data technician and data manager positions are filled at the CNMI DFW.

#### **1.1.3 Meta-Data Dashboard Statistics**

The meta-data dashboard statistics describe the amount of data used or available to calculate the fishery-dependent information. Creel surveys are sampling-based systems that require a random-stratified design applied to pre-scheduled surveys. The number of sampling days, participation runs, and catch interviews can be used to determine if there are sufficient samples to run the expansion algorithm. The trends of these parameters over time may infer survey performance. Monitoring the survey performance is critical for explaining the reliability of the expanded information.

Commercial receipt book information depends on the amount of invoices submitted and the number of vendors participating in the program. Variations in these meta-data affect the commercial landing and revenue estimates.

##### **1.1.3.1 Creel Survey Meta-Data Statistics**

**Calculations:** Shore-based data

# Interview Days: Count of the number of actual days that Creel Survey Data were collected. It's a count of the number of unique dates found in the interview sampling data (the actual sampling date data, include opportunistic interviews).

# Participation Runs: Count of the number of unique occurrences of the combination of survey date and run number in the participation detail data.

# Catch Interviews: Count of the number of unique occurrences of the combination of date and run number in the participation detail data/ count of unique surveyor initials and date in PAR. This is divided into two categories, interviews conducted during scheduled survey days (Regular), and opportunistic interviews (Opp.) which are collected on non-scheduled days.



**Calculation:** Boat-based data

# Sample days: Count of the total number of unique dates found in the boat log data sampling date data.

# Catch Interviews: Count of the total number of data records found in the interview header data (number of interview headers). This is divided into two categories, interviews conducted during scheduled survey days (Regular), and opportunistic interviews (Opportunistic) which are collected on non-scheduled days.

**Table 1. Summary of CNMI creel survey meta-data**

| Year             | Shore-based      |                      |                    |               | Boat-based    |                    |               |
|------------------|------------------|----------------------|--------------------|---------------|---------------|--------------------|---------------|
|                  | # Interview Days | # Participation Runs | # Catch Interviews |               | # Sample Days | # Catch Interviews |               |
|                  |                  |                      | Regular            | Opportunistic |               | Regular            | Opportunistic |
| 2000             |                  |                      |                    |               | 44            | 168                | 9             |
| 2001             |                  |                      |                    |               | 67            | 285                | 0             |
| 2002             |                  |                      |                    |               | 75            | 200                | 25            |
| 2003             |                  |                      |                    |               | 90            | 299                | 40            |
| 2004             |                  |                      |                    |               | 77            | 272                | 16            |
| 2005             | 59               | 157                  | 258                | 42            | 78            | 417                | 29            |
| 2006             | 105              | 337                  | 597                | 248           | 71            | 342                | 22            |
| 2007             | 127              | 413                  | 601                | 36            | 62            | 314                | 1             |
| 2008             | 157              | 340                  | 911                | 24            | 55            | 250                | 1             |
| 2009             | 184              | 324                  | 870                | 24            | 64            | 241                | 25            |
| 2010             | 132              | 294                  | 374                | 29            | 65            | 161                | 82            |
| 2011             | 119              | 327                  | 388                | 14            | 67            | 162                | 87            |
| 2012             | 80               | 273                  | 230                | 10            | 72            | 166                | 0             |
| 2013             | 108              | 277                  | 297                | 2             | 71            | 191                | 0             |
| 2014             | 50               | 209                  | 108                | 1             | 71            | 166                | 0             |
| 2015             | 44               | 193                  | 83                 | 15            | 57            | 119                | 2             |
| 2016             | 44               | 256                  | 88                 | 20            | 65            | 117                | 3             |
| 2017             | 37               | 241                  | 122                | 57            | 66            | 120                | 6             |
| 2018             | 102              | 383                  | 254                | 21            | 53            | 113                | 5             |
| <b>10 yr avg</b> | <b>90</b>        | <b>278</b>           | <b>281</b>         | <b>19</b>     | <b>65</b>     | <b>156</b>         | <b>21</b>     |
| <b>10 yr SD</b>  | <b>45</b>        | <b>54</b>            | <b>224</b>         | <b>15</b>     | <b>6</b>      | <b>38</b>          | <b>33</b>     |
| <b>20 yr avg</b> | <b>96</b>        | <b>287</b>           | <b>370</b>         | <b>39</b>     | <b>67</b>     | <b>216</b>         | <b>19</b>     |
| <b>20 yr SD</b>  | <b>44</b>        | <b>70</b>            | <b>266</b>         | <b>60</b>     | <b>10</b>     | <b>84</b>          | <b>25</b>     |

**1.1.3.2 Commercial Receipt Book Statistics****Calculations:**

# Vendors: Count of the number of unique buyer codes found in the commercial purchase header data from the Commercial Receipt Book.







# Invoices: Count of the number of unique invoice numbers found in the commercial header data from the Commercial Receipt Book.

**Table 2. Summary of commercial receipt book meta-data**



























| <b>Year</b>      | <b>Number of Vendors</b> | <b>Total Invoices Collected</b> |
|------------------|--------------------------|---------------------------------|
| 1998             | 52                       | 5,369                           |
| 1999             | 49                       | 4,649                           |
| 2000             | 47                       | 6,030                           |
| 2001             | 39                       | 4,914                           |
| 2002             | 32                       | 4,759                           |
| 2003             | 24                       | 4,261                           |
| 2004             | 25                       | 3,507                           |
| 2005             | 23                       | 3,945                           |
| 2006             | 21                       | 4,002                           |
| 2007             | 18                       | 3,387                           |
| 2008             | 13                       | 3,054                           |
| 2009             | 6                        | 2,513                           |
| 2010             | 5                        | 1,612                           |
| 2011             | 3                        | 1,198                           |
| 2012             | 19                       | 1,565                           |
| 2013             | 17                       | 2,161                           |
| 2014             | 15                       | 1,665                           |
| 2015             | 10                       | 752                             |
| 2016             | 16                       | 2,100                           |
| 2017             | 29                       | 1,913                           |
| 2018             | 29                       | 2,180                           |
| <b>10 yr avg</b> | <b>15</b>                | <b>1,766</b>                    |
| <b>10 yr SD</b>  | <b>9</b>                 | <b>494</b>                      |
| <b>20 yr avg</b> | <b>22</b>                | <b>3,008</b>                    |
| <b>20 yr SD</b>  | <b>12</b>                | <b>1,418</b>                    |











































#### 1.1.4 Fishery Summary Dashboard Statistics

















The Fishery Summary Dashboard Statics section consolidates all fishery-dependent information comparing the most recent year with short-term (recent 10 years) and long-term (recent 20 years) average (shown bolded in [brackets]). Trend analysis of the past 10 years will dictate the trends (increasing, decreasing, or no trend). The right-most symbol indicates whether the mean of the short-term and long-term years were above, below, or within one standard deviation of the mean of the full time series.

|   |   |
|---|---|
| Legend Key:   |   |
|  - increasing trend in the time series |  - above 1 standard deviation  |
|  - decreasing trend in the time series |  - below 1 standard deviation  |
|  - no trend in the time series         |  - within 1 standard deviation |
| 10,000 [1,000] – point estimate of fishery statistic [difference from short/long term average]                          |   |

**Table 3. 2018 annual indicators for the coral reef and bottomfish fisheries describing performance relative to short-term (10-year) and long-term (20-year) averages**

| Fishery                                 | Fishery statistics   | Short-term (10 years)   | Long-term (20 years)  |
|---|--|---|---|
| <b>Bottomfish</b>                       | <b>Estimated catch (lbs.)</b>  |   |   |
| All species caught in the BF gear       | Boat and shore creel data estimated (expanded) total lbs. (all BF trips) | 858 [▼98%]         | 858 [▼98%]        |
|   | Estimated total lbs. (all species) commercial purchase data              | 4,048[▼72%]      | 4,048[▼82%]     |
| Bottomfish management unit species only | Boat-based creel data Estimated (expanded) total lbs. (all BF trips)     | 858 [▼98%]     | 858 [▼98%]    |
|   | Estimated total lbs. (all species) commercial purchase data              | 3,909[▼70%]    | 3,909[▼76%]   |
|   | <b>Catch-per-unit effort (lbs./gear hours)</b>                           |   |   |
|   | CPUE (creel data only)   | 0.0289[▼85%]   | N/A   |
|   | <b>Fishing effort (only available for creel data)</b>                    |   |   |
|   | Estimated (expanded) total bottomfish trips                              | 96[▼74%]       | N/A   |
|   | Estimated total bottomfishing gear hours                                 | 1,530[▼99%]    | N/A   |
|   | <b>Fishing participants</b>  |   |   |
|   | Estimated total # of fishers that went bottomfishing                     | 1,195[▼48%]    | N/A   |
|   | <b>Bycatch</b>   |   |   |
|   | Total number of bycatch caught   | 100[▼88%]      | N/A   |
|   | # bycatch released   | N/A   | N/A   |

| Fishery           | Fishery statistics                                     | Short-term (10 years)  | Long-term (20 years)   |
|-------------------|--|--|--|
|                   | # bycatch kept   | 100[▼88%]           | N/A  |
| <b>Coral Reef</b> | <b>Estimated catch (lbs.)</b>                          |  |  |
|                   | Boat-based creel data<br>(expanded estimate all gears) | 12,958[▼58%]        | 12,958[▼64%]   |
|                   | Shore-based creel<br>(expanded estimate all gears)     | 13,171[▼52%]        | N/A  |
|                   | Commercial Purchase                                    | 27,791[▼60%]        | 27,791[▼78%]   |
|                   | <b>Catch-per-unit-effort (lbs./gear hours)</b>         |  |  |
|                   | BB spear   | 0.1333 [▼92%]       | N/A  |
|                   | BB troll   | 0.1209[▲15%]        | N/A  |
|                   | BB atulai  | 0.2143 [▼42%]       | N/A  |
|                   | BB cast nets   | N/A  | N/A  |
|                   | SB hook and line                                       | 0.0014[▲17%]        | N/A  |
|                   | SB spear   | 0.0773[▼27%]      | N/A  |
|                   | SB cast nets   | 0.0404[▼49%]    | N/A  |
|                   | <b>Fishing effort (# of gear-hours by gear type)</b>   |  |  |
|                   | BB spear   | 12[▼88%]        | N/A  |
|                   | BB troll   | 75,117[▼48%]    | N/A  |
|                   | BB atulai  | 96[▼87%]        | N/A  |
|                   | BB cast nets   | N/A  | N/A  |
|                   | SB hook and line                                       | 118,668[▼55%]   | N/A  |
|                   | SB spear   | 2,394[▲78%]     | N/A  |
|                   | SB cast nets   | 544[▲547%]      | N/A  |
|                   | <b>Fishing participants (# of gear)</b>                |  |  |
|                   | BB spear   | 1,095[▲50%]     | N/A  |
|                   | BB troll   | 636[▼5%]        | N/A  |
|                   | BB atulai  | 852[▼14%]       | N/A  |
|                   | BB cast nets   | N/A  | N/A  |

| Fishery                    | Fishery statistics | Short-term (10 years)   | Long-term (20 years) |
|----------------------------|--------------------|---|----------------------|
|                            | SB hook and line   | 11,390[▼38%]   | N/A                  |
|                            | SB spear           | 1,544[▼43%]    | N/A                  |
|                            | SB cast nets       | 1,235[▼36%]    | N/A                  |
| <b>Boat-based Bycatch</b>  |                    |   |                      |
|                            | # bycatch caught   | 2,914[▼33%]    | N/A                  |
|                            | # bycatch released | N/A   | N/A                  |
|                            | # bycatch kept     | 2,914[▼33%]    | N/A                  |
| <b>Shore-based Bycatch</b> |                    |   |                      |
|                            | # bycatch caught   | 2,326[▼37%]    | N/A                  |
|                            | # bycatch released | 3[▲6%]         | N/A                  |
|                            | # bycatch kept     | 2,318[▲28%]    | N/A                  |

### 1.1.5 Catch Statistics

The following section summarizes the catch statistics for the bottomfish and coral reef fisheries in CNMI. Estimates of catch are summarized from the creel survey and commercial receipt book data collection programs. Catch statistics provide estimates of annual harvest from the different fisheries. Estimates of fishery removals can provide proxies for the level of fishing mortality and a reference level relative to established quotas. This section also provides detailed levels of catch for fishing methods and the top species complexes harvested in the coral reef and bottomfish fisheries.

#### 1.1.5.1 Catch by Data Stream

This section describes the estimated total catch from the shore- and boat-based creel survey programs as well as the commercial landings from the commercial receipt book system. The difference between the creel total and the commercial landings is assumed to be the non-commercial component. However, there are cases where the commercial landing may be higher than the estimated creel total of the commercial receipt book program. In this case, the commercial receipt books are able to capture the fishery better than the creel surveys.

**Calculations:** Estimated landings are based on all bottomfish species harvested, regardless of the gear used, for all data collection programs (e.g. shore-based creel, boat-based creel and the commercial purchase reports).

**Table 4. Summary of catch (lbs.) for all species caught by bottomfishing gear from 1983-2018 as recorded by creel surveys and the commercial purchase system**

| Year             | Creel survey Estimates |               | Creel Total   | Commercial Landings |
|------------------|------------------------|---------------|---------------|---------------------|
|                  | Shore-Based            | Boat-Based    |               |                     |
| 1983             |                        |               |               | 16,405              |
| 1984             |                        |               |               | 24,434              |
| 1985             |                        |               |               | 24,126              |
| 1986             |                        |               |               | 18,350              |
| 1987             |                        |               |               | 32,818              |
| 1988             |                        |               |               | 44,235              |
| 1989             |                        |               |               | 19,913              |
| 1990             |                        |               |               | 8,205               |
| 1991             |                        |               |               | 5,077               |
| 1992             |                        |               |               | 6,150               |
| 1993             |                        |               |               | 8,778               |
| 1994             |                        |               |               | 18,478              |
| 1995             |                        |               |               | 28,513              |
| 1996             |                        |               |               | 40,292              |
| 1997             |                        |               |               | 26,131              |
| 1998             |                        |               |               | 34,945              |
| 1999             |                        |               |               | 41,652              |
| 2000             |                        | 78,914        | 78,914        | 28,419              |
| 2001             |                        | 29,781        | 29,781        | 42,749              |
| 2002             |                        | 26,895        | 26,895        | 30,587              |
| 2003             |                        | 13,562        | 13,562        | 24,588              |
| 2004             |                        | 33,812        | 33,812        | 33,805              |
| 2005             | 214                    | 38,336        | 38,550        | 42,667              |
| 2006             | 79                     | 39,209        | 39,288        | 19,537              |
| 2007             | 648                    | 62,430        | 63,078        | 24,904              |
| 2008             | 300                    | 23,033        | 23,333        | 26,333              |
| 2009             | 140                    | 69,460        | 69,600        | 25,221              |
| 2010             | 2                      | 58,608        | 58,610        | 15,157              |
| 2011             | 556                    | 29,044        | 29,600        | 17,159              |
| 2012             | 84                     | 137,061       | 137,145       | 11,897              |
| 2013             | 1,332                  | 22,873        | 24,205        | 18,601              |
| 2014             | 166                    | 8,284         | 8,450         | 25,001              |
| 2015             | 215                    | 10,906        | 11,121        | 6,260               |
| 2016             | 36                     | 49,534        | 49,570        | 12,455              |
| 2017             | 59                     | 46,231        | 46,290        | 8,959               |
| 2018             | 299                    | 559           | 858           | 4,048               |
| <b>10 yr avg</b> | <b>289</b>             | <b>43,256</b> | <b>43,545</b> | <b>14,476</b>       |

|                  |            |               |               |               |
|------------------|------------|---------------|---------------|---------------|
| <b>10 yr SD</b>  | <b>380</b> | <b>38,053</b> | <b>37,927</b> | <b>6,843</b>  |
| <b>20 yr avg</b> | <b>295</b> | <b>40,975</b> | <b>41,193</b> | <b>23,000</b> |
| <b>20 yr SD</b>  | <b>340</b> | <b>30,685</b> | <b>30,628</b> | <b>11,297</b> |

**Calculations:** Estimated landings are based on a pre-determined list of species (Appendix A) identified as the BMUS Complex regardless of the gear used, for each data collection (shore-based creel, boat-based creel and the commercial purchase reports).

**Table 5. Summary of BMUS catch (lbs.) for from 1983-2018 for creel surveys and the commercial purchase system**

| Year | Creel survey Estimates |            | Creel Total | Commercial Landings |
|------|------------------------|------------|-------------|---------------------|
|      | Shore-Based            | Boat-Based |             |                     |
| 1983 |                        |            |             | 3,407               |
| 1984 |                        |            |             | 3,463               |
| 1985 |                        |            |             | 2,222               |
| 1986 |                        |            |             | 3,822               |
| 1987 |                        |            |             | 1,889               |
| 1988 |                        |            |             | 2,412               |
| 1989 |                        |            |             | 4,022               |
| 1990 |                        |            |             | 1,274               |
| 1991 |                        |            |             | 781                 |
| 1992 |                        |            |             | 607                 |
| 1993 |                        |            |             | 1,723               |
| 1994 |                        |            |             | 5,476               |
| 1995 |                        |            |             | 17,735              |
| 1996 |                        |            |             | 32,446              |
| 1997 |                        |            |             | 22,133              |
| 1998 |                        |            |             | 27,594              |
| 1999 |                        |            |             | 34,648              |
| 2000 |                        | 78,914     | 78,914      | 14,968              |
| 2001 |                        | 29,781     | 29,781      | 25,264              |
| 2002 |                        | 26,895     | 26,895      | 24,869              |
| 2003 |                        | 13,481     | 13,481      | 18,062              |
| 2004 |                        | 33,812     | 33,812      | 12,974              |
| 2005 | 168                    | 38,266     | 38,434      | 16,539              |
| 2006 | 67                     | 39,200     | 39,267      | 12,238              |
| 2007 | 648                    | 62,389     | 63,037      | 18,606              |
| 2008 | 69                     | 23,033     | 23,102      | 18,387              |
| 2009 | 140                    | 69,447     | 69,587      | 20,419              |
| 2010 | 2                      | 58,608     | 58,610      | 14,729              |
| 2011 | 556                    | 29,044     | 29,600      | 16,931              |
| 2012 | 84                     | 136,769    | 136,853     | 11,747              |

|                  |            |               |               |               |
|------------------|------------|---------------|---------------|---------------|
| 2013             | 1,332      | 22,733        | 24,065        | 17,770        |
| 2014             | 166        | 8,284         | 8,450         | 19,333        |
| 2015             | 215        | 10,906        | 11,121        | 4,197         |
| 2016             | 36         | 49,331        | 49,367        | 12,260        |
| 2017             | 59         | 46,231        | 46,290        | 8,918         |
| 2018             | 299        | 559           | 858           | 3,909         |
| <b>10 yr avg</b> | <b>289</b> | <b>43,191</b> | <b>43,480</b> | <b>13,021</b> |
| <b>10 yr SD</b>  | <b>380</b> | <b>37,984</b> | <b>37,858</b> | <b>5,615</b>  |
| <b>20 yr avg</b> | <b>274</b> | <b>40,931</b> | <b>41,133</b> | <b>16,338</b> |
| <b>20 yr SD</b>  | <b>347</b> | <b>30,641</b> | <b>30,591</b> | <b>6,911</b>  |

**Calculations:** Estimated landings are based on a pre-determined list of species (Appendix A) identified as the CREMUS Complex regardless of the gear used, for each data collection (shore-based creel, boat-based creel and the commercial purchase reports). It is required to finalize the CREMUS list to use for Creel and commercial landings and verify non-overlap between Bottomfish Complex and CREMUS. It is also required to verify all shallow bottomfish are not included in CREMUS list.

**Table 6. Summary of CREMUS catch (lbs.) from 1983-2018 for creel surveys and the commercial purchase system**

| Year | Creel survey Estimates |            | Creel Total | Commercial Landings |
|------|------------------------|------------|-------------|---------------------|
|      | Shore-Based            | Boat-Based |             |                     |
| 1983 |                        |            |             | 167,816             |
| 1984 |                        |            |             | 215,326             |
| 1985 |                        |            |             | 191,359             |
| 1986 |                        |            |             | 206,054             |
| 1987 |                        |            |             | 190,747             |
| 1988 |                        |            |             | 224,821             |
| 1989 |                        |            |             | 345,519             |
| 1990 |                        |            |             | 259,846             |
| 1991 |                        |            |             | 143,921             |
| 1992 |                        |            |             | 188,622             |
| 1993 |                        |            |             | 193,673             |
| 1994 |                        |            |             | 253,053             |
| 1995 |                        |            |             | 210,842             |
| 1996 |                        |            |             | 218,936             |
| 1997 |                        |            |             | 244,917             |
| 1998 |                        |            |             | 274,227             |
| 1999 |                        |            |             | 227,245             |
| 2000 |                        | 84,643     | 84,643      | 236,025             |
| 2001 |                        | 33,239     | 33,239      | 235,432             |
| 2002 |                        | 34,766     | 34,766      | 223,426             |



|                  |               |               |               |                |
|------------------|---------------|---------------|---------------|----------------|
| 2003             |               | 38,551        | 38,551        | 147,500        |
| 2004             |               | 27,698        | 27,698        | 127,517        |
| 2005             | 52,367        | 37,204        | 89,571        | 181,261        |
| 2006             | 55,657        | 42,893        | 98,550        | 176,349        |
| 2007             | 42,626        | 44,556        | 87,182        | 148,110        |
| 2008             | 43,380        | 64,320        | 107,700       | 160,542        |
| 2009             | 37,461        | 70,087        | 107,548       | 125,404        |
| 2010             | 23,425        | 49,505        | 72,930        | 89,567         |
| 2011             | 20,049        | 59,218        | 79,267        | 95,087         |
| 2012             | 19,069        | 49,401        | 68,470        | 68,158         |
| 2013             | 77,232        | 17,306        | 94,538        | 77,120         |
| 2014             | 8,541         | 15,482        | 24,023        | 75,062         |
| 2015             | 20,161        | 10,723        | 30,884        | 41,832         |
| 2016             | 40,531        | 17,990        | 58,521        | 56,192         |
| 2017             | 13,746        | 8,990         | 22,736        | 36,049         |
| 2018             | 13,171        | 12,958        | 26,129        | 27,791         |
| <b>10 yr avg</b> | <b>27,339</b> | <b>31,166</b> | <b>58,505</b> | <b>69,226</b>  |
| <b>10 yr SD</b>  | <b>19,217</b> | <b>21,962</b> | <b>29,532</b> | <b>28,415</b>  |
| <b>20 yr avg</b> | <b>23,371</b> | <b>35,977</b> | <b>59,347</b> | <b>127,783</b> |
| <b>20 yr SD</b>  | <b>22,116</b> | <b>22,055</b> | <b>32,538</b> | <b>67,750</b>  |

### 1.1.5.2 Expanded catch estimates by fishing methods

Catch information is provided for the top shore-based and boat-based fishing methods that comprises a majority of the annual catch.

**Calculations:** The creel survey catch time series are the sum of the estimated weight for selected gear in all strata for all species (except for trolling, which exclude PMUS as well as any other pelagic species complex).

**Table 7. Expanded CNMI creel survey catch estimates (lbs.) from 2005-2018**

| Year | Shore-Based methods |       |          | Boat-Based Methods |       |        |        |          |
|------|---------------------|-------|----------|--------------------|-------|--------|--------|----------|
|      | H&L                 | Spear | Cast Net | Bottomfish         | Spear | Troll* | Atulai | Cast net |
| 2005 | 130                 | 259   | 50       | 3,231              | 12    | 34,575 | 520    | 2        |
| 2006 | 262                 | 320   | 114      | 1,802              | 91    | 29,504 | 340    | 23       |
| 2007 | 203                 | 74    | 110      | 2,220              | 105   | 28,464 | 482    | 0        |
| 2008 | 335                 | 161   | 65       | 914                | 197   | 20,080 | 263    | 48       |
| 2009 | 295                 | 235   | 68       | 1,974              | 113   | 13,147 | 407    | 78       |
| 2010 | 105                 | 102   | 93       | 1,353              | 19    | 14,592 | 74     | 13       |
| 2011 | 136                 | 78    | 18       | 1,521              | 6     | 10,589 | 152    | 33       |
| 2012 | 93                  | 40    | 36       | 2,807              | 1     | 17,921 | 128    | 0        |
| 2013 | 170                 | 94    | 17       | 1,324              | 53    | 19,814 | 98     | 0        |
| 2014 | 55                  | 0     | 9        | 299                | 16    | 16,835 | 99     | 0        |

|                  |            |            |           |              |           |               |            |           |
|------------------|------------|------------|-----------|--------------|-----------|---------------|------------|-----------|
| 2015             | 27         | 123        | 10        | 470          | 81        | 15,491        | 76         | 0         |
| 2016             | 25         | 370        | 10        | 1,388        | 0         | 8,202         | 0          | 0         |
| 2017             | 108        | 82         | 22        | 1,046        | 153       | 14,131        | 0          | 0         |
| 2018             | 170        | 185        | 50        | 53           | 6         | 10,877        | 18         | 0         |
| <b>10 yr avg</b> | <b>118</b> | <b>131</b> | <b>33</b> | <b>1,224</b> | <b>45</b> | <b>14,160</b> | <b>105</b> | <b>12</b> |
| <b>10 yr SD</b>  | <b>77</b>  | <b>102</b> | <b>27</b> | <b>777</b>   | <b>51</b> | <b>3,395</b>  | <b>112</b> | <b>24</b> |
| <b>20 yr avg</b> | <b>151</b> | <b>152</b> | <b>48</b> | <b>1,457</b> | <b>61</b> | <b>18,159</b> | <b>190</b> | <b>14</b> |
| <b>20 yr SD</b>  | <b>92</b>  | <b>105</b> | <b>36</b> | <b>875</b>   | <b>61</b> | <b>7,480</b>  | <b>173</b> | <b>23</b> |

\*Excluding pelagic species

### 1.1.5.3 Top species in the catch for the boat and shore-based fisheries

Catch time series can act as indicators of fishery performance. Variations in the catch can be attributed to various factors, and there is no single explanatory variable for the observed trends. The ten species groups in the shore and boat-based catch records from the coral reef fishery make up a majority of the total annual catches.

**Calculations:** Catch by species complex is tallied directly from the boat-based expanded species composition data combining all gear types and species for all strata.

The averages for the table below were calculated from catch estimates for the entire time series across each of the CREMUS groupings. The average catch for each grouping is ranked from the highest to lowest. The dominant groups that make up more than half of the total annual catch are reported.

**Calculations:** Catch by species complex is tallied directly from the boat-based expanded species composition data combining all gear types and species, for all strata.

The averages for the table below were calculated from catch estimates from the entire time series for each of the CREMUS groupings. The average catch is ranked from the highest to lowest catch. The dominant groups that make up more than 60% of the catch are reported.

**Table 8. Catch for top species complexes in CNMI boat-based creel data from 2000-2018**

| Year             | Boat-Based Estimated Pounds |               |              |              |              |              |            |              |              |            |              |              |
|------------------|-----------------------------|---------------|--------------|--------------|--------------|--------------|------------|--------------|--------------|------------|--------------|--------------|
|                  | Bottomfish                  | BMUS          | Emperors     | Jacks        | Atulai       | Grouper      | Snapper    | Surgeonfish  | Parrotfish   | Mullet     | Squirrelfish | Rudderfish   |
| 2000             | 82,358                      | 82,358        | 34,850       | 14,242       | 967          | 2,035        | 2,193      | 3,648        | 1,266        | 176        | 99           | 0            |
| 2001             | 33,938                      | 33,938        | 9,774        | 3,094        | 4,456        | 1,156        | 623        | 5,569        | 1,453        | 0          | 69           | 13           |
| 2002             | 41,651                      | 41,651        | 9,946        | 5,904        | 613          | 260          | 645        | 3,831        | 2,032        | 818        | 879          | 0            |
| 2003             | 17,319                      | 17,238        | 1,339        | 10,958       | 13,579       | 883          | 240        | 3,924        | 935          | 0          | 2,030        | 265          |
| 2004             | 37,792                      | 37,792        | 3,675        | 11,215       | 1,008        | 1,186        | 1,020      | 2,153        | 1,306        | 0          | 503          | 600          |
| 2005             | 41,410                      | 41,340        | 3,242        | 17,733       | 0            | 1,617        | 2,282      | 1,722        | 776          | 0          | 47           | 925          |
| 2006             | 42,118                      | 42,109        | 8,086        | 8,700        | 2,932        | 1,336        | 590        | 4,260        | 1,792        | 0          | 340          | 235          |
| 2007             | 77,315                      | 77,274        | 9,934        | 4,280        | 7,336        | 2,424        | 2,716      | 3,948        | 2,778        | 0          | 4,391        | 985          |
| 2008             | 23,633                      | 23,633        | 15,785       | 6,939        | 14,039       | 1,025        | 595        | 5,572        | 4,378        | 0          | 1,104        | 520          |
| 2009             | 74,883                      | 74,870        | 18,669       | 2,197        | 20,622       | 3,501        | 548        | 7,506        | 3,910        | 0          | 635          | 3,189        |
| 2010             | 62,529                      | 62,529        | 10,980       | 12,847       | 6,195        | 745          | 1,430      | 3,934        | 1,364        | 0          | 780          | 0            |
| 2011             | 32,552                      | 32,552        | 15,534       | 10,238       | 7,847        | 5,160        | 178        | 4,016        | 205          | 0          | 542          | 3,715        |
| 2012             | 137,118                     | 136,826       | 16,418       | 974          | 14,438       | 4,231        | 123        | 974          | 1,147        | 0          | 1,150        | 88           |
| 2013             | 23,068                      | 22,928        | 5,221        | 1,400        | 720          | 1,011        | 64         | 955          | 60           | 0          | 2            | 175          |
| 2014             | 8,284                       | 8,284         | 4,638        | 5,161        | 330          | 8            | 37         | 2,063        | 695          | 0          | 236          | 0            |
| 2015             | 10,906                      | 10,906        | 1,436        | 1,037        | 111          | 2,068        | 325        | 4,218        | 277          | 0          | 345          | 127          |
| 2016             | 49,534                      | 49,330        | 1,689        | 9,467        | 0            | 47           | 194        | 0            | 0            | 0          | 57           | 0            |
| 2017             | 46,231                      | 46,231        | 691          | 2,219        | 3,122        | 0            | 0          | 0            | 17           | 0          | 4            | 0            |
| 2018             | 559                         | 559           | 8,356        | 855          | 573          | 209          | 30         | 1,149        | 0            | 0          | 0            | 0            |
| <b>10 yr avg</b> | <b>44,502</b>               | <b>44,566</b> | <b>8,363</b> | <b>4,640</b> | <b>5,396</b> | <b>1,698</b> | <b>293</b> | <b>2,482</b> | <b>768</b>   | <b>0</b>   | <b>375</b>   | <b>729</b>   |
| <b>10 yr SD</b>  | <b>38,441</b>               | <b>38,507</b> | <b>6,360</b> | <b>4,305</b> | <b>6,732</b> | <b>1,840</b> | <b>410</b> | <b>2,271</b> | <b>1,148</b> | <b>0</b>   | <b>374</b>   | <b>1,368</b> |
| <b>20 yr avg</b> | <b>44,334</b>               | <b>44,379</b> | <b>9,488</b> | <b>6,814</b> | <b>5,205</b> | <b>1,521</b> | <b>728</b> | <b>3,129</b> | <b>1,284</b> | <b>52</b>  | <b>695</b>   | <b>570</b>   |
| <b>20 yr SD</b>  | <b>31,324</b>               | <b>31,365</b> | <b>8,084</b> | <b>4,964</b> | <b>6,052</b> | <b>1,410</b> | <b>810</b> | <b>1,973</b> | <b>1,233</b> | <b>185</b> | <b>1,010</b> | <b>1,037</b> |

**Table 9. Catch for top species complexes in CNMI shore-based creel data from 2000-2018**

| Year             | Shore-based Estimated Pounds |              |              |              |               |              |              |              |              |              |              |
|------------------|------------------------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|
|                  | Jacks                        | Emperors     | Rabbitfish   | Surgeonfish  | Goatfish      | Atulai       | Parrotfish   | Mollusks     | Mullet       | Wrasse       | Rudderfish   |
| 2005             | 7,482                        | 16,943       | 5,619        | 12,378       | 8,822         | 631          | 5,261        | 14,262       | 6,172        | 4,721        | 145          |
| 2006             | 12,487                       | 18,142       | 4,771        | 16,613       | 14,981        | 8,990        | 7,228        | 2,072        | 7,380        | 3,887        | 1,004        |
| 2007             | 7,833                        | 16,551       | 3,982        | 3,428        | 3,605         | 9,992        | 3,978        | 8,025        | 8,321        | 3,328        | 360          |
| 2008             | 26,271                       | 12,850       | 3,209        | 8,178        | 5,568         | 4,821        | 10,751       | 325          | 3,878        | 2,534        | 52           |
| 2009             | 14,409                       | 13,502       | 2,723        | 10,185       | 7,539         | 1,124        | 6,471        | 2,734        | 4,419        | 2,327        | 3,529        |
| 2010             | 3,393                        | 3,315        | 659          | 2,196        | 1,697         | 26,130       | 1,484        | 0            | 1,981        | 1,296        | 155          |
| 2011             | 10,958                       | 5,234        | 504          | 4,204        | 1,378         | 1,176        | 1,394        | 0            | 2,702        | 1,103        | 87           |
| 2012             | 7,275                        | 3,005        | 9,593        | 472          | 3,755         | 303          | 368          | 3,588        | 2,487        | 852          | 0            |
| 2013             | 8,239                        | 8,902        | 12,715       | 33,136       | 38,353        | 11,033       | 23,290       | 0            | 1,052        | 4,120        | 2,286        |
| 2014             | 8,233                        | 2,513        | 55           | 229          | 182           | 73           | 0            | 0            | 742          | 42           | 1,354        |
| 2015             | 2,096                        | 6,226        | 6,855        | 6,114        | 5,529         | 320          | 7,471        | 0            | 844          | 387          | 400          |
| 2016             | 3,947                        | 6,829        | 3,969        | 5,394        | 2,851         | 0            | 13,342       | 0            | 616          | 0            | 0            |
| 2017             | 12,046                       | 1,307        | 926          | 2,819        | 1,006         | 0            | 3,065        | 2,370        | 1,203        | 267          | 1,250        |
| 2018             | 1,256                        | 3,020        | 2,245        | 2,038        | 3,718         | 1,767        | 8,176        | 182          | 537          | 40           | 1,464        |
| <b>10 yr avg</b> | <b>7,185</b>                 | <b>5,385</b> | <b>6,679</b> | <b>4,024</b> | <b>6,601</b>  | <b>4,193</b> | <b>6,506</b> | <b>887</b>   | <b>1,658</b> | <b>1,043</b> | <b>1,053</b> |
| <b>10 yr SD</b>  | <b>4,221</b>                 | <b>3,486</b> | <b>9,258</b> | <b>4,105</b> | <b>10,791</b> | <b>7,967</b> | <b>6,895</b> | <b>1,346</b> | <b>1,180</b> | <b>1,239</b> | <b>1,111</b> |
| <b>20 yr avg</b> | <b>8,995</b>                 | <b>8,453</b> | <b>7,670</b> | <b>4,130</b> | <b>7,070</b>  | <b>4,740</b> | <b>6,591</b> | <b>2,397</b> | <b>3,024</b> | <b>1,779</b> | <b>863</b>   |
| <b>20 yr SD</b>  | <b>6,125</b>                 | <b>5,772</b> | <b>8,397</b> | <b>3,506</b> | <b>9,436</b>  | <b>7,072</b> | <b>5,986</b> | <b>3,944</b> | <b>2,539</b> | <b>1,622</b> | <b>1,005</b> |

### 1.1.6 Catch-per-Unit-Effort (CPUE) Statistics

This section summarizes the estimates for CPUE in the boat- and shore-based fisheries. The boat-based fisheries include the bottomfishing (handline gear), spearfishing (snorkel), troll, atulai nets, and cast nets, which comprise 84% of the total catch. Trolling is primarily a pelagic fishing method but also catches coral reef fishes like jacks and gray jobfish. The shore-based fisheries include the hook-and-line, spearfishing and cast nets, which comprise 99% of the total coral reef fish catch. CPUE is reported as pounds per gear-hour for the shore-based fishery, and pounds per fishing trip in the boat-based fishery.

**Calculations:** CPUE is calculated from interview data by gear type using  $\sum \text{catch} / \sum (\text{hours fished} * \text{number of fishers})$  for boat based and  $\sum \text{catch} / \sum (\text{hours fished} * \text{number of gears used})$  for shore based. If the value is blank (empty), then there was no interview collected for that method. Landings from interviews without fishing hours are excluded from the calculations.

**Table 10. CPUE for top fishing methods in CNMI shore-based fisheries from 2005-2018**

| Year             | Shore-Based Gear CPUE (lbs./gear hour) |               |               |
|------------------|--|---------------|---------------|
|                  | H&L                                    | Spear         | Cast Net      |
| 2005             | 0.0009                                 | 0.0654        | 0.0321        |
| 2006             | 0.0002                                 | 0.0434        | 0.0158        |
| 2007             | 0.0003                                 | 0.0705        | 0.0340        |
| 2008             | 0.0002                                 | 0.0658        | 0.0074        |
| 2009             | 0.0002                                 | 0.0623        | 0.0280        |
| 2010             | 0.0004                                 | 0.0567        | 0.1771        |
| 2011             | 0.0005                                 | 0.0556        | 0.0557        |
| 2012             | 0.0004                                 | 0.0465        | 0.1000        |
| 2013             | 0.0009                                 | 0.1302        | 0.0833        |
| 2014             | 0.0024                                 | 0.0000        | 0.1500        |
| 2015             | 0.0017                                 | 0.1538        | 0.1042        |
| 2016             | 0.0020                                 | 0.2864        | 0.0526        |
| 2017             | 0.0016                                 | 0.1911        | 0.0404        |
| 2018             | 0.0014                                 | 0.0773        | 0.0058        |
| <b>10 yr avg</b> | <b>0.0012</b>                          | <b>0.1060</b> | <b>0.0797</b> |
| <b>10 yr SD</b>  | <b>0.0007</b>                          | <b>0.0807</b> | <b>0.0514</b> |
| <b>20 yr avg</b> | <b>0.0009</b>                          | <b>0.0932</b> | <b>0.0633</b> |
| <b>20 yr SD</b>  | <b>0.0007</b>                          | <b>0.0714</b> | <b>0.0510</b> |

**Table 11. CPUE for top fishing methods in CNMI boat-based fisheries from 2000-2018**

| Year | Boat-Based Gear CPUE (lbs./gear hours) |        |        |        |          |
|------|--|--------|--------|--------|----------|
|      | Bottomfishing                          | Spear  | Troll  | Atulai | Cast Net |
| 2000 | 0.1102                                 | 2.3929 | 0.0837 | 0.1326 | 0.0000   |
| 2001 | 0.0301                                 | 1.4844 | 0.0588 | 0.1067 | 0.0000   |
| 2002 | 0.0485                                 | 3.9000 | 0.0608 | 0.1079 | 0.0000   |

|                  |               |               |               |               |               |
|------------------|---------------|---------------|---------------|---------------|---------------|
| 2003             | 0.0345        | 0.1009        | 0.0371        | 0.2284        | 1.4000        |
| 2004             | 0.0307        | 0.0839        | 0.0568        | 0.0480        | 0.0000        |
| 2005             | 0.0137        | 1.0000        | 0.0372        | 0.0704        | 0.1250        |
| 2006             | 0.0126        | 0.1071        | 0.0545        | 0.0437        | 1.1500        |
| 2007             | 0.0289        | 0.3182        | 0.0726        | 0.0311        | 0.0000        |
| 2008             | 0.0125        | 0.0533        | 0.0718        | 0.1927        | 0.6667        |
| 2009             | 0.0069        | 0.1495        | 0.0745        | 0.0755        | 5.5714        |
| 2010             | 0.0022        | 3.1667        | 0.1065        | 0.2284        | 1.4444        |
| 2011             | 0.0021        | 1.0000        | 0.0855        | 0.6609        | 0.3929        |
| 2012             | 0.3558        | 0.2500        | 0.1113        | 0.0914        | 0.0000        |
| 2013             | 0.1445        | 0.3155        | 0.0982        | 0.2917        | 0.0000        |
| 2014             | 0.1286        | 3.2000        | 0.0866        | 0.5789        | 0.0000        |
| 2015             | 0.2318        | 27.0000       | 0.1594        | 0.7917        | 0.0000        |
| 2016             | 0.3541        | 0.0000        | 0.0893        | 0.0000        | 0.0000        |
| 2017             | 0.6671        | 1.1333        | 0.1184        | 0.0000        | 0.0000        |
| 2018             | 0.0289        | 0.3333        | 0.1209        | 0.2143        | 0.0000        |
| <b>10 yr avg</b> | <b>0.1922</b> | <b>4.0609</b> | <b>0.1051</b> | <b>0.3666</b> | <b>1.8522</b> |
| <b>10 yr SD</b>  | <b>0.2049</b> | <b>8.1883</b> | <b>0.0232</b> | <b>0.2551</b> | <b>2.2113</b> |
| <b>20 yr avg</b> | <b>0.1181</b> | <b>2.5549</b> | <b>0.0834</b> | <b>0.2291</b> | <b>1.1945</b> |
| <b>20 yr SD</b>  | <b>0.1690</b> | <b>6.0490</b> | <b>0.0301</b> | <b>0.2228</b> | <b>1.6392</b> |

### 1.1.7 Effort Statistics

This section summarizes the effort trends in the coral reef and bottomfish fishery. Fishing effort trends provide insights on the level of fishing pressure through time. Effort information is provided for the top shore-based and boat-based fishing methods that comprise a majority of the annual catch.

**Calculations:** Effort estimates (hours) are generated by summing the effort data collected from interviews by gear type. For shore-based estimates, data collection started in 2005.

**Table 12. Effort estimates in CNMI coral reef and bottomfish fisheries from 2000-2018**

| Year | Estimated Effort by Gear Type |       |          |                       |       |         |        |          |
|------|-------------------------------|-------|----------|-----------------------|-------|---------|--------|----------|
|      | Shore-Based Gear Hours        |       |          | Boat-Based Gear Hours |       |         |        |          |
|      | H&L                           | Spear | Cast Net | Bottom                | Spear | Troll   | Atulai | Cast Net |
| 2000 |                               |       |          | 15,194                | 21    | 131,472 | 2,379  | 0        |
| 2001 |                               |       |          | 26,076                | 16    | 475,304 | 2,400  | 0        |
| 2002 |                               |       |          | 23,547                | 10    | 286,520 | 1,888  | 0        |
| 2003 |                               |       |          | 16,492                | 3,420 | 841,750 | 918    | 5        |
| 2004 |                               |       |          | 40,633                | 666   | 462,027 | 4,620  | 0        |
| 2005 | 143,992                       | 3,960 | 1,560    | 230,736               | 12    | 899,028 | 7,062  | 16       |
| 2006 | 1,145,508                     | 7,380 | 7,216    | 145,722               | 918   | 505,362 | 7,020  | 8        |

|                  |                |              |              |                |              |                |              |           |
|------------------|----------------|--------------|--------------|----------------|--------------|----------------|--------------|-----------|
| 2007             | 677,265        | 1,050        | 3,233        | 70,168         | 352          | 359,047        | 14,602       | 6         |
| 2008             | 1,464,036      | 2,448        | 8,736        | 71,463         | 3,780        | 261,960        | 1,521        | 36        |
| 2009             | 1,494,570      | 3,774        | 2,432        | 305,064        | 714          | 173,600        | 5,159        | 7         |
| 2010             | 238,815        | 1,800        | 525          | 658,504        | 6            | 136,413        | 297          | 9         |
| 2011             | 286,144        | 1,403        | 323          | 869,240        | 6            | 117,576        | 230          | 56        |
| 2012             | 216,905        | 860          | 360          | 8,211          | 4            | 169,278        | 1,200        | 0         |
| 2013             | 182,684        | 722          | 204          | 9,480          | 168          | 212,346        | 392          | 0         |
| 2014             | 23,023         | 2            | 60           | 2,625          | 10           | 216,425        | 171          | 3         |
| 2015             | 15,624         | 800          | 96           | 2,340          | 6            | 107,514        | 96           | 0         |
| 2016             | 12,402         | 1,292        | 190          | 5,376          | 0            | 99,828         | 0            | 0         |
| 2017             | 67,081         | 429          | 544          | 1,568          | 81           | 124,845        | 0            | 0         |
| 2018             | 118,668        | 2,394        | 8,667        | 1,530          | 12           | 75,117         | 96           | 0         |
| <b>10 yr avg</b> | <b>265,592</b> | <b>1,348</b> | <b>1,340</b> | <b>186,394</b> | <b>101</b>   | <b>143,294</b> | <b>764</b>   | <b>8</b>  |
| <b>10 yr SD</b>  | <b>420,380</b> | <b>1,038</b> | <b>2,529</b> | <b>305,785</b> | <b>211</b>   | <b>45,362</b>  | <b>1,502</b> | <b>16</b> |
| <b>20 yr avg</b> | <b>434,766</b> | <b>2,022</b> | <b>2,439</b> | <b>131,788</b> | <b>537</b>   | <b>297,653</b> | <b>2,634</b> | <b>8</b>  |
| <b>20 yr SD</b>  | <b>518,328</b> | <b>1,869</b> | <b>3,161</b> | <b>234,001</b> | <b>1,087</b> | <b>235,198</b> | <b>3,616</b> | <b>14</b> |

### 1.1.8 Participants

This section summarizes the estimated number of participants in each fishery. The information presented here can be used in the impact analysis of potential amendments in the FEPs associated with the bottomfish and coral reef fisheries. The trend in the number of participants over time can also be used as an indicator for fishing pressure.

**Calculations:** Estimated number of participants is calculated by using an average number of fishers out fishing per day multiplied by the numbers of dates in the calendar year by gear type. The total is a combination of weekend and weekday stratum estimates.

**Table 13. Number of fishermen participating in the CNMI bottomfish fishery and number of gears in the CNMI coral reef fishery from 2000-2018**

| Year             | Bottomfish   |              | Coral Reef Boat-Based |            |              |            | Coral Reef Shore-Based |              |              |
|------------------|--------------|--------------|-----------------------|------------|--------------|------------|------------------------|--------------|--------------|
|                  | # Fishers    | # Gears      | Spear                 | Troll      | Atulai       | Cast Net   | H&L                    | Spear        | Cast Net     |
| 2000             | 1,161        | 1,119        | 1,464                 | 803        | 1,577        | 0          |                        |              |              |
| 2001             | 993          | 898          | 1,460                 | 806        | 1,095        | 0          |                        |              |              |
| 2002             | 1,259        | 1,287        | 730                   | 851        | 1,156        | 0          |                        |              |              |
| 2003             | 1,374        | 1,331        | 816                   | 930        | 913          | 730        |                        |              |              |
| 2004             | 1,319        | 1,236        | 993                   | 793        | 1,313        | 0          |                        |              |              |
| 2005             | 1,369        | 1,342        | 1,095                 | 850        | 1,007        | 730        | 43,884                 | 7,058        | 4,798        |
| 2006             | 1,130        | 1,155        | 830                   | 870        | 973          | 1,825      | 49,116                 | 8,448        | 5,251        |
| 2007             | 883          | 807          | 782                   | 800        | 1,186        | 1,095      | 41,127                 | 6,554        | 3,521        |
| 2008             | 1,888        | 1,843        | 848                   | 723        | 1,423        | 976        | 58,569                 | 5,270        | 4,547        |
| 2009             | 3,043        | 3,224        | 821                   | 671        | 1,345        | 730        | 42,908                 | 4,137        | 2,771        |
| 2010             | 6,375        | 6,727        | 730                   | 660        | 876          | 1,095      | 17,505                 | 3,039        | 2,145        |
| 2011             | 6,246        | 7,581        | 730                   | 758        | 913          | 730        | 24,927                 | 2,049        | 3,134        |
| 2012             | 690          | 718          | 366                   | 738        | 1,281        | 0          | 17,198                 | 2,751        | 2,075        |
| 2013             | 728          | 753          | 728                   | 655        | 874          | 0          | 22,960                 | 2,870        | 2,728        |
| 2014             | 666          | 751          | 365                   | 626        | 1,095        | 730        | 13,601                 | 2,452        | 1,656        |
| 2015             | 678          | 782          | 365                   | 641        | 730          | 0          | 8,374                  | 2,769        | 817          |
| 2016             | 641          | 878          | 0                     | 633        | 0            | 0          | 11,804                 | 3,225        | 1,544        |
| 2017             | 786          | 786          | 1,369                 | 650        | 0            | 0          | 13,376                 | 2,108        | 1,290        |
| 2018             | 1,195        | 995          | 1,095                 | 636        | 852          |            | 11,390                 | 1,544        | 1,235        |
| <b>10 yr avg</b> | <b>2,320</b> | <b>2,105</b> | <b>730</b>            | <b>667</b> | <b>996</b>   | <b>821</b> | <b>18,404</b>          | <b>2,694</b> | <b>1,940</b> |
| <b>10 yr SD</b>  | <b>2,529</b> | <b>2,213</b> | <b>325</b>            | <b>43</b>  | <b>206</b>   | <b>158</b> | <b>9,527</b>           | <b>683</b>   | <b>722</b>   |
| <b>20 yr avg</b> | <b>1,801</b> | <b>1,707</b> | <b>866</b>            | <b>742</b> | <b>1,095</b> | <b>960</b> | <b>26,910</b>          | <b>3,877</b> | <b>2,679</b> |
| <b>20 yr SD</b>  | <b>1,924</b> | <b>1,670</b> | <b>327</b>            | <b>93</b>  | <b>227</b>   | <b>341</b> | <b>16,073</b>          | <b>2,049</b> | <b>1,362</b> |



| Year             | Bottomfish |            | Coral Reef Boat-Based |              |           |          | Coral Reef Shore-Based |              |              |
|------------------|------------|------------|-----------------------|--------------|-----------|----------|------------------------|--------------|--------------|
|                  | # Gears    | # Trips    | Spears                | Troll        | Atulai    | Cast Net | H&L                    | Cast Net     | Spears       |
| 2000             | 366        | 441        | 10                    | 903          | 133       | 0        |                        |              |              |
| 2001             | 365        | 425        | 6                     | 1,401        | 119       | 0        |                        |              |              |
| 2002             | 365        | 185        | 5                     | 976          | 83        | 0        |                        |              |              |
| 2003             | 365        | 231        | 112                   | 1,913        | 53        | 3        |                        |              |              |
| 2004             | 366        | 390        | 37                    | 934          | 112       | 0        |                        |              |              |
| 2005             | 365        | 824        | 2                     | 531          | 53        | 2        | 45,558                 | 4,715        | 7,058        |
| 2006             | 365        | 843        | 30                    | 553          | 73        | 2        | 52,248                 | 5,160        | 8,448        |
| 2007             | 365        | 695        | 33                    | 793          | 200       | 4        | 42,591                 | 3,479        | 6,872        |
| 2008             | 366        | 592        | 57                    | 377          | 50        | 5        | 60,468                 | 4,482        | 5,167        |
| 2009             | 365        | 587        | 7                     | 101          | 27        | 0        | 44,638                 | 2,744        | 4,137        |
| 2010             | 365        | 421        | 0                     | 35           | 2         | 1        | 18,980                 | 2,086        | 3,069        |
| 2011             | 365        | 452        | 0                     | 27           | 1         | 1        | 26,575                 | 3,054        | 2,036        |
| 2012             | 366        | 320        | 14                    | 1,916        | 215       | 0        | 18,388                 | 2,236        | 2,751        |
| 2013             | 364        | 292        | 85                    | 1,711        | 85        | 0        | 24,536                 | 2,649        | 2,870        |
| 2014             | 365        | 211        | 21                    | 2,868        | 94        | 10       | 14,062                 | 1,656        | 2,410        |
| 2015             | 365        | 173        | 24                    | 2,409        | 97        | 0        | 8,828                  | 817          | 2,769        |
| 2016             | 366        | 113        | 0                     | 1,433        | 0         | 0        | 12,455                 | 1,733        | 3,637        |
| 2017             | 365        | 88         | 117                   | 2,646        | 0         | 0        | 13,594                 | 1,327        | 2,145        |
| 2018             | 365        | 96         | 24                    | 1,789        | 97        | 0        | 11,688                 | 1,349        | 1,567        |
| <b>10 yr avg</b> | <b>275</b> | <b>365</b> | <b>29</b>             | <b>1,494</b> | <b>62</b> | <b>1</b> | <b>19,374</b>          | <b>1,965</b> | <b>2,739</b> |
| <b>10 yr SD</b>  | <b>161</b> | <b>1</b>   | <b>38</b>             | <b>1,029</b> | <b>66</b> | <b>3</b> | <b>9,988</b>           | <b>678</b>   | <b>722</b>   |
| <b>20 yr avg</b> | <b>388</b> | <b>365</b> | <b>31</b>             | <b>1,227</b> | <b>79</b> | <b>1</b> | <b>28,186</b>          | <b>2,678</b> | <b>3,924</b> |
| <b>20 yr SD</b>  | <b>228</b> | <b>1</b>   | <b>36</b>             | <b>855</b>   | <b>60</b> | <b>2</b> | <b>16,700</b>          | <b>1,306</b> | <b>2,066</b> |

### 1.1.9 Bycatch Estimates

This section focuses on MSA § 303(a)(11), which requires that all FMPs establish a standardized reporting methodology to assess the amount and type of bycatch occurring in the fishery, and include conservation and management measures that, to the extent practicable, minimize bycatch and bycatch mortality. The MSA § 303(a)(11) standardized reporting methodology is commonly referred to as a “Standardized Bycatch Reporting Methodology” (SBRM) and was added to the MSA by the Sustainable Fisheries Act of 1996 (SFA). The Council implemented omnibus amendments to FMPs in 2003 to address MSA bycatch provisions and established SBRMs at that time.

**Calculations:** The number caught is the sum of the total number of individuals found in the raw data including bycatch. The number kept is the total number of individuals in the raw data that are not marked as bycatch. The number released is bycatch caught minus the number of bycatch kept. Percent bycatch is the sum of all bycatch divided by the total catch.

**Table 14. Bycatch estimates in CNMI non-bottomfishing boat-based fisheries from 2000-2018**

| Year             | Boat-Based Non-Bottomfishing Bycatch |              |          |               |
|------------------|--------------------------------------|--------------|----------|---------------|
|                  | # Caught                             | Kept         | Released | % Bycatch     |
| 2000             | 3,089                                | 3,086        | 3        | 0.0010        |
| 2001             | 5,732                                | 5,731        | 1        | 0.0002        |
| 2002             | 4,885                                | 4,885        | 0        | 0.0000        |
| 2003             | 8,785                                | 8,785        | 0        | 0.0000        |
| 2004             | 5,717                                | 5,717        | 0        | 0.0000        |
| 2005             | 6,772                                | 6,772        | 0        | 0.0000        |
| 2006             | 6,761                                | 6,759        | 2        | 0.0003        |
| 2007             | 6,683                                | 6,683        | 0        | 0.0000        |
| 2008             | 4,463                                | 4,463        | 0        | 0.0000        |
| 2009             | 3,792                                | 3,792        | 0        | 0.0000        |
| 2010             | 3,462                                | 3,462        | 0        | 0.0000        |
| 2011             | 2,515                                | 2,515        | 0        | 0.0000        |
| 2012             | 3,963                                | 3,963        | 0        | 0.0000        |
| 2013             | 3,732                                | 3,732        | 0        | 0.0000        |
| 2014             | 2,600                                | 2,600        | 0        | 0.0000        |
| 2015             | 2,693                                | 2,693        | 0        | 0.0000        |
| 2016             | 1,812                                | 1,812        | 0        | 0.0000        |
| 2017             | 2,632                                | 2,632        | 0        | 0.0000        |
| 2018             | 1,940                                | 1,940        | 0        | 0.0000        |
| <b>10 yr avg</b> | <b>2,914</b>                         | <b>2,914</b> | <b>0</b> | <b>0.0000</b> |
| <b>10 yr SD</b>  | <b>734</b>                           | <b>734</b>   | <b>0</b> | <b>0.0000</b> |
| <b>20 yr avg</b> | <b>4,317</b>                         | <b>4,317</b> | <b>0</b> | <b>0.0001</b> |
| <b>20 yr SD</b>  | <b>1,898</b>                         | <b>1,898</b> | <b>1</b> | <b>0.0002</b> |

**Table 15. Bycatch estimates in the CNMI bottomfish fishery from 2000-2018**

| Year             | Boat-Based Bottomfishing Bycatch |              |           |               |
|------------------|----------------------------------|--------------|-----------|---------------|
|                  | # Caught                         | Kept         | Released  | % Bycatch     |
| 2000             | 818                              | 797          | 21        | 0.0257        |
| 2001             | 931                              | 930          | 1         | 0.0011        |
| 2002             | 904                              | 890          | 14        | 0.0155        |
| 2003             | 877                              | 841          | 36        | 0.0410        |
| 2004             | 1,379                            | 1,359        | 20        | 0.0145        |
| 2005             | 3,225                            | 3,221        | 4         | 0.0012        |
| 2006             | 1,845                            | 1,842        | 3         | 0.0016        |
| 2007             | 2,110                            | 2,110        | 0         | 0.0000        |
| 2008             | 1,158                            | 1,158        | 0         | 0.0000        |
| 2009             | 1,779                            | 1,779        | 0         | 0.0000        |
| 2010             | 1,474                            | 1,474        | 0         | 0.0000        |
| 2011             | 1,734                            | 1,734        | 0         | 0.0000        |
| 2012             | 782                              | 782          | 0         | 0.0000        |
| 2013             | 857                              | 857          | 0         | 0.0000        |
| 2014             | 216                              | 216          | 0         | 0.0000        |
| 2015             | 196                              | 196          | 0         | 0.0000        |
| 2016             | 721                              | 721          | 0         | 0.0000        |
| 2017             | 314                              | 314          | 0         | 0.0000        |
| 2018             | 100                              | 100          | 0         | 0.0000        |
| <b>10 yr avg</b> | <b>817</b>                       | <b>817</b>   | <b>0</b>  | <b>0.0000</b> |
| <b>10 yr SD</b>  | <b>610</b>                       | <b>610</b>   | <b>0</b>  | <b>0.0000</b> |
| <b>20 yr avg</b> | <b>1,127</b>                     | <b>1,122</b> | <b>5</b>  | <b>0.0053</b> |
| <b>20 yr SD</b>  | <b>757</b>                       | <b>757</b>   | <b>10</b> | <b>0.0109</b> |

**Table 16. Bycatch estimates in the CNMI shore-based fishery from 2000-2018**

| Year | Shore-Based Bycatch |       |          |           |
|------|---------------------|-------|----------|-----------|
|      | # Caught            | Kept  | Released | % Bycatch |
| 2000 |                     |       |          |           |
| 2001 |                     |       |          |           |
| 2002 |                     |       |          |           |
| 2003 |                     |       |          |           |
| 2004 |                     |       |          |           |
| 2005 | 3,170               | 3,104 | 66       | 0.0208    |
| 2006 | 6,015               | 5,987 | 28       | 0.0047    |
| 2007 | 2,670               | 2,660 | 10       | 0.0037    |
| 2008 | 7,142               | 7,135 | 7        | 0.0010    |

|                  |              |              |           |               |
|------------------|--------------|--------------|-----------|---------------|
| 2009             | 4,412        | 4,411        | 1         | 0.0002        |
| 2010             | 1,839        | 1,839        | 0         | 0.0000        |
| 2011             | 2,601        | 2,601        | 0         | 0.0000        |
| 2012             | 1,466        | 1,465        | 1         | 0.0007        |
| 2013             | 2,007        | 2,001        | 6         | 0.0030        |
| 2014             | 544          | 544          | 0         | 0.0000        |
| 2015             | 687          | 687          | 0         | 0.0000        |
| 2016             | 723          | 723          | 0         | 0.0000        |
| 2017             | 1,450        | 1,447        | 3         | 0.0021        |
| 2018             | 2,326        | 2,318        | 8         | 0.0034        |
| <b>10 yr avg</b> | <b>1,806</b> | <b>1,804</b> | <b>2</b>  | <b>0.0009</b> |
| <b>10 yr SD</b>  | <b>1,095</b> | <b>1,095</b> | <b>3</b>  | <b>0.0013</b> |
| <b>20 yr avg</b> | <b>2,647</b> | <b>2,637</b> | <b>9</b>  | <b>0.0028</b> |
| <b>20 yr SD</b>  | <b>1,907</b> | <b>1,901</b> | <b>17</b> | <b>0.0052</b> |

### 1.1.10 Number of Federal Permit Holders

The Code of Federal Regulations (CFR), Title 50, Part 665 requires the following Federal permits for fishing in the exclusive economic zone (EEZ) under the Mariana FEP:

#### 1.1.10.1 Northern Mariana Island Bottomfish Permit

Regulations require this permit for any vessel commercially fishing for, landing, or transshipping bottomfish management unit species (MUS) or bottomfish ecosystem component species (ECS) in the EEZ around the Commonwealth of the Northern Mariana Islands (CNMI). Commercial fishing is also prohibited within the boundaries of the Islands Unit of the Marianas Trench Marine National Monument.

#### 1.1.10.2 Special Coral Reef Ecosystem Permit

Regulations require the coral reef ecosystem special permit for anyone fishing for coral reef ECS in a low-use marine protected area (MPA), fishing for species on the list of Potentially Harvested Coral Reef Taxa, or using fishing gear not specifically allowed in the regulations. NMFS will make an exception to this permit requirement for any person issued a permit to fish under any fishery ecosystem plan who incidentally catches CNMI coral ECS while fishing for bottomfish MUS, crustacean ECS, western Pacific pelagic MUS, precious coral, or seamount groundfish. Regulations require a transshipment permit for any receiving vessel used to land or transship potentially harvested coral reef taxa, or any coral reef ECS caught in a low-use MPA.

#### 1.1.10.3 Western Pacific Precious Corals Permit

Regulations require this permit for anyone harvesting or landing black, bamboo, pink, red, or gold corals in the EEZ in the western Pacific.

#### 1.1.10.4 Western Pacific Crustaceans Permit (Lobster or Deepwater Shrimp)

Regulations require a permit by the owner of a U.S. fishing vessel used to fish for lobster or deepwater shrimp in the EEZ around American Samoa, Guam, Hawaii, and the Pacific Remote

Islands Areas, and in the EEZ seaward of 3 nautical miles of the shoreline of the Northern Mariana Islands.

There is no record of special coral reef or precious coral fishery permits issued for the EEZ around Northern Mariana Islands since 2007. Table 17 provides the number of permits issued for CNMI fisheries between 2008 and 2018. Historical data are from the PIFSC, and 2018 data are from the PIRO Sustainable Fisheries Division permits program as of February 27, 2019.

**Table 17. Number of federal permits holders for the CNMI crustacean and bottomfish fisheries from 2009-2018**

| CNMI Fisheries | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|----------------|------|------|------|------|------|------|------|------|------|------|
| Lobster        | 4*   | 0    | 0    | 0    | 0    | 0    | 0    | 1**  | 0    | 1**  |
| Shrimp         | 0    | 2*   | 1*   | 0    | 0    | 0    | 0    | 1    | 0    | 0    |
| Bottomfish     | 2    | 13   | 10   | 13   | 5    | 6    | 7    | 17   | 20   | 13   |

\* Permits apply to multiple areas and may include American Samoa, Guam, CNMI, and PRIA.

\*\*Area 5 CNMI and Guam.

### 1.1.11 Status Determination Criteria

#### 1.1.11.1 Bottomfish Fishery

Overfishing criteria and control rules are specified and applied to individual species within the multi-species stock whenever possible. When this is not possible, they are based on an indicator species for the multi-species stock. It is important to recognize that individual species would be affected differently based on this type of control rule, and it is important that for any given species, fishing mortality does not currently exceed a level that would result in excessive depletion of that species. No indicator species are used for the bottomfish multi-species stock complexes and the coral reef species complex. Instead, the control rules are applied to each stock complex as a whole.

The MSY control rule is used as the maximum fishing mortality threshold (MFMT). The MFMT and minimum stock size threshold (MSST) are specified based on the recommendations of Restrepo et al. (1998) and both are dependent on the natural mortality rate (M). The value of M used to determine the reference point values are not specified in this document. The latest estimate, published annually in the SAFE report, is used and the value is occasionally re-estimated using the best available information. The range of M among species within a stock complex is taken into consideration when estimating and choosing the M to be used for the purpose of computing the reference point values.

In addition to the thresholds MFMT and MSST, a warning reference point,  $B_{FLAG}$ , is specified at some point above the MSST to provide a trigger for consideration of management action prior to B reaching the threshold. MFMT, MSST, and  $B_{FLAG}$  are specified as indicated in Table 18.

**Table 18. Overfishing threshold specifications for the BMUS in CNMI**

| MFMT | MSST | $B_{FLAG}$ |
|------|------|------------|
|------|------|------------|

|  |             |           |
|--|-------------|-----------|
| $F(B) = \frac{F_{MSY} B}{c B_{MSY}} \quad \text{for } B \leq c B_{MSY}$ $F(B) = F_{MSY} \quad \text{for } B > c B_{MSY}$ | $c B_{MSY}$ | $B_{MSY}$ |
| where $c = \max(1-M, 0.5)$   |             |           |

Standardized values of fishing effort (E) and catch-per-unit-effort (CPUE) are used as proxies for F and B, respectively, so  $E_{MSY}$ ,  $CPUE_{MSY}$ , and  $CPUE_{FLAG}$  are used as proxies for  $F_{MSY}$ ,  $B_{MSY}$ , and  $B_{FLAG}$ , respectively.

In cases where reliable estimates of  $CPUE_{MSY}$  and  $E_{MSY}$  are not available, they would be estimated from catch and effort time series, standardized for all identifiable biases.  $CPUE_{MSY}$  would be calculated as half of a multi-year average reference CPUE, called  $CPUE_{REF}$ . The multi-year reference window would be objectively positioned in time to maximize the value of  $CPUE_{REF}$ .  $E_{MSY}$  would be calculated using the same approach or, following Restrepo et al. (1998), by setting  $E_{MSY}$  equal to  $E_{AVE}$ , where  $E_{AVE}$  represents the long-term average effort prior to declines in CPUE. When multiple estimates are available, the more precautionary one is used.

Since the MSY control rule specified here applies to multi-species stock complexes, it is important to ensure that no particular species within the complex has a mortality rate that leads to excessive depletion. In order to accomplish this, a secondary set of reference points is specified to evaluate stock status with respect to recruitment overfishing. A secondary “recruitment overfishing” control rule is specified to control fishing mortality with respect to that status. The rule applies only to those component stocks (species) for which adequate data are available. The ratio of a current spawning stock biomass proxy ( $SSBP_t$ ) to a given reference level ( $SSBP_{REF}$ ) is used to determine if individual stocks are experiencing recruitment overfishing.  $SSBP$  is CPUE scaled by percent mature fish in the catch. When the ratio  $SSBP_t/SSBP_{REF}$ , or the “SSBP ratio” ( $SSBPR$ ) for any species drops below a certain limit ( $SSBPR_{MIN}$ ), that species is considered to be recruitment overfished and management measures will be implemented to reduce fishing mortality on that species. The rule applies only when the SSBP ratio drops below the  $SSBPR_{MIN}$ , but it will continue to apply until the ratio achieves the “SSBP ratio recovery target” ( $SSBPR_{TARGET}$ ), which is set at a level no less than  $SSBPR_{MIN}$ . These two reference points and their associated recruitment overfishing control rule, which prescribe a target fishing mortality rate ( $F_{RO-REBUILD}$ ) as a function of the SSBP ratio, are specified as indicated in Table 19. Again,  $E_{MSY}$  is used as a proxy for  $F_{MSY}$ .

**Table 19. Rebuilding control rules for the BMUS in CNMI**

| $F_{RO-REBUILD}$   | $SSBPR_{MIN}$ | $SSBPR_{TARGET}$ |
|--|---------------|------------------|
| $F(SSBPR) = 0 \quad \text{for } SSBPR \leq 0.10$<br>$F(SSBPR) = 0.2 F_{MSY} \quad \text{for } 0.10 < SSBPR \leq SSBPR_{MIN}$<br>$F(SSBPR) = 0.4 F_{MSY} \quad \text{for } SSBPR_{MIN} < SSBPR \leq SSBPR_{TARGET}$ | 0.20          | 0.30             |

### 1.1.11.2 Coral Reef Fishery

Available biological and fishery data are poor for all coral reef ecosystem management unit species in the Mariana Islands. There is scant information on the life histories, ecosystem dynamics, fishery impact, community structure changes, yield potential, and management reference points for many coral reef ecosystem species. Additionally, total fishing effort cannot be adequately partitioned between the various management unit species (MUS) for any fishery or area. Biomass, maximum sustainable yield, and fishing mortality estimates are not available for any single MUS. Once these data are available, fishery managers can establish limits and reference points based on the multi-species coral reef ecosystem as a whole.

The MSY control rule should be applied to the individual species in a multi-species stock when possible. When this is not possible, MSY may be specified for one or more species; these values can be used as indicators for the multi-species stock's MSY.

Individual species that are part of a multi-species complex will respond differently to an OY-determined level of fishing effort ( $F_{OY}$ ). Thus, for a species complex that is fished at  $F_{OY}$ , managers still must track individual species' mortality rates in order to prevent species-specific population declines that would lead to depletion.

For the coral reef fishery, the multi-species complex as a whole is used to establish limits and reference points for each area. Available data for a particular species are used to evaluate the status of individual MUS stocks in order to prevent recruitment overfishing when possible. When better data and the appropriate multi-species stock assessment methodologies become available, all stocks will be evaluated independently, without proxy.

#### Establishing Reference Point Values

Standardized values of catch per unit effort (CPUE) and effort (E) are used to establish limit and reference point values, which act as proxies for relative biomass and fishing mortality, respectively. Limits and reference points are calculated in terms of  $CPUE_{MSY}$  and  $E_{MSY}$  included in Table 20.

**Table 20. Status determination criteria for CNMI CREMUS using CPUE based proxies**

| Value                | Proxy             | Explanation                                     |
|----------------------|-------------------|---|
| MaxFMT ( $F_{MSY}$ ) | $E_{MSY}$         | $0.91 CPUE_{MSY}$                               |
| $F_{OY}$             | $0.75 E_{MSY}$    | suggested default scaling for target            |
| $B_{MSY}$            | $CPUE_{MSY}$      | operational counterpart                         |
| $B_{OY}$             | $1.3 CPUE_{MSY}$  | simulation results from Mace (1994)             |
| MinSST               | $0.7 CPUE_{MSY}$  | suggested default $(1-M)B_{MSY}$ with $M=0.3^*$ |
| $B_{FLAG}$           | $0.91 CPUE_{MSY}$ | suggested default $(1-M)B_{OY}$ with $M=0.3^*$  |

When reliable estimates of  $E_{MSY}$  and  $CPUE_{MSY}$  are not available, they are generated from time series of catch and effort values, standardized for all identifiable biases using the best available

analytical tools.  $CPUE_{MSY}$  is calculated as one-half a multi-year moving average reference CPUE ( $CPUE_{REF}$ ).

### 1.1.11.3 Current Stock Status

#### Bottomfish

Biological and other fishery data are poor for all bottomfish species in the Mariana Archipelago. Generally, data are only available on commercial landings by species and catch-per-unit-effort (CPUE) for the multi-species complexes as a whole. At this time it is not possible to partition these effort measures among the various bottomfish MUS. The most recent stock assessment update (Yau et al., 2015) for the CNMI bottomfish management unit species complex (comprised of 17 species of shallow and deep species of snapper, grouper, jacks, and emperors) was based on estimate of total catch, an abundance index derived from the nominal CPUE generated from the creel surveys, and a fishery independent point estimate of MSY from the Our Living Oceans Report (Humphreys and Moffitt, 1999; Moffitt and Humphreys, 2009). The assessment utilized a state-space surplus production model with explicit process and observation error terms (Meyer and Millar, 1999). Determinations of overfishing and overfished status can then be made by comparing current biomass and harvest rates to MSY level reference points. To date, the CNMI BMUS is not subject to overfishing and is not overfished (Table 21).

**Table 21. Stock assessment parameters for the CNMI BMUS complex (Yau et al., 2015).**

| Parameter   | Value             | Notes                                       | Status                   |
|-------------|-------------------|---|--------------------------|
| MSY         | $173.1 \pm 32.19$ | Expressed in 1000 lbs. ( $\pm$ std. error)  |                          |
| $H_{2013}$  | 0.022             | Expressed in percentage                     |                          |
| $H_{MSY}$   | $0.261 \pm 0.063$ | Expressed in percentage ( $\pm$ std. error) |                          |
| $H/H_{MSY}$ | 0.088             |   | No overfishing occurring |
| $B_{2013}$  | 1,262             | Expressed in thousand pounds                |                          |
| $B_{MSY}$   | $683.5 \pm 126.7$ | Expressed in 1000 lbs. ( $\pm$ std. error)  |                          |
| $B/B_{MSY}$ | 1.85              |   | Not overfished           |

#### Coral Reef

The application of the SDCs for the management unit species in the coral reef fisheries is limited due to various challenges. First, the thousands of species included in the coral reef MUS makes the SDC and status determination impractical. Second, the CPUE derived from the creel survey is based on the fishing method and there is no species-specific CPUE information available. In order to allocate the fishing method level CPUE to individual species, the catch data (the value of catch is derived from CPUE hence there is collinearity) will have to be identified to species level and CPUE will be parsed out by species composition. The third challenge is that there is very little species level identification applied to the creel surveys. There has been no attempt to estimate MSY for the coral reef MUS until the 2007 re-authorization of MSA that requires the Council to specify ACLs for species in the FEPs.



For ACL specification purposes, MSYs in the coral reef fisheries are determined by using the Biomass-Augmented Catch-MSY approach (Sabater and Kleiber, 2014). This method estimates MSY using plausible combination rates of population increase (denoted by  $r$ ) and carrying capacity (denoted by  $k$ ) assumed from the catch time series, resilience characteristics (from FishBase), and biomass from existing underwater census surveys done by PIFSC. This method was applied to species complexes grouped by taxonomic families. The most recent MSY estimates are found in Table 22. The SSC utilized the MSYs for the coral reef MUS complexes as the OFLs.

**Table 22. Best available MSY estimates for the coral reef MUS in CNMI**

| <b>Coral Reef MUS Complex</b>  | <b>MSY (lbs.)</b> |
|--|-------------------|
| <i>Selar crumenophthalmus</i> – atulai or bigeye scad  | 122,500           |
| Acanthuridae – surgeonfish   | 361,200           |
| Carangidae – jacks   | 55,300            |
| Crustaceans – crabs  | 9,100             |
| Holocentridae – squirrelfish   | 78,500            |
| Kyphosidae – chubs/rudderfish  | 29,500            |
| Labridae – wrasses <sup>1</sup>  | 73,500            |
| Lethrinidae – emperors   | 69,700            |
| Lutjanidae – snappers  | 225,800           |
| Mollusks – turbo snail; octopus; giant clams   | 16,700            |
| Mugilidae – mullets  | 7,700             |
| Mullidae – goatfish  | 31,000            |
| Scaridae – parrotfish <sup>2</sup>   | 189,900           |
| Serranidae – groupers  | 110,300           |
| Siganidae – rabbitfish   | 12,000            |
| All Other CREMUS Combined (other coral reef ecosystem finfish, other invertebrates, misc. bottomfish, misc. reef fish, and misc. shallow bottomfish) | 14,500            |
| <i>Cheilinus undulatus</i> – humphead (Napoleon) wrasse  | N.A.              |
| <i>Bolbometopon muricatum</i> – bumphead parrotfish  | N.A.              |
| Carcharhinidae – reef sharks   | N.A.              |

## 1.1.12 Overfishing Limit, Acceptable Biological Catch, and Annual Catch Limits

### 1.1.12.1 Brief Description of the ACL Process

The Council developed a Tiered system of control rules to guide the specification of ACLs and Accountability Measures (AMs) (WPRFMC, 2011). The process starts with the use of the best scientific information available (BSIA) in the form of, but not limited to, stock assessments, published paper, reports, or available data. These information are classified to the different Tiers in the control rule ranging from Tier 1 (most information available typically an assessment) to Tier 5 (catch-only information). The control rules are applied to the BSIA. Tiers 1 to 3 would involve conducting a Risk of Overfishing Analysis (denoted by P\*) to quantify the scientific uncertainties around the assessment to specify the Acceptable Biological Catch (ABC). This

would lower the ABC from the OFL (MSY-based). A Social, Ecological, Economic, and Management (SEEM) Uncertainty Analysis is performed to quantify the uncertainties from the SEEM factors. The buffer is used to lower the ACL from the ABC. For Tier 4, which contains stocks with MSY estimates but no active fisheries, the control rule is 91% of MSY. For Tier 5, which contains catch only information, the control rule is a third reduction in the median catch depending on the qualitative evaluation on what the stock status is based on expert opinion. ACL specification can choose from a variety of method including the above mentioned SEEM analysis or a percentage buffer (percent reduction from ABC based on expert opinion) or the use of an Annual Catch Target. Specifications are done on an annual basis but the Council normally specifies a multi-year specification.

The Accountability Measure for the coral reef and bottomfish fisheries in CNMI is an overage adjustment. The ACL is downward adjusted with the amount of overage from the ACL based on a three year running average.

### 1.1.12.2 Current OFL, ABC, ACL, and Recent Catch

The most recent multiyear specification of OFL, ABC, and ACL for the coral reef fishery was completed in the 160<sup>th</sup> Council meeting on June 25 to 27, 2014. The specification covers fishing year 2015, 2016, 2017, and 2018 for the coral reef MUS complexes. A P\* and SEEM analysis was performed for this multiyear specification (NMFS, 2015). For the bottomfish, it was a roll over from the previous specification since an assessment update was not available for fishing year 2015. ACLs were not specified by NMFS for the coral reef ecosystem MUS because NMFS has recently acquired new information that require additional environmental analyses to support the Council's ACL recommendations for these management unit species (50 CFR Part 665).

**Table 23. CNMI ACL table with three-year average catch in pounds**

| <b>Fishery</b> | <b>MUS</b>                        | <b>OFL</b> | <b>ABC</b> | <b>ACL</b> | <b>Catch</b> |
|----------------|-----------------------------------|------------|------------|------------|--------------|
| Bottomfish     | Bottomfish multi-species complex  | 293,000    | 228,000    | 228,000    | 32,370       |
| Crustacean     | Deepwater shrimp                  | N.A.       | 275,570    | 275,570    | N.A.F        |
|                | Spiny lobster                     | 9,600      | 7,800      | 7,410      | 829          |
|                | Slipper lobster                   | N.A.       | 60         | 60         | <b>172</b>   |
|                | Kona crab                         | N.A.       | 6,300      | 6,300      | N.A.F        |
| Precious Coral | Black coral                       | 8,250      | 2,100      | 2,100      | N.A.F        |
|                | Precious coral in CNMI expl. area | N.A.       | 2,205      | 2,205      | N.A.F        |

Notes: The MUS highlighted in red have a three-year recent average catch that exceeds the prescribed ACL.

The catch shown in Table 23 takes the average of the recent three years as recommended by the Council at its 160<sup>th</sup> meeting to avoid large fluctuations in catch due to data quality and outliers. "N.A.F." indicates no active fisheries to date. "N.D." indicates that there are no data available.

### 1.1.13 Best Scientific Information Available

#### 1.1.13.1 Bottomfish Fishery

##### Stock Assessment Benchmark

The benchmark stock assessment for the Territory Bottomfish Management Unit Species complex was developed and finalized in October 2007 (Moffitt et al., 2007). This benchmark utilized a Bayesian statistical framework to estimate parameters of a Schaefer model fit to a time series of annual CPUE statistics. The surplus production model included process error in biomass production dynamics and observation error in the CPUE data. This was an improvement to the previous approach of using index-based proxies for  $B_{MSY}$  and  $F_{MSY}$ . Best available information for the bottomfish stock assessment is as follows:

Input data: The CPUE and catch data used were from the Guam off-shore creel survey. The catch and CPUE were expanded on annual level. CPUE was expressed in line-hours. The data was screened for trips that landed more than 50% BMUS species using the handline gear.

Model: state-space model with explicit process and observation error terms (see Meyer and Millar, 1999).

Fishery independent source for biomass: point estimate of MSY from the Our Living Oceans Report (Humphreys and Moffitt, 1999; Moffitt and Humphreys, 2009)

##### Stock Assessment Updates

Updates to the 2007 benchmark done in 2012 (Brodziak et al., 2012) and 2015 (Yau et al., 2015). These included a two-year stock projection table used for selecting the level of risk the fishery will be managed under ACLs. Yau et al. (2015) is considered the best scientific information available for the Territory bottomfish MUS complex after undergoing a WPSAR Tier 3 panel review (Franklin et al., 2015). This was the basis for the P\* analysis and SEEM analysis the determined the risk levels to specify ABCs and ACLs.

##### Other Information Available

Approximately every five years PIFSC administers a socioeconomic survey to small boat fishermen in CNMI. This survey consists of about 60 questions regarding a variety of topics, including fishing experiences, market participation, vessels and gear, demographics and household income, and fishermen perspectives. The survey requests participants to identify which MUS they primarily targeted during the previous 12 months, by percentage of trips. Full reports of these surveys can be found at the PIFSC Socioeconomics webpage (Hospital and Beavers, 2011).

#### 1.1.13.2 Coral Reef Fishery

##### 1.1.13.2.1 Stock Assessment Benchmark

No stock assessment has been generated for the coral reef fisheries. The SDCs using index-based proxies were tested for its applicability in the different MUS in the coral reef fisheries (Hawhee, 2007). This analysis was done on a gear level. It paints a dire situation for the shore-based fishery with 43% of the gear/species combination fell below  $B_{flag}$  and 33% below MSST with most catch and CPUE trends showing a decline over time. The off-shore fisheries were shown to be less dire with 50% of the gear/species combination fell below  $B_{flag}$  and 38% below MSST but

the catch and CPUE trends were increasing over time. The inconsistency in the CPUE and catch trends with the SDC results makes this type of assessment to be unreliable.

The first attempt to use a model based approach in assessing the coral reef MUS complexes was done in 2014 using a biomass-based population dynamics model (Sabater and Kleiber, 2014). This model was based on the original Martell and Froese (2012) model but was augmented with biomass information to relax the assumption behind carrying capacity. It estimates MSY based on a range of rate of population growth ( $r$ ) and carrying capacity ( $k$ ) values. The best available information for the coral reef stock assessment is as follows:

**Input data:** The catch data was derived from the inshore and off-shore creel surveys. Commercial receipt book information was also used in combination of the creel data. A downward adjustment was done to address for potential overlap due to double reporting.

**Model:** Biomass Augmented Catch MSY approach based on the original catch-MSY model (Martell and Froese, 2013; Sabater and Kleiber, 2014).

**Fishery independent source for biomass:** biomass density from the Rapid Assessment and Monitoring Program (RAMP) of NMFS' Coral Reef Ecosystem Division (CRED) was expanded to the hard bottom habitat from 0-30 m (Williams, 2010).

This model had undergone a CIE review in 2014 (Cook, 2014; Haddon, 2014; Jones, 2014). This was the basis for the P\* analysis that determined the risk levels to specify ABCs

#### **1.1.13.2.2 Stock Assessment Updates**

No updates available for the coral reef MUS complex. However, NMFS-PIFSC is finalizing a length-based model for estimating sustainable yield levels and various biological reference points (Nadon et al., 2015). This can be used on a species level. The Council is also working with a contractor to enhance the BAC-MSY model to incorporate catch, biomass, CPUE, effort, length-based information in an integrated framework (Martell 2015)

#### **1.1.13.2.3 Other Information Available**

Approximately every five years PIFSC administers a socioeconomic survey to small boat fishermen in CNMI. This survey consists of about 60 questions regarding a variety of topics, including fishing experiences, market participation, vessels and gear, demographics and household income, and fishermen perspectives. The survey requests participants to identify which MUS they primarily targeted during the previous 12 months, by percentage of trips. Full reports of these surveys can be found at the PIFSC Socioeconomics webpage (Hospital and Beavers, 2011).

PIFSC and the Council conducted a workshop with various stakeholders in CNMI to identify factors and quantify uncertainties associated with the social, economic, ecological, and management of the coral reef fisheries (Sievanen and McCaskey 2014). This was the basis for the SEEM analysis that determined the risk levels to specify ACLs.

### 1.1.14 Harvest Capacity and Extent

The MSA defines the term “optimum,” with respect to the yield from a fishery, as the amount of fish which:

- Will provide the greatest overall benefit to the Nation, particularly with respect to food production and recreational opportunities, and taking into account the protection of marine ecosystems.
- Is prescribed on the basis of the MSY from the fishery, as reduced by any relevant social, economic, or ecological factor.
- In the case of an overfished fishery, provides for rebuilding to a level consistent with producing the MSY in such fishery [50 CFR §600.310(f)(1)(i)].

Optimum yield in the coral reef and bottomfish fisheries is prescribed based on the MSY from the stock assessment and the best available scientific information. In the process of specifying ACLs, social, economic, and ecological factors were considered and the uncertainties around those factors defined the management uncertainty buffer between the ABC and ACL. OY for the bottomfish and coral reef fish MUS complexes is defined to be the level of harvest equal to the ACL consistent with the goals and objectives of the Fishery Ecosystem Plans and used by the Council to manage the stock.

The Council recognizes that MSY and OY are long term values whereas the ACLs are yearly snapshots based on the level of fishing mortality at  $F_{MSY}$ . There are situations when the long-term means around MSY are going to be lower than ACLs especially if the stock is known to be productive or relatively pristine or lightly fished. One can have catch levels and catch rates exceeding that of MSY over short-term enough to lower the biomass to a level around the estimated MSY and still not jeopardize the stock. In this situation is true for the territory bottomfish multi-species complex.

The harvest extent, in this case, is defined as the level of catch harvested in a fishing year relative to the ACL or OY. The harvest capacity is the level of catch remaining in the annual catch limit that can potentially be used for the total allowable level of foreign fishing (TALFF).

Table 24 summarizes the harvest extent and harvest capacity information for CNMI in 2017.

**Table 24. CNMI proportion of harvest capacity and extent for 2018**

| Fishery        | MUS                               | ACL     | Catch  | Harvest extent (%) | Harvest capacity (%) |
|----------------|-----------------------------------|---------|--------|--------------------|----------------------|
| Bottomfish     | Bottomfish multi-species complex  | 228,000 | 32,370 | 14.2               | 85.8                 |
| Crustacean     | Deepwater shrimp                  | 275,570 | N.A.F. | 0.0                | 100.0                |
|                | Spiny lobster                     | 7,410   |        |                    |                      |
|                | Slipper lobster                   | 60      | 172    | 286                | 0.0                  |
|                | Kona crab                         | 6,300   | N.A.F. | 0.0                | 100.0                |
| Precious coral | Black coral                       | 2,100   | N.A.F. | 0.0                | 100.0                |
|                | Precious coral in CNMI expl. area | 2,205   | N.A.F. | 0.0                | 100.0                |

### 1.1.15 Administrative and Regulatory Actions

This summary describes management actions NMFS has taken for CNMI fisheries since the April 2018 Joint FEP Plan Team meeting.

June 14, 2018. Final rule. 5-Year Extension of Moratorium on Harvest of Gold Corals. This final rule extends the region-wide moratorium on the harvest of gold corals in the U.S. Pacific Islands through June 30, 2023. NOAA Fisheries intends this final rule to prevent overfishing and to stimulate research on gold corals.

February 8, 2019. Final rule. Reclassifying Management Unit Species to Ecosystem Component Species. This final rule reclassifies certain management unit species in the Pacific Islands as ecosystem component species. The rule also updates the scientific and local names of certain species. The intent of this final rule is to prioritize conservation and management efforts and to improve efficiency of fishery management in the region. This rule is effective March 11, 2019.

### 1.1.16 References

- Bowers, N.M., 2001. Problems of resettlement on Saipan, Tinian, and Rota, Mariana Islands. Occasional Historical Papers Series No. 7. CNMI Division of Historic Preservation, Saipan, MP.
- Cook, R., 2014. Report on the Review of the Biomass Augmented Catch-MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report to Center for Independent Experts.
- Fricke, H. and Meischner, D., 1985. Depth limits of Bermudan scleractinian corals: a submersible survey. *Marine Biology*, 88(2), pp.175-187.
- Haddon, M., 2014. Center for Independent Experts (CIE) Peer Review of the Biomass Augmented Catch-MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report submitted to the Center for Independent Experts.
- Hart, S.R., Coetzee, M., Workman, R.K., Blusztajn, J., Johnson, K.T.M., Sinton, J.M., Steinberger, B. and Hawkins, J.W., 2004. Genesis of the Western Samoa seamount province: age, geochemical fingerprint and tectonics. *Earth and Planetary Science Letters*, 227(1), pp.37-56.
- Humphreys, R. and Moffitt, R., 1999. Unit 17-Western Pacific Bottomfish and Armorhead Fisheries. DOC, NOAA, NMFS Our Living Oceans—Report on the Status of US Living Marine Resources, pp.189-192.
- Jones, C., 2014. Biomass Augmented Catch---MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report submitted to the Center for Independent Experts.
- Kahng, S.E. and Maragos, J.E., 2006. The deepest, zooxanthellate scleractinian corals in the world? *Coral Reefs*, 25(2), pp.254-254.

- Konter, J.G. and Jackson, M.G., 2012. Large volumes of rejuvenated volcanism in Samoa: Evidence supporting a tectonic influence on late-stage volcanism. *Geochemistry, Geophysics, Geosystems*, 13(6).
- Lang, J.C., 1974. Biological Zonation at the Base of a Reef: Observations from the submersible Nekton Gamma have led to surprising revelations about the deep fore-reef and island slope at Discovery Bay, Jamaica. *American Scientist*, 62(3), pp.272-281.
- Martell, S. and Froese, R., 2013. A simple method for estimating MSY from catch and resilience. *Fish and Fisheries*, 14(4), pp.504-514.
- Meyer, R. and Millar, R.B., 1999. Bayesian stock assessment using a state-space implementation of the delay difference model. *Canadian Journal of Fisheries and Aquatic Sciences*, 56(1), pp.37-52.
- Moffitt, R.B., Brodziak, J., and Flores, T., 2007. Status of the Bottomfish Resources of American Samoa, Guam, and Commonwealth of the Northern Mariana Islands, 2005. NOAA Pacific Islands Fisheries Science Center Administrative Report H-07-04.
- Nadon, M.O., Ault, J.S., Williams, I.D., Smith, S.G. and DiNardo, G.T., 2015. Length-based assessment of coral reef fish populations in the main and northwestern Hawaiian Islands. *PloS one*, 10(8), p.e0133960.
- Neall, V.E. and Trewick, S.A., 2008. The age and origin of the Pacific islands: a geological overview. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1508), pp.3293-3308.
- NMFS, 2015. Specification of Annual Catch Limits and Accountability Measures for Pacific Island Coral Reef Ecosystem Fisheries in Fishing Years 2015 through 2018. Honolulu, HI 96813. 228 p.
- Oram, R., TuiSamoa, N., Tomanogi, J., Sabater, M., Quach, M., Hamm, D., and Graham, C., 2011. American Samoa Shore-Based Creel Survey Documentation. NOAA, National Marine Fishery Service, Pacific Island Fishery Science Center.
- Oram, R., TuiSamoa, N., Tomanogi, J., Sabater, M., Quach, M., Hamm, D., and Graham, C., 2011. American Samoa Boat-Based Creel Survey Documentation. NOAA, National Marine Fishery Service, Pacific Island Fishery Science Center.
- Restrepo, V.R., Thompson, G.G., Mace, P.M., Gabriel, W.L., Low, L.L., MacCall, A.D., Methot, R.D., Powers, J.E., Taylor, B.L., Wade, P.R. and Witzig, J.F., 1998. Technical guidance on the use of precautionary approaches to implementing National Standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. NOAA Technical Memorandum NMFS-F/SPO-31, pp.1-54.
- Russell, S., 1999. Tiempón Alemán: A Look Back at German Rule of the Northern Mariana Islands, 1899-1914. University of Guam, Micronesian Area Research Center.

- Sabater, M. and Kleiber, P., 2014. Augmented catch-MSY approach to fishery management in coral-associated fisheries. *Interrelationships between Corals and Fisheries*. CRC Press, Boca Raton, FL, pp.199-218.
- Williams, I., 2010. U.S. Pacific Reef Fish Estimates Based on Visual Survey Data. NOAA Pacific Islands Fisheries Science Center Internal Report IR-10-024.
- WPRFMC, 2011. Omnibus Amendment for the Western Pacific Region to Establish a Process for Specifying Annual Catch Limits and Accountability Measures. Honolulu, HI 96813.
- Yau, A., Nadon, M., Richards, B., Brodziak, J., Fletcher, E., 2016. Stock assessment updates of the Bottomfish Management Unit species of American Samoa, the Commonwealth of the Northern Mariana Islands, and Guam in 2015 using data through 2013. U.S. Dept. of Commerce, NOAA Technical Memorandum, NMFS-PIFSC-51, 54 p.

## 1.2 GUAM FISHERY DESCRIPTIONS

### 1.2.1 Bottomfish Fishery

Bottomfishing on Guam is a combination of recreational, subsistence, and small-scale commercial fishing. It can be separated into two distinct fisheries separated by depth and species composition. The shallow water complex (< 500 feet) comprises the largest portion of the total bottomfish harvest and effort, and primarily includes: reef-dwelling snappers of the genera *Lutjanus*, *Aphareus*, and *Aprion*; groupers of the genera *Epinephelus*, *Variola*, and *Cephalopholis*; jacks of the genera *Caranx* and *Carangoides*; Holocentrids (*Myripristis* spp. and *Sargocentron* spp.); emperors of the genera *Lethrinus* and *Gymnocranius*; and Dogtooth Tuna (*Gymnosarda unicolor*). The deep water complex (> 500 feet) consists primarily of groupers of the genera *Hyporthodus* and *Cephalopholis*, jacks of the genera *Caranx* and *Seriola*, and snappers of the genera *Pristipomoides*, *Etelis*, and *Aphareus*. In recent years, deep water species have made up a significant portion of the total expanded bottomfishing catch.

The majority of people that participate in the bottomfish fishery are either subsistence or part-time commercial fishermen, operate boats less than 25 feet in length, and target primarily the shallow water bottomfish complex. It is not uncommon to intercept fishermen combining bottomfishing with other methods such as trolling, spearing, and jigging to maximize their catch. High demand has made it profitable to sell locally caught bottomfish, although overhead costs including fuel and gear may be significant factors for in determining a fisherman's selection of fishing method. The demand for local bottomfish, when combined with environmental pressures, however, may cause stress to local bottomfish stocks.

The majority of bottomfishing around Guam takes place on offshore banks, though practically no information exists on the condition of the reefs on offshore banks. On the basis of anecdotal information, most of the offshore banks are in good condition due to their isolation. According to Myers (1997), less than 20 percent of the total coral reef resources harvested in Guam are taken from the EEZ, primarily because the reefs are often associated with less accessible offshore banks. As such, finfish make up most of the catch in the EEZ. Most offshore banks are deep, remote, and subject to strong currents. Generally, these banks are only accessible during calm weather in the summer months (May to August/September). Galvez Bank is the closest and most



accessible and, consequently, fished most frequently. In contrast, other banks (White Tuna and Santa Rose, Rota) are remote and generally are fished only during exceptional weather conditions (Green 1997). Local fishermen report that up to ten commercial boats, with two to three people per boat, and some recreational boats, make use of the banks when the weather is good (Green 1997).

At present, the banks are fished using two methods: bottomfishing by hook and line, and jigging at night for bigeye scad (*Selar crumenophthalmus*; Myers 1997). In recent years, the estimated annual catch in these fisheries has ranged from 14 to 22 metric tons of shallow bottomfish and 3 to 15 metric tons of bigeye scad (Green 1997). The shallow water component accounted for nearly 68 percent (35,002 to 65,162 lbs.) of the aggregate bottomfish landings in fiscal years 1992–1994 (Myers 1997). Catch composition of the shallow water bottomfish complex (and coral reef species) is dominated by lethrinids, with a single species (*Lethrinus rubrioperculatus*) alone accounting for 28 percent of the total catch. Other important components of the bottomfish catch include lutjanids, carangids, other lethrinids, and serranids. Holocentrids, mullids, labrids, scombrids, and balistids are minor components of the shallow water bottomfish complex. It should be noted that at least two of these species (*Aprion virescens* and *Caranx lugubris*) are also found in deeper waters, and as a result comprise a portion of the catch of the deep water fishery.

Species that are commonly taken in the shallow-bottom fishery of Guam are:

*Aphareus furca*

*Aprion virescens*

*Lutjanus kasmira*, *L. fulvus*

*Carangoides orthogrammus*

*Caranx lugubris*, *C. melampygus*, *C. ignobilis*

*Selar crumenophthalmus*

*Cephalopholis argus*, *C. spiloparaea*, *C. urodeta*

*Epinephelus fasciatus*

*Gymnocranius* spp.

*Lethrinus atkinsoni*, *L. erythracanthus*, *L. olivaceus*, *L. rubrioperculatus*,

*L. xanthochilus*

*Gymnosarda unicolor*

*Sargocentron* spp.

*Myripristis* spp.

*Variola albimarginata*, *V. louti*

Species that are commonly taken in the deep-bottom fishery of Guam are:

*Aphareus rutilans*

*Aprion virescens*

*Caranx lugubris*

*Seriola dumerilii*

*Cephalopholis igarashiensis*, *C. sonnerati*

*Hyporthodus octofasciatus*

*Etelis carbunculus*, *E. coruscans*  
*Pristipimoides* spp.

### 1.2.2 Coral Reef Fishery

Shore-based fishing accounts for most of the fish and invertebrate harvest from coral reefs around Guam. The coral reef fishery harvests more than 100 species of fish, including members of the families Acanthuridae, Carangidae, Gerreidae, Holocentridae, Kyphosidae, Labridae, Lethrinidae, Lutjanidae, Mugilidae, Mullidae, Scaridae, and Siganidae (Hensley and Sherwood 1993). There are several pulse fisheries for juvenile fish that can be major components of the coral reef fishery, but totals in these can vary year to year. These include juvenile rabbitfish (manahak and lessó'), juvenile jacks (i'e), and juvenile goatfish (ti'ao).

Species that are commonly taken in the coral reef fishery of Guam are:

*Naso unicornis*, *N. lituratus*  
*Acanthurus xanthopterus*, *A. lineatus*, *A. triostegus*  
*Caranx melampygus*, *C. papuensis*, i'e  
*Selar crumenophthalmus*  
*Gerres acinaces*  
*Myripristis* spp.  
*Sargocentron* spp.  
*Neoniphon* spp.  
*Kyphosus cinerascens*, *K. vaigiensis*  
*Cheilinus undulatus*, *Cheilinus* spp., *Halichoeres* spp.  
*Lethrinus harak*, *L. obseletus*, *L. atkinsoni*, *Gnathodentex aurolineatus*  
*Lutjanus fulvus*, *L. monostigma*, *L. bohar*, *L. argentimaculatus*  
*Mulloidichthys flavolineatus*, *M. vanicolensis*, ti'ao  
*Parupeneus multifasciatus*, *P. barberinus*, *P. cyclostomus*  
*Ellechelon vaigiensis*, *Moolgarda engeli*, *M. seheli*  
*Chlorurus spilurus*, *C. frontalis*,  
*Scarus psittacus*, *S. altipinnis*, *S. rubrioviolaceus*, *S. ghobban*, *S. schlegeli*  
*Siganus spinus*, *S. argenteus*, manahak, lessó

Hook and line is the most common method of fishing for coral reef fish on Guam. In 2018, hook and line fishing accounted for around 75% of fishers and gear. Throw net (talaya) is the second most common method, accounting for about 10% of fishers and gear. Other methods include gill net, snorkel spearfishing, SCUBA spearfishing, surround net, drag net, hooks and gaffs, and gleaning.

### 1.2.3 Fishery Data Collection System

Guam currently has three fishery-dependent collection programs which can be described as long-term data collection programs with different approaches for gathering important information on fishery harvest methods performed by fishermen. The three programs are the offshore data

program, the inshore data program, and the commercial fishery program. The Sportfish Restoration Grant from the U.S. Fish and Wildlife Service provides the significant portion of the funding for these programs. Training of the fishery staff to collect information is rigorous, and year-end totals are calculated by an expansion process done with in collaboration with NOAA PIFSC. Identification of fish to the species level is the goal of Guam's fishery staff.

The offshore and inshore programs, boat- and shore-based creel surveys, respectively, are long-term programs that collect participation, effort, and catch data from fishermen. Collaboration with PIFSC has resulted in a reproducible computer database program that can analyze the data to produce various types of trends that describe status of both charter and non-charter fisheries in federal and local waters. The commercial receipt book program is an important source of information for fish that enter the commercial market; however, obtaining information from dealers has been sporadic, occasionally with less than three dealers providing data. In order to improve this situation, the Council, DAWR, and PIFSC partnered to increase vendor participation in the data collection program through the Territory Science Initiative.

Guam has continued to experience high levels of commercial activity targeting reef fish. This has primarily been performed by recent migrants from the Federated States of Micronesia. The fishers are generally hired by retail shops to fish six days per week; there have been as many as eight or nine of these stores open at a time. Gathering commercial sales data from these vendors has been difficult due to vendor anxiety surrounding the reason data is being collected and the lack of perceived benefit to the vendor for reporting sales. There have been several instances during data collection where the vendors were not able to comfortably communicate in English. Data collected from these vendors is of limited value, as fish are not identified to species level, and are frequently labeled simply as "reef fish". In 2018, there were seven vendors reporting sales. In order to improve this situation, the Council, DAWR, and PIFSC partnered to increase vendor participation in the data collection program through the Territory Science Initiative. Extensive training, follow-ups, education, and outreach efforts were conducted to vendors and fishermen to increase participation in data collection.

Oram et al. (2011a) and (2011b) describe the fishery data collection process for the offshore and inshore programs on Guam. In general, DAWR staff collect fishery information through a series of random-stratified surveys for participation (i.e. accounting for fishing effort) and catch interviews (i.e. accounting for catch composition, size frequency, and catch-per-unit effort, CPUE). These data are transcribed into the WPacFIN database, and the annual catch estimates are expanded from the effort and CPUE information. Monthly commercial vendor reports are tallied at the end of the year and adjusted based on the coverage estimates provided by the vendor and/or the data collection program staff.

#### **1.2.4 Meta-Data Dashboard Statistics**

The meta-data dashboard statistics describe the amount of data used or available to calculate the fishery-dependent information. Creel surveys are sampling-based systems that require random-stratified design applied to pre-scheduled surveys. The number of sampling days, participation runs, and catch interviews would determine if there are sufficient samples to run the expansion algorithm. The trends of these parameters over time may infer survey performance. Monitoring the survey performance is critical for explaining the reliability of the expanded information.

Commercial receipt book information depends on the amount of invoices submitted and the number of vendors participating in the program. Variations in these meta-data affect the commercial landing and revenue estimates.

### 1.2.5 Creel Survey Meta-Data Statistics

#### Calculations: Shore-based data

# Interview Days: Count of the number of actual days that Creel Survey Data were collected. It's a count of the number of unique dates found in the interview sampling data (the actual sampling date data, include opportunistic interviews).

# Participation Runs: Count of the number of unique occurrences of the day/night shift combined with surveyor's initials (the person assigned to conduct the participation survey on a given date). This is compiled annually from the participation header data.

# Catch Interviews: Count of the total number of data records found in the interview header data (number of interview headers). This is divided into two categories, interviews conducted during scheduled survey days (Regular), and opportunistic interviews (Opportunistic) which are collected on non-scheduled days.

#### Calculation: Boat-based data

# Sample days: Count of the total number of unique dates found in the boat log data sampling date data.

# Catch Interviews: Count of the total number of data records found in the interview header data (number of interview headers). This is divided into two categories, interviews conducted during scheduled survey days (Regular), and opportunistic interviews (Opportunistic) which are collected on non-scheduled days.

**Table 25. Summary of Guam creel survey meta-data describing survey performance parameters with potential influence on the creel survey expansion from 1982-2018**

| Year | Shore-Based      |                      |                    |               | Boat-Based    |                    |               |
|------|------------------|----------------------|--------------------|---------------|---------------|--------------------|---------------|
|      | # Interview Days | # Participation Runs | # Catch Interviews |               | # Sample Days | # Catch Interviews |               |
|      |                  |                      | Regular            | Opportunistic |               | Regular            | Opportunistic |
| 1982 |                  |                      |                    |               | 46            | 469                | 8             |
| 1983 |                  |                      |                    |               | 47            | 431                | 34            |
| 1984 | 12               | 23                   | 56                 | 0             | 53            | 531                | 0             |
| 1985 | 51               | 78                   | 367                | 0             | 66            | 812                | 0             |
| 1986 | 47               | 74                   | 291                | 0             | 49            | 522                | 0             |
| 1987 | 45               | 62                   | 245                | 0             | 48            | 612                | 0             |
| 1988 | 48               | 62                   | 280                | 0             | 48            | 949                | 0             |
| 1989 | 49               | 63                   | 297                | 0             | 48            | 931                | 2             |
| 1990 | 47               | 62                   | 485                | 0             | 48            | 1,028              | 0             |

|                  |           |           |            |           |           |            |          |
|------------------|-----------|-----------|------------|-----------|-----------|------------|----------|
| 1991             | 48        | 54        | 497        | 0         | 48        | 1,019      | 1        |
| 1992             | 48        | 55        | 611        | 0         | 48        | 1,110      | 0        |
| 1993             | 48        | 48        | 598        | 0         | 52        | 1,119      | 0        |
| 1994             | 47        | 48        | 702        | 0         | 55        | 1,168      | 0        |
| 1995             | 48        | 49        | 764        | 0         | 96        | 1,613      | 4        |
| 1996             | 48        | 53        | 679        | 0         | 96        | 1,608      | 0        |
| 1997             | 48        | 67        | 915        | 0         | 96        | 1,358      | 0        |
| 1998             | 49        | 73        | 880        | 0         | 96        | 1,581      | 0        |
| 1999             | 48        | 68        | 939        | 1         | 96        | 1,367      | 3        |
| 2000             | 48        | 84        | 791        | 0         | 96        | 1,246      | 1        |
| 2001             | 48        | 96        | 753        | 0         | 96        | 908        | 6        |
| 2002             | 47        | 94        | 439        | 4         | 84        | 610        | 1        |
| 2003             | 48        | 96        | 518        | 10        | 78        | 446        | 0        |
| 2004             | 47        | 93        | 337        | 35        | 95        | 530        | 1        |
| 2005             | 48        | 96        | 371        | 3         | 97        | 552        | 0        |
| 2006             | 49        | 96        | 300        | 0         | 96        | 556        | 0        |
| 2007             | 48        | 96        | 243        | 118       | 96        | 500        | 0        |
| 2008             | 46        | 96        | 282        | 0         | 96        | 571        | 2        |
| 2009             | 47        | 94        | 321        | 1         | 96        | 803        | 0        |
| 2010             | 48        | 94        | 299        | 0         | 96        | 902        | 0        |
| 2011             | 43        | 96        | 250        | 0         | 96        | 645        | 0        |
| 2012             | 47        | 92        | 272        | 0         | 74        | 371        | 0        |
| 2013             | 49        | 94        | 257        | 0         | 96        | 561        | 1        |
| 2014             | 48        | 92        | 227        | 0         | 90        | 635        | 9        |
| 2015             | 45        | 96        | 279        | 46        | 97        | 651        | 13       |
| 2016             | 48        | 96        | 281        | 9         | 93        | 900        | 2        |
| 2017             | 45        | 92        | 245        | 1         | 92        | 820        | 10       |
| 2018             | 46        | 94        | 221        | 1         | 89        | 795        | 11       |
| <b>10 yr avg</b> | <b>47</b> | <b>94</b> | <b>265</b> | <b>6</b>  | <b>92</b> | <b>708</b> | <b>5</b> |
| <b>10 yr SD</b>  | <b>2</b>  | <b>2</b>  | <b>30</b>  | <b>14</b> | <b>7</b>  | <b>159</b> | <b>5</b> |
| <b>20 yr avg</b> | <b>47</b> | <b>93</b> | <b>381</b> | <b>11</b> | <b>92</b> | <b>718</b> | <b>3</b> |
| <b>20 yr SD</b>  | <b>1</b>  | <b>6</b>  | <b>203</b> | <b>27</b> | <b>6</b>  | <b>248</b> | <b>4</b> |

### 1.2.5.1 Commercial receipt book statistics

#### Calculations:

# Vendors: Count of the number of unique buyer codes found in the commercial purchase header data from the Commercial Receipt Book.

# Invoices: Count of the number of unique invoice numbers found in the commercial header data from the Commercial Receipt Book.

**Table 26. Summary of Guam commercial receipt book meta-data describing reporting performance parameters with potential influence on total commercial landing estimates from 1980-2018**

| <b>Year</b> | <b>Number of Vendors</b> | <b>Total Invoices Collected</b> |
|-------------|--------------------------|---------------------------------|
| 1980        | *                        | *                               |
| 1981        | *                        | *                               |
| 1982        | *                        | *                               |
| 1983        | 3                        | 2,311                           |
| 1984        | 3                        | 2,587                           |
| 1985        | *                        | *                               |
| 1986        | *                        | *                               |
| 1987        | *                        | *                               |
| 1988        | *                        | *                               |
| 1989        | *                        | *                               |
| 1990        | 4                        | 2,803                           |
| 1991        | 3                        | 2,512                           |
| 1992        | 3                        | 2,737                           |
| 1993        | 3                        | 2,664                           |
| 1994        | *                        | *                               |
| 1995        | 3                        | 1,565                           |
| 1996        | 6                        | 1,965                           |
| 1997        | 7                        | 2,923                           |
| 1998        | 4                        | 3,591                           |
| 1999        | 5                        | 3,410                           |
| 2000        | 3                        | 3,868                           |
| 2001        | 3                        | 4,155                           |
| 2002        | 3                        | 3,494                           |
| 2003        | *                        | *                               |
| 2004        | 3                        | 3,104                           |
| 2005        | 3                        | 2,649                           |
| 2006        | 4                        | 2,589                           |
| 2007        | *                        | *                               |
| 2008        | *                        | *                               |
| 2009        | *                        | *                               |
| 2010        | *                        | *                               |
| 2011        | *                        | *                               |
| 2012        | *                        | *                               |
| 2013        | *                        | *                               |
| 2014        | 8                        | 1,353                           |
| 2015        | 9                        | 1,335                           |







|                  |          |              |
|------------------|----------|--------------|
| 2016             | 8        | 1,661        |
| 2017             | 11       | 1,969        |
| 2018             | 10       | 1,674        |
| <b>10 yr avg</b> | <b>5</b> | <b>1,586</b> |
| <b>10 yr SD</b>  | <b>4</b> | <b>265</b>   |
| <b>20 yr avg</b> | <b>4</b> | <b>2,293</b> |
| <b>20 yr SD</b>  | <b>3</b> | <b>886</b>   |

\*Less than three vendors

### 1.2.6 Fishery Summary Dashboard Statistics









The Fishery Summary Dashboard Statics section consolidates all fishery-dependent information comparing the most recent year with short-term (recent 10-year) and long-term (recent 20-year) average (shown bolded in [brackets]). Trend analysis of the past 10 years will dictate the trends (increasing, decreasing, or no trend). The right-most symbol indicates whether the mean of the short-term and long-term years were above, below, or within one standard deviation of the mean of the full time series.

Legend Key:

|   |   |
|---|---|
| <p> - increasing trend in the time series</p> <p> - decreasing trend in the time series</p> <p> - no trend in the time series</p> | <p> - above 1 standard deviation</p> <p> - below 1 standard deviation</p> <p> - within 1 standard deviation</p> |
|---|---|

10,000 [**1,000**] – point estimate of fishery statistic [*difference from short/long term average*]



**Table 27. 2018 annual indicators for the coral reef and bottomfish fishery describing fishery performance comparing current estimates with short-term (10 year) and long-term (20 year) average values**

| Fishery                                 | Fishery statistics   | Short-term (10 years)   | Long-term (20 years)   |
|---|--|---|--|
| <b>Bottomfish</b>                       | <b>Estimated catch (lbs.)</b>  |   |  |
| All species caught in the BF gear       | Boat and shore creel data estimated (expanded) total lbs. (all BF trips) | 32,751 [▲ 6%]    | 22,962 [▼ 10%]   |
|   | Estimated total lbs. (all species) commercial purchase data              | No trends available due to confidentiality  | No trends available due to confidentiality   |
| Bottomfish management unit species only | Total creel data Estimated (expanded) total lbs. (all BF trips)          | 32,750 [▲ 11%]   | 32,750 [▼ 8%]    |
|   | Estimated total lbs. (all  | No trends available due to  | No trends available due to   |

|   |   |  |  |
|---|---|--|--|
|   | species) commercial purchase data                   | confidentiality                            | confidentiality                            |
| <b>Catch-per-unit effort (lbs./gear-hours)</b>        |   |  |  |
|   | CPUE (creel data only)                              | 0.0076[▼58%]                               | 0.0076[▼52%]                               |
| <b>Fishing effort (only available for creel data)</b> |   |  |  |
|   | Estimated (expanded) total bottomfish # of trips    | 697[▼36%]                                  | 697[▼36%]                                  |
| <b>Fishing participants</b>                           |   |  |  |
|   | Estimated total # of fishers                        | 1,158[▲6%]                                 | 1,158[▲6%]                                 |
| <b>Bycatch</b>  |   |  |  |
|   | # bycatch caught                                    | 985[▼47%]                                  | 985[▼53%]                                  |
|   | # bycatch kept                                      | 1,492[▼19%]                                | 1,492[▼29%]                                |
|   | # bycatch released                                  | 57[▼49%]                                   | N/A  |
| <b>Coral Reef</b>                                     | <b>Estimated catch (lbs.)</b>                       |  |  |
|   | Boat-based creel data (expanded estimate all gears) | 136,385[▲12%]                              | 136,385[▼6%]                               |
|   | Shore-based creel (expanded estimate all gears)     | 106,707[▼16%]                              | 106,707[▼5%]                               |
|   | Commercial Purchase                                 | No trends available due to confidentiality | No trends available due to confidentiality |
| <b>Catch-per-unit-effort (lbs./gear hours)</b>        |   |  |  |
|   | BB spear  | 0.1607[▼24%]                               | 0.1607[▼5%]                                |
|   | BB SCUBA  | 0.4026[▼67%]                               | 0.4026[▼61%]                               |
|   | BB Gill Net   | 0.6667[▼57%]                               | 0.6667[▼45%]                               |
|   | BB Troll  | 0.0098[▼7%]                                | 0.0098[▼8%]                                |
|   | SB Hook and Line                                    | 0.0017[▼29%]                               | 0.0017[▲13%]                               |
|   | SB Throw/Cast Net                                   | 0.0935[▲129%]                              | 0.0935 [▲285%]                             |
|   | SB Gill Net   | 1.2284[▲350%]                              | 1.2284[▲673%]                              |
|   | SB Spear  | 0.9091[▲184%]                              | 0.9091[▲335%]                              |
|   | SB Hook and Gaff                                    | N/A  | N/A  |
| <b>Fishing effort (# of gear-hours by gear type)</b>  |   |  |  |



|   |                    |                 |                  |
|---|--------------------|-----------------|------------------|
|   | BB spear           | 3,960[▼81%]     | 3,960[▼29%]      |
|   | BB SCUBA           | 3,200[▲194%]    | 3,200[▲49%]      |
|   | BB Gill Net        | 15[▼94%]        | 15[▼98%]         |
|   | BB Troll           | 6,807,570[▲23%] | 6,807,570 [▲34%] |
|   | SB Hook and Line   | 144,364[▼5%]    | 144,364[▼52%]    |
|   | SB Throw/Cast Net  | 3,713[▼36%]     | 3,713[▼54%]      |
|   | SB Gill Net        | 162[▼87%]       | 162[▼98%]        |
|   | SB Spear           | 88[▼91%]        | 88[▼97%]         |
|   | SB Hook and Gaff   | N/A             | N/A              |
| <b>Fishing participants (# of gear)</b> |                    |                 |                  |
|   | BB spear           | 1,153[▲7%]      | 1,153[▲14%]      |
|   | BB SCUBA           | 1,460[▲12%]     | 1,460[▲24%]      |
|   | BB Gill Net        | 1,825[▲174%]    | 1,825[▲199%]     |
|   | BB Troll           | 1,178[▼1%]      | 1,178[▼7%]       |
|   | SB Hook and line   | 64,768[▼15%]    | 64,768[▼28%]     |
|   | SB Throw/cast net  | 8,014[▼37%]     | 8,014[▼46%]      |
|   | SB Gill net        | 2,796[▼55%]     | 2,796[▼63%]      |
|   | SB Spear           | 7,642[▼17%]     | 7,642[▼32%]      |
|   | SB Hook and Gaff   | 373[▼83%]       | 373[▼87%]        |
| <b>Boat- Based Bycatch</b>              |                    |                 |                  |
|   | # bycatch caught   | 9,580[▼15%]     | 9,580[▼10%]      |
|   | # bycatch kept     | 9,578[▼15%]     | 9,578[▼10%]      |
|   | # bycatch released | 2[▼71%]         | 2[▼88%]          |
| <b>Shore- Based Bycatch</b>             |                    |                 |                  |
|   | # bycatch caught   | 16,981[▲183%]   | 16,981[▲194%]    |
|   | # bycatch kept     | 16,965[▲188%]   | 16,965[▲198%]    |

|  |                    |   |  |
|--|--------------------|---|--|
|  | # bycatch released | 16[▼84%]   | 16[▼79%]   |
|--|--------------------|---|--|

### 1.2.7 Catch statistics

The following section summarizes the catch statistics for the bottomfish and coral reef fisheries in Guam. Estimates of catch are summarized from the creel survey and commercial receipt book data collection programs. Catch statistics provide estimates of annual harvest from the different fisheries. Estimates of fishery removals can provide proxies for the level of fishing mortality and a reference level relative to established quotas. This section also provides detailed levels of catch for fishing methods and the top species complexes harvested in the coral reef and bottomfish fisheries.

#### 1.2.7.1 Catch by Data Stream

This section describes the estimated total catch from the shore- and boat-based creel survey programs as well as the commercial landings from the commercial receipt book system. The difference between the creel total and the commercial landings is assumed to be the non-commercial component. However, there are cases where the commercial landing may be higher than the estimated creel total of the commercial receipt book program. In this case, the commercial receipt books are able to capture the fishery better than the creel surveys.

**Calculations:** Estimated landings are based on all bottomfish species harvested, regardless of the gear used, for all data collection programs (e.g. shore-based creel, boat-based creel and the commercial purchase reports).

**Table 28. Summary of time series of catch (lbs.) for all species caught using the bottomfishing gear in Guam from 1980-2018**

| Year | Creel Survey Estimates |             | Creel Total | Commercial landings |
|------|------------------------|-------------|-------------|---------------------|
|      | Boat-Based             | Shore-Based |             |                     |
| 1980 |                        |             |             | *                   |
| 1981 |                        |             |             | *                   |
| 1982 | 24,943                 | 0           | 24,943      | *                   |
| 1983 | 38,823                 | NULL        | 38,823      | 6,255               |
| 1984 | 39,146                 | NULL        | 39,146      | 5,329               |
| 1985 | 49,399                 | 333         | 49,732      | *                   |
| 1986 | 19,145                 | 451         | 19,596      | *                   |
| 1987 | 27,937                 | 12          | 27,949      | *                   |
| 1988 | 44,807                 | 3,100       | 47,907      | *                   |
| 1989 | 57,949                 | 76          | 58,025      | *                   |
| 1990 | 41,846                 | 3,872       | 45,718      | 5,664               |
| 1991 | 38,744                 | 6,957       | 45,701      | 3,061               |
| 1992 | 49,231                 | 4,233       | 53,464      | 2,994               |
| 1993 | 53,803                 | 1,348       | 55,151      | 4,621               |
| 1994 | 48,822                 | 545         | 49,367      | *                   |
| 1995 | 40,709                 | 2,108       | 42,817      | 7,695               |

|                  |               |            |               |              |
|------------------|---------------|------------|---------------|--------------|
| 1996             | 52,667        | 2,798      | 55,465        | 2,205        |
| 1997             | 30,232        | 1,946      | 32,178        | 2,687        |
| 1998             | 37,391        | 812        | 38,203        | 5,277        |
| 1999             | 52,795        | 1,066      | 53,861        | 22,025       |
| 2000             | 66,108        | 906        | 67,014        | 13,696       |
| 2001             | 50,864        | 178        | 51,042        | 11,900       |
| 2002             | 23,832        | 2,573      | 26,405        | 6,245        |
| 2003             | 41,677        | 439        | 42,116        | *            |
| 2004             | 37,266        | 1,040      | 38,306        | 10,453       |
| 2005             | 36,477        | 223        | 36,700        | 13,552       |
| 2006             | 37,713        | 1,769      | 39,482        | 9,436        |
| 2007             | 26,558        | 195        | 26,753        | *            |
| 2008             | 36,847        | 168        | 37,015        | *            |
| 2009             | 38,834        | 960        | 39,794        | *            |
| 2010             | 28,320        | 224        | 28,544        | *            |
| 2011             | 58,343        | 682        | 59,025        | *            |
| 2012             | 21,718        | 466        | 22,184        | *            |
| 2013             | 29,777        | 1,137      | 30,914        | *            |
| 2014             | 26,824        | 1,491      | 28,315        | 1,714        |
| 2015             | 15,142        | 499        | 15,641        | 923          |
| 2016             | 27,167        | 614        | 27,781        | 1,619        |
| 2017             | 22,267        | 695        | 22,962        | 5,153        |
| 2018             | 31,634        | 1,117      | 32,751        | 3,556        |
| <b>10 yr avg</b> | <b>30,003</b> | <b>789</b> | <b>30,791</b> | <b>4,863</b> |
| <b>10 yr SD</b>  | <b>11,199</b> | <b>362</b> | <b>11,257</b> | <b>3,248</b> |
| <b>20 yr avg</b> | <b>35,508</b> | <b>822</b> | <b>36,330</b> | <b>7,662</b> |
| <b>20 yr SD</b>  | <b>12,832</b> | <b>597</b> | <b>12,801</b> | <b>5,064</b> |

\*Less than three vendors

**Calculations:** Estimated landings are based on a pre-determined list of species (Appendix 3) identified as the BMUS Complex regardless of the gear used, for each data collection (shore-based creel, boat-based creel, and the commercial purchase reports).

**Table 29. Summary of time series of BMUS catch (lbs.) in Guam from 1980-2018**

| Year | Creel Survey Estimates |             | Creel Total | Commercial Landings |
|------|------------------------|-------------|-------------|---------------------|
|      | Boat-Based             | Shore-Based |             |                     |
| 1980 |                        |             |             | *                   |
| 1981 |                        |             |             | *                   |
| 1982 | 24,032                 |             | 24,032      | *                   |
| 1983 | 38,794                 |             | 38,794      | 6,255               |
| 1984 | 16,205                 |             | 16,205      | 5,329               |
| 1985 | 46,574                 | 4           | 46,578      | *                   |

|                  |               |            |               |              |
|------------------|---------------|------------|---------------|--------------|
| 1986             | 19,145        | 386        | 19,531        | *            |
| 1987             | 27,831        | 12         | 27,843        | *            |
| 1988             | 43,982        | 3,092      | 47,074        | *            |
| 1989             | 57,580        | 76         | 57,656        | *            |
| 1990             | 41,653        | 3,723      | 45,376        | 5,664        |
| 1991             | 38,253        | 6,849      | 45,102        | 3,061        |
| 1992             | 48,960        | 4,169      | 53,129        | 2,994        |
| 1993             | 53,457        | 1,184      | 54,641        | 4,621        |
| 1994             | 48,621        | 396        | 49,017        | *            |
| 1995             | 40,233        | 1,900      | 42,133        | 7,657        |
| 1996             | 52,484        | 2,718      | 55,202        | 2,205        |
| 1997             | 29,765        | 1,467      | 31,232        | 2,687        |
| 1998             | 36,966        | 409        | 37,375        | 5,267        |
| 1999             | 52,531        | 117        | 52,648        | 22,025       |
| 2000             | 65,682        | 768        | 66,450        | 13,534       |
| 2001             | 50,370        | 175        | 50,545        | 11,900       |
| 2002             | 23,803        | 2,572      | 26,375        | 6,245        |
| 2003             | 41,567        | 301        | 41,868        | *            |
| 2004             | 36,008        | 865        | 36,873        | 10,453       |
| 2005             | 36,431        | 129        | 36,560        | 13,552       |
| 2006             | 37,704        | 1,768      | 39,472        | 9,436        |
| 2007             | 26,558        | 194        | 26,752        | *            |
| 2008             | 36,847        | 168        | 37,015        | *            |
| 2009             | 38,342        | 905        | 39,247        | *            |
| 2010             | 26,821        | 223        | 27,044        | *            |
| 2011             | 58,343        | 680        | 59,023        | *            |
| 2012             | 21,718        | 464        | 22,182        | *            |
| 2013             | 29,741        | 1,128      | 30,869        | *            |
| 2014             | 23,466        | 1,399      | 24,865        | 1,651        |
| 2015             | 13,532        | 305        | 13,837        | 804          |
| 2016             | 26,380        | 512        | 26,892        | 1,619        |
| 2017             | 18,904        | 239        | 19,143        | 5,095        |
| 2018             | 31,634        | 1,116      | 32,750        | 3,556        |
| <b>10 yr avg</b> | <b>28,888</b> | <b>697</b> | <b>29,585</b> | <b>4,750</b> |
| <b>10 yr SD</b>  | <b>11,799</b> | <b>397</b> | <b>11,914</b> | <b>3,161</b> |
| <b>20 yr avg</b> | <b>34,819</b> | <b>701</b> | <b>35,521</b> | <b>7,586</b> |
| <b>20 yr SD</b>  | <b>13,235</b> | <b>628</b> | <b>13,170</b> | <b>5,060</b> |

\*Less than three vendors

**Calculations:** Estimated landings are based on a pre-determined list of species (Appendix 3) identified as the CREMUS Complex regardless of the gear used, for each data collection (shore-based creel, boat-based creel, and the commercial purchase reports).

**Table 30. Summary of time series of CREMUS catch (lbs.) in Guam from 1980-2018**

| Year | Creel Survey Estimates |             | Creel Total | Commercial Landings |
|------|------------------------|-------------|-------------|---------------------|
|      | Boat-Based             | Shore-Based |             |                     |
| 1980 |                        |             |             | *                   |
| 1981 |                        |             |             | *                   |
| 1982 | 29,248                 |             | 29,248      | *                   |
| 1983 | 53,077                 |             | 53,077      | 80,171              |
| 1984 | 95,924                 |             | 95,924      | 118,390             |
| 1985 | 131,353                | 401,187     | 532,540     | *                   |
| 1986 | 69,133                 | 236,498     | 305,631     | *                   |
| 1987 | 62,967                 | 229,383     | 292,350     | *                   |
| 1988 | 111,436                | 217,126     | 328,562     | *                   |
| 1989 | 156,378                | 153,837     | 310,215     | *                   |
| 1990 | 121,793                | 125,914     | 247,707     | 50,769              |
| 1991 | 171,220                | 261,531     | 432,751     | 38,322              |
| 1992 | 123,803                | 184,287     | 308,090     | 38,793              |
| 1993 | 174,809                | 100,143     | 274,952     | 33,320              |
| 1994 | 154,312                | 142,562     | 296,874     | *                   |
| 1995 | 267,515                | 189,515     | 457,030     | 26,304              |
| 1996 | 386,366                | 101,281     | 487,647     | 50,376              |
| 1997 | 219,166                | 191,563     | 410,729     | 72,762              |
| 1998 | 230,905                | 231,903     | 462,808     | 169,663             |
| 1999 | 374,272                | 277,098     | 651,370     | 258,789             |
| 2000 | 268,191                | 68,611      | 336,802     | 262,194             |
| 2001 | 256,389                | 84,594      | 340,983     | 267,622             |
| 2002 | 122,999                | 54,439      | 177,438     | 197,642             |
| 2003 | 152,096                | 117,200     | 269,296     | *                   |
| 2004 | 166,830                | 80,487      | 247,317     | 155,223             |
| 2005 | 88,942                 | 72,068      | 161,010     | 179,408             |
| 2006 | 86,051                 | 92,737      | 178,788     | 194,229             |
| 2007 | 72,870                 | 69,105      | 141,975     | *                   |
| 2008 | 103,971                | 67,362      | 171,333     | *                   |
| 2009 | 126,473                | 411,859     | 538,332     | *                   |
| 2010 | 76,133                 | 80,402      | 156,535     | *                   |
| 2011 | 260,962                | 77,422      | 338,384     | *                   |
| 2012 | 87,746                 | 149,342     | 237,088     | *                   |
| 2013 | 87,812                 | 181,043     | 268,855     | *                   |
| 2014 | 142,326                | 48,592      | 190,918     | 87,801              |
| 2015 | 122,065                | 81,157      | 203,222     | 58,762              |
| 2016 | 97,872                 | 56,971      | 154,843     | 73,250              |
| 2017 | 75,373                 | 72,055      | 147,428     | 273,375             |

|                  |                |                |                |                |
|------------------|----------------|----------------|----------------|----------------|
| 2018             | 136,385        | 106,707        | 243,092        | 133,941        |
| <b>10 yr avg</b> | <b>121,315</b> | <b>126,555</b> | <b>247,870</b> | <b>129,166</b> |
| <b>10 yr SD</b>  | <b>52,104</b>  | <b>102,847</b> | <b>112,010</b> | <b>61,775</b>  |
| <b>20 yr avg</b> | <b>145,288</b> | <b>112,463</b> | <b>257,750</b> | <b>164,488</b> |
| <b>20 yr SD</b>  | <b>79,819</b>  | <b>85,998</b>  | <b>129,944</b> | <b>65,388</b>  |

\*Less than three vendors

### 1.2.7.2 Expanded catch estimates by fishing methods

Catch information is provided for the top shore-based and boat-based fishing methods that contribute to a majority of the annual catch.

**Calculations:** The creel survey catch time series are the sum of the estimated weight for selected gear in all strata for all species (except for trolling, which exclude PMUS as well as any other pelagic species complex).

Table 31. Summary of expanded creel survey time series of catch (lbs.) by gear type in Guam for 1982-2018

| Year | Shore-Based Methods |        |         |        |        |        | Boat-Based Methods |        |         |        |
|------|---------------------|--------|---------|--------|--------|--------|--------------------|--------|---------|--------|
|      | Cast Net            | H&L    | Gillnet | Spear  | SCUBA  | H&G    | Bottom             | Spear  | SCUBA   | Troll* |
| 1982 |                     |        |         |        |        |        | 41,328             | 420    | 3,135   | 14,747 |
| 1983 |                     |        |         |        |        |        | 50,416             | 1,355  | 4,400   | 14,586 |
| 1984 |                     |        |         |        |        |        | 57,412             | 14,108 | 5,460   | 6,867  |
| 1985 | 83,628              | 41,488 | 59,241  | 83,182 | 3,136  | 6,900  | 88,045             | 18,737 | 12,761  | 18,692 |
| 1986 | 72,685              | 34,137 | 77,319  | 35,638 | 0      | 3,582  | 34,515             | 12,545 | 5,145   | 14,918 |
| 1987 | 75,312              | 31,262 | 78,088  | 31,650 | 0      | 2,076  | 44,459             | 12,448 | 7,474   | 12,440 |
| 1988 | 28,197              | 44,121 | 84,778  | 44,074 | 3,862  | 6,820  | 67,037             | 24,712 | 10,649  | 24,956 |
| 1989 | 38,948              | 40,012 | 40,550  | 13,435 | 1,282  | 8,267  | 79,972             | 30,930 | 20,839  | 15,349 |
| 1990 | 33,648              | 43,856 | 37,089  | 10,430 | 441    | 1,883  | 61,401             | 28,871 | 22,273  | 10,895 |
| 1991 | 105,524             | 52,137 | 51,556  | 18,085 | 70     | 3,748  | 60,753             | 27,898 | 37,027  | 19,522 |
| 1992 | 40,493              | 41,928 | 67,799  | 26,380 | 260    | 1,484  | 78,175             | 35,162 | 25,226  | 8,533  |
| 1993 | 20,711              | 14,840 | 21,458  | 30,996 | 497    | 4,053  | 107,130            | 39,434 | 22,848  | 5,611  |
| 1994 | 44,410              | 33,176 | 27,242  | 25,453 | 1,247  | 3,386  | 105,283            | 37,555 | 27,244  | 12,080 |
| 1995 | 81,934              | 22,492 | 25,148  | 38,939 | 14,452 | 2,207  | 101,073            | 40,554 | 74,734  | 17,045 |
| 1996 | 47,587              | 19,758 | 13,423  | 14,498 | 688    | 1,953  | 129,708            | 67,447 | 91,810  | 34,810 |
| 1997 | 61,155              | 34,158 | 16,456  | 20,248 | 237    | 2,159  | 109,346            | 37,363 | 41,920  | 16,396 |
| 1998 | 54,412              | 27,401 | 15,276  | 88,172 | 1,844  | 20,082 | 99,600             | 56,443 | 68,197  | 17,957 |
| 1999 | 100,194             | 26,485 | 33,541  | 75,345 | 320    | 15,294 | 122,930            | 45,200 | 82,024  | 30,561 |
| 2000 | 21,196              | 14,780 | 14,216  | 15,265 | 117    | 763    | 115,836            | 42,403 | 116,071 | 20,367 |
| 2001 | 22,304              | 7,362  | 8,934   | 21,083 | 106    | 5,670  | 123,975            | 74,369 | 65,103  | 17,581 |
| 2002 | 22,352              | 12,867 | 5,913   | 13,374 | 89     | 444    | 55,448             | 21,711 | 34,766  | 10,922 |
| 2003 | 40,729              | 16,174 | 10,975  | 50,456 | 157    | 177    | 82,223             | 22,649 | 42,685  | 30,524 |
| 2004 | 31,462              | 11,932 | 6,530   | 27,397 | 70     | 200    | 61,874             | 33,601 | 51,237  | 52,619 |
| 2005 | 23,509              | 8,286  | 22,033  | 8,073  | 394    | 7,944  | 62,651             | 15,037 | 32,375  | 13,387 |
| 2006 | 33,873              | 39,707 | 6,120   | 16,550 | 552    | 765    | 89,865             | 12,796 | 6,359   | 13,755 |
| 2007 | 28,815              | 6,066  | 15,867  | 12,053 | 137    | 5,131  | 57,750             | 24,704 | 29,989  | 10,567 |

|                  |               |               |               |               |            |              |               |               |               |               |
|------------------|---------------|---------------|---------------|---------------|------------|--------------|---------------|---------------|---------------|---------------|
| 2008             | 29,866        | 13,432        | 20,403        | 3,209         | 0          | 362          | 59,639        | 31,433        | 25,449        | 6,037         |
| 2009             | 44,133        | 342,402       | 6,569         | 2,329         | 0          | 13,746       | 89,997        | 22,669        | 37,424        | 12,185        |
| 2010             | 6,440         | 19,873        | 50,294        | 2,063         | 0          | 706          | 56,164        | 23,635        | 32,608        | 14,026        |
| 2011             | 38,331        | 33,663        | 2,607         | 1,619         | 211        | 378          | 88,694        | 26,483        | 67,431        | 6,637         |
| 2012             | 95,362        | 31,598        | 15,335        | 6,361         | 30         | 6,886        | 40,214        | 23,986        | 14,087        | 1,630         |
| 2013             | 44,113        | 98,377        | 26,579        | 6,675         | 148        | 4,090        | 42,601        | 20,816        | 5,390         | 26,073        |
| 2014             | 37,436        | 8,796         | 576           | 1,009         | 30         | 181          | 69,300        | 28,088        | 36,140        | 21,027        |
| 2015             | 49,829        | 10,332        | 8,140         | 45,819        | 0          | 1,755        | 29,395        | 22,371        | 34,607        | 19,750        |
| 2016             | 11,300        | 12,603        | 8,063         | 25,645        | 0          | 712          | 51,475        | 28,985        | 21,891        | 17,619        |
| 2017             | 29,163        | 33,063        | 2,873         | 4,000         | 307        | 762          | 46,715        | 17,792        | 11,201        | 12,095        |
| 2018             | 33,976        | 48,188        | 10,700        | 13,549        | 183        | 72           | 57,903        | 23,051        | 65,997        | 11,951        |
| <b>10 yr avg</b> | <b>39,008</b> | <b>63,890</b> | <b>13,174</b> | <b>10,907</b> | <b>152</b> | <b>2,929</b> | <b>57,246</b> | <b>23,788</b> | <b>32,678</b> | <b>14,299</b> |
| <b>10 yr SD</b>  | <b>23,019</b> | <b>96,116</b> | <b>14,283</b> | <b>13,658</b> | <b>99</b>  | <b>4,149</b> | <b>19,060</b> | <b>3,185</b>  | <b>20,060</b> | <b>6,773</b>  |
| <b>20 yr avg</b> | <b>37,219</b> | <b>39,799</b> | <b>13,813</b> | <b>17,594</b> | <b>190</b> | <b>3,302</b> | <b>70,232</b> | <b>28,089</b> | <b>40,642</b> | <b>17,466</b> |
| <b>20 yr SD</b>  | <b>22,766</b> | <b>72,432</b> | <b>11,737</b> | <b>18,958</b> | <b>141</b> | <b>4,455</b> | <b>26,653</b> | <b>13,141</b> | <b>26,831</b> | <b>10,947</b> |



### **1.2.7.3 Top Species in the Catch for the Boat- and Shore-Based Fisheries**

The time series for catch is an indicator of fishery performance. Fluctuations in the catch can be attributed to various factors and there is no single explanatory variable for the trends. The 10 species group in the boat and shore-based catch for the coral reef fishery make up 67% and 76%, respectively, of the total annual catches.

**Calculations:** Catch by species complex is tallied directly from the boat-based expanded species composition data combining all gear types and species, for all strata.

The averages for the table below were calculated from catch estimates for the entire time series across each of the CREMUS groupings. The average catch for each grouping is ranked from the highest to lowest. The dominant groups that make up more than half of the total annual catch are reported.

**Calculations:** Catch by species complex is tallied directly from the boat-based expanded species composition data combining all gear types and species, for all strata.

The averages for the table below were calculated from catch estimates from the entire time series for each of the CREMUS grouping. The average catch is ranked from the highest to lowest catch. The dominant groups that make up a majority of the catch are reported

**Table 32. Catch time series of 11 top CREMUS from Guam boat-based creel survey expansion data from 2000-2018**

| Year | Boat-Based Estimated Pounds |        |         |          |             |        |            |          |          |          |            |
|------|-----------------------------|--------|---------|----------|-------------|--------|------------|----------|----------|----------|------------|
|      | Bottomfish                  | BMUS   | Atulai  | Emperors | Surgeonfish | Jacks  | Parrotfish | Groupers | Snappers | Goatfish | Rabbitfish |
| 1982 | 24,944                      | 24,033 | 204     | 991      | 55          | 4,823  | 197        | 5,034    | 1,710    | 372      | 11         |
| 1983 | 38,824                      | 38,794 | 28,099  | 929      | 949         | 3,425  | 1,049      | 804      | 2,102    | 805      | 0          |
| 1984 | 39,144                      | 16,203 | 37,342  | 3,774    | 1,023       | 2,869  | 1,768      | 1,423    | 556      | 377      | 0          |
| 1985 | 49,401                      | 46,576 | 51,625  | 5,443    | 3,792       | 6,237  | 9,014      | 4,706    | 3,975    | 1,810    | 140        |
| 1986 | 19,147                      | 19,147 | 22,004  | 2,719    | 2,559       | 6,585  | 4,819      | 1,708    | 2,693    | 274      | 60         |
| 1987 | 27,938                      | 27,832 | 14,913  | 2,152    | 1,431       | 6,170  | 6,074      | 2,686    | 2,697    | 612      | 104        |
| 1988 | 44,808                      | 43,983 | 33,000  | 3,094    | 7,510       | 15,149 | 9,479      | 3,559    | 3,742    | 1,404    | 267        |
| 1989 | 57,946                      | 57,578 | 60,347  | 5,665    | 13,994      | 8,790  | 9,910      | 1,559    | 4,470    | 4,611    | 1,769      |
| 1990 | 41,846                      | 41,653 | 9,602   | 15,752   | 19,415      | 6,537  | 12,651     | 8,749    | 4,547    | 6,482    | 2,890      |
| 1991 | 38,744                      | 38,252 | 34,101  | 10,986   | 12,797      | 5,693  | 24,141     | 5,310    | 8,319    | 5,325    | 925        |
| 1992 | 49,231                      | 48,961 | 10,077  | 13,306   | 20,403      | 6,381  | 22,345     | 4,789    | 7,915    | 2,722    | 662        |
| 1993 | 53,805                      | 53,460 | 29,291  | 10,245   | 12,141      | 7,467  | 15,689     | 11,450   | 6,009    | 10,341   | 2,535      |
| 1994 | 48,822                      | 48,621 | 4,063   | 18,064   | 16,635      | 13,499 | 17,515     | 9,702    | 6,184    | 3,782    | 1,247      |
| 1995 | 40,706                      | 40,231 | 52,171  | 22,603   | 39,683      | 16,533 | 24,169     | 8,278    | 5,869    | 9,210    | 3,736      |
| 1996 | 52,669                      | 52,486 | 98,881  | 27,165   | 56,172      | 40,254 | 22,232     | 6,931    | 9,500    | 6,257    | 3,950      |
| 1997 | 30,233                      | 29,766 | 32,958  | 26,672   | 28,141      | 13,975 | 19,358     | 9,229    | 4,230    | 7,808    | 2,867      |
| 1998 | 37,390                      | 36,965 | 31,118  | 19,340   | 47,571      | 10,501 | 22,108     | 6,496    | 5,938    | 7,459    | 5,079      |
| 1999 | 52,795                      | 52,531 | 135,337 | 19,394   | 44,710      | 25,812 | 25,786     | 7,287    | 6,666    | 10,098   | 3,925      |
| 2000 | 66,109                      | 65,682 | 14,008  | 29,076   | 52,732      | 18,161 | 30,770     | 12,056   | 8,019    | 9,056    | 5,147      |
| 2001 | 50,866                      | 50,371 | 7,974   | 34,764   | 31,109      | 15,731 | 27,856     | 9,845    | 5,902    | 3,775    | 8,545      |
| 2002 | 23,835                      | 23,806 | 438     | 24,871   | 20,462      | 15,934 | 16,497     | 4,151    | 1,934    | 5,166    | 3,072      |
| 2003 | 41,677                      | 41,567 | 502     | 18,569   | 18,640      | 38,377 | 18,237     | 5,909    | 3,618    | 2,990    | 1,553      |
| 2004 | 37,266                      | 36,008 | 1,768   | 13,274   | 35,195      | 37,328 | 19,616     | 6,396    | 5,593    | 1,009    | 731        |
| 2005 | 36,479                      | 36,432 | 160     | 9,857    | 18,382      | 17,195 | 8,953      | 6,775    | 1,462    | 3,656    | 156        |
| 2006 | 37,713                      | 37,705 | 1,155   | 6,321    | 4,258       | 19,979 | 2,222      | 3,917    | 4,702    | 4,732    | 204        |
| 2007 | 26,558                      | 26,558 | 848     | 10,572   | 8,695       | 10,489 | 7,968      | 1,417    | 2,043    | 1,274    | 19         |

|                  |               |               |               |               |               |               |               |              |              |              |              |
|------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|--------------|--------------|--------------|
| 2008             | 36,844        | 36,844        | 10,335        | 7,560         | 24,395        | 8,460         | 7,524         | 7,205        | 5,538        | 6,599        | 1,486        |
| 2009             | 38,834        | 38,342        | 11,337        | 16,494        | 24,717        | 7,155         | 7,988         | 10,265       | 2,946        | 2,355        | 272          |
| 2010             | 28,320        | 26,821        | 5,887         | 11,940        | 11,518        | 7,706         | 6,788         | 3,884        | 3,623        | 1,460        | 485          |
| 2011             | 58,342        | 58,342        | 120,766       | 12,529        | 12,235        | 6,172         | 4,394         | 3,192        | 3,399        | 565          | 304          |
| 2012             | 21,718        | 21,718        | 24,936        | 7,210         | 3,313         | 3,083         | 5,206         | 1,950        | 2,857        | 2,470        | 1,349        |
| 2013             | 29,778        | 29,742        | 19,864        | 11,003        | 9,817         | 12,440        | 9,458         | 3,856        | 3,951        | 972          | 1,167        |
| 2014             | 26,823        | 23,465        | 4,077         | 22,347        | 10,376        | 13,027        | 8,856         | 5,136        | 3,741        | 8,399        | 3,808        |
| 2015             | 15,142        | 13,531        | 28,707        | 8,053         | 4,966         | 14,375        | 1,440         | 2,090        | 5,061        | 3,145        | 782          |
| 2016             | 27,165        | 26,379        | 2,523         | 9,419         | 7,672         | 18,661        | 10,493        | 3,352        | 2,561        | 1,615        | 784          |
| 2017             | 22,271        | 18,908        | 6,063         | 5,506         | 3,714         | 11,707        | 1,707         | 9,392        | 3,317        | 704          | 915          |
| 2018             | 31,636        | 31,636        | 8,503         | 5,417         | 5,408         | 7,560         | 4,117         | 4,321        | 2,986        | 1,070        | 311          |
| <b>10 yr avg</b> | <b>28,888</b> | <b>30,003</b> | <b>23,266</b> | <b>10,992</b> | <b>9,374</b>  | <b>10,189</b> | <b>6,045</b>  | <b>4,744</b> | <b>3,444</b> | <b>2,276</b> | <b>1,018</b> |
| <b>10 yr SD</b>  | <b>11,798</b> | <b>11,199</b> | <b>33,626</b> | <b>4,991</b>  | <b>5,968</b>  | <b>4,400</b>  | <b>3,014</b>  | <b>2,704</b> | <b>678</b>   | <b>2,190</b> | <b>994</b>   |
| <b>20 yr avg</b> | <b>34,819</b> | <b>35,509</b> | <b>20,259</b> | <b>14,209</b> | <b>17,616</b> | <b>15,468</b> | <b>11,294</b> | <b>5,620</b> | <b>3,996</b> | <b>3,556</b> | <b>1,751</b> |
| <b>20 yr SD</b>  | <b>13,235</b> | <b>12,832</b> | <b>36,878</b> | <b>8,047</b>  | <b>13,798</b> | <b>9,184</b>  | <b>8,624</b>  | <b>2,917</b> | <b>1,653</b> | <b>2,852</b> | <b>2,104</b> |

Table 33. Catch time series of 11 top CREMUS from Guam shore-based creel survey expansion data from 1980-2018

| Year | Shore-Based Estimated Pounds |            |          |        |          |        |        |          |            |            |
|------|------------------------------|------------|----------|--------|----------|--------|--------|----------|------------|------------|
|      | Surgeonfish                  | Rabbitfish | Mollusks | Atulai | Goatfish | Jacks  | Mullet | Emperors | Rudderfish | Parrotfish |
| 1985 | 71,019                       | 54,067     | 93,768   | 0      | 17,946   | 8,671  | 4,264  | 26,046   | 6,245      | 14,992     |
| 1986 | 60,497                       | 9,109      | 6,229    | 0      | 37,606   | 11,794 | 12,854 | 10,236   | 6,123      | 3,993      |
| 1987 | 74,665                       | 26,000     | 4,933    | 0      | 40,809   | 17,606 | 5,906  | 5,195    | 5,662      | 1,377      |
| 1988 | 65,455                       | 22,969     | 7,593    | 0      | 26,678   | 8,557  | 13,955 | 13,936   | 15,504     | 3,452      |
| 1989 | 33,304                       | 15,765     | 9,379    | 0      | 31,536   | 5,550  | 9,149  | 6,431    | 8,116      | 554        |
| 1990 | 25,985                       | 15,973     | 5,093    | 0      | 15,159   | 4,871  | 9,641  | 6,183    | 9,561      | 1,527      |
| 1991 | 27,229                       | 38,600     | 40,594   | 0      | 15,520   | 44,313 | 10,591 | 6,197    | 21,608     | 11,970     |
| 1992 | 48,450                       | 30,397     | 11,114   | 0      | 14,553   | 2,815  | 15,182 | 10,983   | 9,104      | 5,547      |
| 1993 | 15,598                       | 9,106      | 6,724    | 0      | 10,199   | 3,270  | 4,472  | 5,537    | 2,188      | 3,339      |
| 1994 | 28,537                       | 29,842     | 12,677   | 0      | 8,003    | 5,715  | 9,781  | 8,194    | 320        | 5,189      |

|                  |               |               |              |          |              |               |               |              |              |              |
|------------------|---------------|---------------|--------------|----------|--------------|---------------|---------------|--------------|--------------|--------------|
| 1995             | 68,150        | 15,855        | 7,966        | 0        | 12,856       | 6,083         | 5,413         | 11,736       | 2,475        | 8,387        |
| 1996             | 30,569        | 16,365        | 4,031        | 0        | 6,133        | 4,931         | 3,024         | 7,205        | 2,997        | 3,900        |
| 1997             | 35,075        | 20,921        | 7,425        | 0        | 5,638        | 11,571        | 3,678         | 8,295        | 3,690        | 5,489        |
| 1998             | 47,832        | 12,584        | 29,751       | 0        | 7,485        | 13,658        | 3,439         | 11,414       | 20,250       | 11,137       |
| 1999             | 67,325        | 24,757        | 22,422       | 0        | 8,295        | 34,539        | 4,999         | 10,080       | 6,022        | 20,155       |
| 2000             | 12,294        | 11,376        | 3,940        | 0        | 4,281        | 3,372         | 1,409         | 4,632        | 5,671        | 2,046        |
| 2001             | 15,010        | 5,845         | 10,165       | 0        | 4,887        | 8,593         | 1,011         | 5,185        | 754          | 1,344        |
| 2002             | 8,036         | 6,622         | 1,934        | 0        | 2,582        | 4,954         | 1,492         | 3,328        | 10,706       | 1,458        |
| 2003             | 27,732        | 12,602        | 16,589       | 0        | 7,615        | 7,799         | 1,495         | 5,135        | 2,111        | 8,533        |
| 2004             | 20,297        | 7,566         | 5,660        | 0        | 2,820        | 12,765        | 677           | 2,353        | 2,317        | 2,613        |
| 2005             | 11,572        | 3,578         | 11,459       | 0        | 7,079        | 11,155        | 2,907         | 3,547        | 734          | 1,931        |
| 2006             | 28,996        | 11,572        | 2,689        | 0        | 9,008        | 8,594         | 531           | 1,367        | 11,950       | 3,083        |
| 2007             | 18,694        | 11,056        | 6,875        | 0        | 1,381        | 12,301        | 1,096         | 5,483        | 944          | 926          |
| 2008             | 19,072        | 12,988        | 1,390        | 0        | 698          | 10,330        | 4,720         | 1,471        | 1,973        | 2,049        |
| 2009             | 201,749       | 86,424        | 14,721       | 0        | 2,211        | 27,106        | 45,336        | 5,705        | 911          | 979          |
| 2010             | 38,327        | 8,257         | 1,489        | 0        | 2,091        | 4,181         | 2,568         | 2,783        | 8,283        | 1,000        |
| 2011             | 12,343        | 9,443         | 1,099        | 0        | 8,193        | 29,650        | 2,195         | 5,724        | 164          | 3,234        |
| 2012             | 7,820         | 2,630         | 9,848        | 0        | 13,338       | 80,189        | 14,455        | 5,974        | 1,542        | 951          |
| 2013             | 43,947        | 26,221        | 5,338        | 0        | 34,187       | 36,409        | 4,750         | 3,422        | 139          | 619          |
| 2014             | 4,194         | 4,349         | 666          | 0        | 3,715        | 19,022        | 70            | 8,742        | 387          | 486          |
| 2015             | 15,609        | 5,654         | 22,082       | 0        | 9,515        | 9,023         | 603           | 3,962        | 863          | 2,527        |
| 2016             | 17,916        | 8,442         | 3,795        | 0        | 4,392        | 3,750         | 719           | 2,032        | 736          | 1,394        |
| 2017             | 28,873        | 6,070         | 2,636        | 0        | 2,761        | 14,558        | 2,584         | 700          | 358          | 410          |
| 2018             | 16,927        | 38,164        | 1,254        | 0        | 7,560        | 20,941        | 59            | 9,653        | 37           | 1,422        |
| <b>10 yr avg</b> | <b>19,565</b> | <b>38,771</b> | <b>6,293</b> | <b>0</b> | <b>8,796</b> | <b>24,483</b> | <b>7,334</b>  | <b>4,870</b> | <b>1,342</b> | <b>1,302</b> |
| <b>10 yr SD</b>  | <b>24,715</b> | <b>55,666</b> | <b>6,787</b> | <b>0</b> | <b>9,153</b> | <b>21,215</b> | <b>13,297</b> | <b>2,706</b> | <b>2,354</b> | <b>866</b>   |
| <b>20 yr avg</b> | <b>15,181</b> | <b>30,837</b> | <b>7,303</b> | <b>0</b> | <b>6,830</b> | <b>17,962</b> | <b>4,684</b>  | <b>4,564</b> | <b>2,830</b> | <b>2,858</b> |
| <b>20 yr SD</b>  | <b>18,441</b> | <b>41,739</b> | <b>6,747</b> | <b>0</b> | <b>7,056</b> | <b>17,357</b> | <b>9,829</b>  | <b>2,583</b> | <b>3,569</b> | <b>4,321</b> |

### 1.2.8 Catch-per-Unit-Effort (CPUE) Statistics

This section summarizes the estimates for catch-per-unit effort in the boat and shore-based fisheries. The boat-based fisheries include the bottomfishing (handline gear), spearfishing (SCUBA and snorkel), gillnets, and troll that comprise 83% of the total catch. Trolling methods are primarily a pelagic fishing method but also catches coral reef fishes like jacks and gray jobfish. The shore-based fisheries include the hook-and-line, throw or cast nets, gillnets, spear, and hook-and-gaff that comprise 88% of the total coral reef fish catch. CPUE is reported as pounds per gear-hours for the shore-based methods whereas in the boat-based methods it's pounds per trip.

**Calculations:** CPUE is calculated from interview data by gear type using  $\sum \text{catch} / \sum (\text{hours fished} * \text{number of fishers})$  for boat based and  $\sum \text{catch} / \sum (\text{hours fished} * \text{number of gears used})$  for shore based. If the value is blank (empty), then there was no interview collected for that method. Landings from interviews without fishing hours are excluded from the calculations.

**Table 34. CPUE time series for dominant fishing methods in Guam shore-based fisheries  
CPUE from 1984-2018**

| Year | Shore-Based Gear CPUE (lbs./gear hour) |          |          |        |        |
|------|--|----------|----------|--------|--------|
|      | H&L                                    | Cast Net | Gill Net | Spear  | H&G    |
| 1984 | 0.0106                                 | 0.1339   | 0.3507   | 0.7500 | 1.1250 |
| 1985 | 0.0029                                 | 0.0224   | 0.0509   | 0.0773 | 0.0975 |
| 1986 | 0.0040                                 | 0.0224   | 0.0441   | 0.0962 | 0.2393 |
| 1987 | 0.0074                                 | 0.0208   | 0.0515   | 0.0747 | 0.0354 |
| 1988 | 0.0027                                 | 0.0213   | 0.0764   | 0.0805 | 0.2444 |
| 1989 | 0.0022                                 | 0.0136   | 0.0548   | 0.0627 | 0.2545 |
| 1990 | 0.0011                                 | 0.0171   | 0.0309   | 0.0590 | 0.0551 |
| 1991 | 0.0017                                 | 0.0128   | 0.0305   | 0.0918 | 0.0690 |
| 1992 | 0.0005                                 | 0.0122   | 0.0255   | 0.0986 | 0.0327 |
| 1993 | 0.0003                                 | 0.0060   | 0.0181   | 0.1621 | 0.0347 |
| 1994 | 0.0004                                 | 0.0160   | 0.0208   | 0.0370 | 0.0734 |
| 1995 | 0.0005                                 | 0.0064   | 0.0117   | 0.0734 | 0.0313 |
| 1996 | 0.0003                                 | 0.0158   | 0.0220   | 0.0659 | 0.0938 |
| 1997 | 0.0004                                 | 0.0060   | 0.0134   | 0.0415 | 0.0544 |
| 1998 | 0.0005                                 | 0.0082   | 0.0067   | 0.0544 | 0.1094 |
| 1999 | 0.0005                                 | 0.0076   | 0.0124   | 0.0316 | 0.1925 |
| 2000 | 0.0004                                 | 0.0083   | 0.0189   | 0.0476 | 0.0381 |
| 2001 | 0.0004                                 | 0.0045   | 0.0204   | 0.0575 | 0.2946 |
| 2002 | 0.0007                                 | 0.0152   | 0.0184   | 0.0906 | 0.4500 |
| 2003 | 0.0007                                 | 0.0034   | 0.0359   | 0.1844 | 0.0256 |
| 2004 | 0.0010                                 | 0.0051   | 0.0290   | 0.1257 | 0.2222 |
| 2005 | 0.0005                                 | 0.0019   | 0.0781   | 0.1333 | 0.2593 |
| 2006 | 0.0015                                 | 0.0169   | 0.0373   | 0.1035 | 0.2889 |
| 2007 | 0.0007                                 | 0.0071   | 0.1264   | 0.1555 | 0.4286 |

|                  |               |               |               |               |               |
|------------------|---------------|---------------|---------------|---------------|---------------|
| 2008             | 0.0009        | 0.0064        | 0.0738        | 0.0489        | 0.1333        |
| 2009             | 0.0010        | 0.1468        | 0.1294        | 0.1222        | 0.3524        |
| 2010             | 0.0003        | 0.0138        | 0.2598        | 0.2708        | 0.2115        |
| 2011             | 0.0018        | 0.0203        | 0.1245        | 0.7429        | 0.5200        |
| 2012             | 0.0020        | 0.0188        | 0.1356        | 0.1527        | 0.2143        |
| 2013             | 0.0017        | 0.0438        | 0.1176        | 0.0988        | 0.2639        |
| 2014             | 0.0030        | 0.0141        | 0.4388        | 0.4688        | 0.2857        |
| 2015             | 0.0102        | 0.0147        | 0.0673        | 0.3298        | 0.4231        |
| 2016             | 0.0006        | 0.0051        | 0.0269        | 0.0290        | 0.4000        |
| 2017             | 0.0013        | 0.0377        | 0.2016        | 0.0806        | 0.0577        |
| 2018             | 0.0017        | 0.0935        | 1.2284        | 0.9091        | 0.0000        |
| <b>10 yr avg</b> | <b>0.0024</b> | <b>0.0409</b> | <b>0.2730</b> | <b>0.3205</b> | <b>0.3032</b> |
| <b>10 yr SD</b>  | <b>0.0027</b> | <b>0.0428</b> | <b>0.3368</b> | <b>0.2844</b> | <b>0.1298</b> |
| <b>20 yr avg</b> | <b>0.0015</b> | <b>0.0243</b> | <b>0.1590</b> | <b>0.2092</b> | <b>0.2664</b> |
| <b>20 yr SD</b>  | <b>0.0021</b> | <b>0.0347</b> | <b>0.2651</b> | <b>0.2324</b> | <b>0.1377</b> |

**Table 35. CPUE time series for dominant fishing methods in Guam boat-based fisheries  
CPUE from 1982-2018**

| Year | Boat-Based Gear CPUE (lbs./gear hours) |        |        |          |        |
|------|--|--------|--------|----------|--------|
|      | Bottomfishing                          | Spear  | SCUBA  | Gill Net | Troll  |
| 1982 | 0.0293                                 | 0.4800 | 0.0000 | 0.0000   | 0.0162 |
| 1983 | 0.0293                                 | 0.2198 | 0.3956 | 0.0000   | 0.0154 |
| 1984 | 0.0230                                 | 0.1159 | 0.3553 | 3.0000   | 0.0135 |
| 1985 | 0.0099                                 | 0.2025 | 0.1598 | 0.5357   | 0.0098 |
| 1986 | 0.0210                                 | 0.2915 | 0.4402 | 0.5000   | 0.0092 |
| 1987 | 0.0223                                 | 0.2312 | 0.5550 | 0.3195   | 0.0086 |
| 1988 | 0.0114                                 | 0.1518 | 0.2097 | 0.6465   | 0.0057 |
| 1989 | 0.0106                                 | 0.1194 | 0.2343 | 0.4050   | 0.0048 |
| 1990 | 0.0116                                 | 0.1515 | 0.6306 | 0.3795   | 0.0037 |
| 1991 | 0.0116                                 | 0.1691 | 0.4482 | 0.3110   | 0.0051 |
| 1992 | 0.0106                                 | 0.0794 | 0.1164 | 0.2381   | 0.0034 |
| 1993 | 0.0102                                 | 0.0637 | 0.4413 | 0.6389   | 0.0041 |
| 1994 | 0.0109                                 | 0.0766 | 0.3632 | 0.3262   | 0.0039 |
| 1995 | 0.0029                                 | 0.0568 | 0.2424 | 0.1213   | 0.0032 |
| 1996 | 0.0035                                 | 0.0586 | 0.2149 | 0.4762   | 0.0034 |
| 1997 | 0.0029                                 | 0.0706 | 0.4460 | 0.2965   | 0.0040 |
| 1998 | 0.0027                                 | 0.0252 | 0.3077 | 0.1199   | 0.0035 |
| 1999 | 0.0035                                 | 0.0334 | 0.2841 | 0.6192   | 0.0031 |
| 2000 | 0.0052                                 | 0.0532 | 0.2758 | 0.0661   | 0.0042 |
| 2001 | 0.0071                                 | 0.1912 | 0.3202 | 0.3005   | 0.0069 |
| 2002 | 0.0069                                 | 0.0857 | 0.5128 | 0.4275   | 0.0117 |
| 2003 | 0.0172                                 | 0.1880 | 0.7129 | 1.8968   | 0.0176 |

|                  |               |               |               |               |               |
|------------------|---------------|---------------|---------------|---------------|---------------|
| 2004             | 0.0143        | 0.2008        | 0.7860        | 1.0195        | 0.0174        |
| 2005             | 0.0171        | 0.0848        | 0.7361        | 0.4407        | 0.0104        |
| 2006             | 0.0230        | 0.1134        | 0.3905        | 1.7500        | 0.0114        |
| 2007             | 0.0226        | 0.2217        | 4.0816        | 0.5214        | 0.0136        |
| 2008             | 0.0162        | 0.1087        | 0.6206        | 1.5606        | 0.0100        |
| 2009             | 0.0164        | 0.0795        | 1.7182        | 0.2311        | 0.0083        |
| 2010             | 0.0081        | 0.0828        | 0.3333        | 0.3787        | 0.0067        |
| 2011             | 0.0270        | 0.2714        | 2.6571        | 0.5000        | 0.0095        |
| 2012             | 0.0341        | 0.8788        | 3.0000        | 10.3504       | 0.0185        |
| 2013             | 0.0254        | 0.1598        | 0.9375        | 0.4643        | 0.0147        |
| 2014             | 0.0172        | 0.1629        | 1.5469        | 1.3313        | 0.0109        |
| 2015             | 0.0163        | 0.1729        | 0.5435        | 0.9467        | 0.0125        |
| 2016             | 0.0137        | 0.0961        | 0.2078        | 0.1993        | 0.0074        |
| 2017             | 0.0151        | 0.0501        | 0.8095        | 0.3646        | 0.0065        |
| 2018             | 0.0076        | 0.1607        | 0.4026        | 0.6667        | 0.0098        |
| <b>10 yr avg</b> | <b>0.0181</b> | <b>0.2115</b> | <b>1.2156</b> | <b>1.5433</b> | <b>0.0105</b> |
| <b>10 yr SD</b>  | <b>0.0080</b> | <b>0.2305</b> | <b>0.9367</b> | <b>2.9540</b> | <b>0.0036</b> |
| <b>20 yr avg</b> | <b>0.0157</b> | <b>0.1698</b> | <b>1.0439</b> | <b>1.2018</b> | <b>0.0106</b> |
| <b>20 yr SD</b>  | <b>0.0078</b> | <b>0.1742</b> | <b>1.0283</b> | <b>2.1617</b> | <b>0.0042</b> |

### 1.2.9 Effort Statistics

This section summarizes the effort trends in the coral reef and bottomfish fishery. Fishing effort trends provide insights on the level of fishing pressure through time. Effort information is provided for the top shore-based and boat-based fishing methods that contribute to a majority of the annual catch. Trolling method is included in this report because coral reef MUS is also caught using trolling method. Pelagic MUS caught using trolling method is reported in the Pelagic Annual/SAFE report module.

**Calculations:** Effort estimates (hours) are generated by summing the effort data collected from interviews by gear type. For shore-based estimates, data collection started in 1985.

**Table 36. Time series of effort estimates (gear hours or no. trips for bottomfish) from Guam coral reef and bottomfish fisheries for 1982-2018**

| Year | Shore-Based Gear Hours |           |          |        |       | Boat-Based Gear Hours |        |        |          |            |
|------|------------------------|-----------|----------|--------|-------|-----------------------|--------|--------|----------|------------|
|      | Cast Net               | H&L       | Gill Net | Spear  | H&G   | Bottom                | Spear  | Scuba  | Gill Net | Troll      |
| 1982 | 15                     | 400       | 0        | 208    | 0     | 81,620                | 65     | 1      | 0        | 3,046,932  |
| 1983 | 0                      | 0         | 0        | 0      | 0     | 59,512                | 143    | 527    | 0        | 2,615,565  |
| 1984 | 224                    | 2,914     | 345      | 24     | 8     | 131,159               | 6,156  | 630    | 15       | 2,548,752  |
| 1985 | 5,673                  | 82,992    | 10,658   | 15,096 | 400   | 532,350               | 4,092  | 5,304  | 21       | 4,709,880  |
| 1986 | 3,430                  | 52,899    | 14,378   | 3,410  | 117   | 98,112                | 1,888  | 304    | 2        | 3,019,692  |
| 1987 | 4,902                  | 18,204    | 8,550    | 9,964  | 4,779 | 113,442               | 2,257  | 624    | 493      | 3,946,710  |
| 1988 | 8,487                  | 34,662    | 9,735    | 6,264  | 225   | 295,911               | 6,375  | 1,920  | 44       | 9,291,900  |
| 1989 | 15,810                 | 42,120    | 6,336    | 2,184  | 224   | 331,525               | 4,416  | 2,655  | 100      | 7,495,286  |
| 1990 | 13,534                 | 253,492   | 20,240   | 2,679  | 272   | 249,280               | 1,794  | 1,200  | 640      | 11,182,260 |
| 1991 | 13,932                 | 368,466   | 17,835   | 1,862  | 1,638 | 197,964               | 2,016  | 2,142  | 918      | 9,667,476  |
| 1992 | 13,900                 | 739,440   | 30,000   | 1,440  | 490   | 202,400               | 5,893  | 6,820  | 414      | 11,705,316 |
| 1993 | 12,604                 | 796,708   | 18,040   | 1,666  | 1,701 | 270,758               | 8,961  | 2,520  | 324      | 11,355,743 |
| 1994 | 6,048                  | 978,945   | 21,070   | 7,520  | 722   | 383,520               | 8,827  | 3,569  | 1,300    | 11,652,024 |
| 1995 | 19,840                 | 673,200   | 40,608   | 7,221  | 384   | 1,258,615             | 24,497 | 16,268 | 5,520    | 17,307,210 |
| 1996 | 4,875                  | 939,333   | 8,601    | 2,684  | 96    | 1,351,026             | 28,310 | 13,959 | 5,244    | 20,231,220 |
| 1997 | 19,760                 | 1,120,575 | 31,692   | 5,328  | 294   | 1,017,597             | 13,144 | 3,713  | 3,080    | 13,812,489 |
| 1998 | 21,976                 | 795,960   | 73,066   | 15,006 | 448   | 1,526,630             | 62,160 | 10,126 | 3,348    | 16,974,006 |
| 1999 | 14,351                 | 1,234,925 | 52,116   | 26,010 | 504   | 1,230,288             | 20,574 | 12,060 | 1,122    | 12,031,104 |
| 2000 | 14,157                 | 838,240   | 27,930   | 9,416  | 315   | 622,364               | 15,930 | 10,856 | 8,064    | 11,211,280 |
| 2001 | 15,125                 | 827,519   | 16,464   | 3,968  | 224   | 483,060               | 5,940  | 4,860  | 1,008    | 6,544,218  |
| 2002 | 7,614                  | 227,813   | 14,691   | 2,352  | 20    | 278,604               | 5,544  | 960    | 384      | 2,681,143  |
| 2003 | 18,900                 | 345,598   | 2,950    | 1,394  | 195   | 148,160               | 3,596  | 1,369  | 147      | 1,405,206  |
| 2004 | 7,885                  | 195,202   | 4,662    | 1,050  | 36    | 168,413               | 2,295  | 1,044  | 66       | 2,336,400  |
| 2005 | 9,400                  | 167,334   | 1,242    | 360    | 54    | 190,400               | 4,368  | 480    | 253      | 2,290,578  |
| 2006 | 6,336                  | 96,074    | 2,091    | 425    | 45    | 147,125               | 3,618  | 117    | 2        | 2,796,184  |
| 2007 | 2,948                  | 343,952   | 546      | 418    | 70    | 92,820                | 1,550  | 49     | 154      | 2,443,480  |



|                  |              |                |               |              |            |                |              |              |              |                  |
|------------------|--------------|----------------|---------------|--------------|------------|----------------|--------------|--------------|--------------|------------------|
| 2008             | 5,976        | 164,300        | 1,720         | 266          | 15         | 127,710        | 8,393        | 289          | 264          | 2,771,390        |
| 2009             | 4,026        | 185,298        | 255           | 180          | 210        | 285,891        | 6,072        | 100          | 532          | 6,262,704        |
| 2010             | 7,313        | 141,860        | 408           | 144          | 156        | 370,360        | 5,250        | 6            | 168          | 7,455,312        |
| 2011             | 5,184        | 103,653        | 988           | 70           | 25         | 136,284        | 1,800        | 196          | 3            | 3,945,474        |
| 2012             | 6,006        | 122,850        | 1,128         | 550          | 70         | 30,084         | 504          | 65           | 45           | 1,194,173        |
| 2013             | 4,221        | 81,774         | 672           | 729          | 72         | 47,061         | 1,710        | 24           | 1,120        | 3,601,465        |
| 2014             | 4,544        | 130,062        | 196           | 224          | 28         | 144,690        | 3,528        | 40           | 210          | 4,490,376        |
| 2015             | 5,858        | 227,766        | 3,358         | 1,980        | 156        | 65,262         | 2,842        | 391          | 65           | 5,278,731        |
| 2016             | 14,040       | 183,219        | 4,717         | 5,520        | 20         | 170,159        | 6,210        | 6,732        | 189          | 9,152,541        |
| 2017             | 3,320        | 191,836        | 506           | 484          | 104        | 176,253        | 8,051        | 140          | 64           | 7,157,862        |
| 2018             | 3,713        | 144,364        | 162           | 88           | 0          | 349,272        | 3,960        | 3,200        | 15           | 6,807,570        |
| <b>10 yr avg</b> | <b>5,823</b> | <b>151,268</b> | <b>1,239</b>  | <b>997</b>   | <b>84</b>  | <b>177,532</b> | <b>3,993</b> | <b>1,089</b> | <b>241</b>   | <b>5,534,621</b> |
| <b>10 yr SD</b>  | <b>2,972</b> | <b>42,622</b>  | <b>1,464</b>  | <b>1,601</b> | <b>67</b>  | <b>115,162</b> | <b>2,259</b> | <b>2,096</b> | <b>328</b>   | <b>2,186,818</b> |
| <b>20 yr avg</b> | <b>8,046</b> | <b>297,682</b> | <b>6,840</b>  | <b>2,781</b> | <b>116</b> | <b>263,213</b> | <b>5,587</b> | <b>2,149</b> | <b>694</b>   | <b>5,092,860</b> |
| <b>20 yr SD</b>  | <b>4,572</b> | <b>298,257</b> | <b>12,497</b> | <b>5,792</b> | <b>122</b> | <b>265,257</b> | <b>4,757</b> | <b>3,568</b> | <b>1,727</b> | <b>3,082,233</b> |

### **1.2.10 Participants**

This section summarizes the estimated number of participants in each fishery. The information presented here can be used in the impact analysis of potential amendments in the FEPs associated with the bottomfish and coral reef fisheries. The trend in the number of participants over time can also be used as an indicator for fishing pressure.

**Calculations:** For boat-based data, the estimated number of participants is calculated by multiplying the average number of fishers per trip by the number of trips per day, and then by the number of dates in the calendar year by gear type. The total is a combination of weekend and weekday stratum estimates.

For shore-based data, the estimated number of participants is calculated by using an average number of fishers per day multiplied by the numbers of dates in the calendar year across gear types. The total is a combination of weekend, weekday, day, and night stratum estimates.

**Table 37. Number of boats participating in the Guam bottomfish fishery and number of gears in the Guam boat- and shore-based coral reef fisheries from 1982-2018**

| Year | Bottomfish |         | Coral Reef Boat-Based |       |          |       | Coral Reef Shore-Based |        |          |        |       |
|------|------------|---------|-----------------------|-------|----------|-------|------------------------|--------|----------|--------|-------|
|      | # Fishers  | # Gears | Spear                 | SCUBA | Gill Net | Troll | H&L                    | Throw  | Gill Net | Spear  | H&G   |
| 1982 | 865        | 798     | 1,095                 | 365   | 0        | 920   |                        |        |          |        |       |
| 1983 | 820        | 709     | 852                   | 533   | 0        | 955   |                        |        |          |        |       |
| 1984 | 977        | 847     | 1,519                 | 701   | 732      | 1,022 | 101,016                | 18,141 | 18,523   | 7,065  | 2,101 |
| 1985 | 971        | 883     | 1,326                 | 852   | 1,460    | 952   | 120,562                | 32,345 | 37,904   | 21,282 | 3,931 |
| 1986 | 918        | 794     | 913                   | 1,049 | 1,095    | 975   | 90,441                 | 21,308 | 46,996   | 19,236 | 2,072 |
| 1987 | 874        | 829     | 712                   | 830   | 1,095    | 964   | 108,511                | 25,715 | 49,381   | 18,297 | 1,978 |
| 1988 | 975        | 903     | 987                   | 864   | 824      | 1,151 | 98,891                 | 23,518 | 42,645   | 25,360 | 5,242 |
| 1989 | 931        | 869     | 1,156                 | 1,065 | 730      | 1,122 | 125,421                | 26,558 | 28,505   | 10,985 | 4,310 |
| 1990 | 1,002      | 883     | 1,338                 | 1,116 | 1,004    | 1,247 | 101,800                | 23,666 | 32,991   | 11,233 | 2,896 |
| 1991 | 1,049      | 843     | 1,241                 | 1,136 | 962      | 1,287 | 215,674                | 39,177 | 64,483   | 15,087 | 6,002 |
| 1992 | 1,067      | 886     | 1,330                 | 1,243 | 1,098    | 1,335 | 186,939                | 38,170 | 76,740   | 18,606 | 3,673 |
| 1993 | 1,028      | 910     | 1,191                 | 1,359 | 776      | 1,236 | 189,891                | 41,884 | 46,720   | 19,527 | 6,296 |
| 1994 | 1,103      | 947     | 1,204                 | 1,278 | 791      | 1,217 | 217,996                | 33,762 | 43,891   | 18,615 | 4,015 |
| 1995 | 1,327      | 1,275   | 1,062                 | 1,362 | 1,137    | 1,239 | 246,531                | 37,900 | 48,269   | 21,453 | 7,956 |
| 1996 | 1,609      | 1,562   | 1,074                 | 1,311 | 864      | 1,253 | 252,664                | 24,115 | 32,650   | 16,408 | 7,127 |
| 1997 | 1,816      | 1,581   | 1,033                 | 1,406 | 1,000    | 1,215 | 210,044                | 27,784 | 29,222   | 12,944 | 2,550 |
| 1998 | 1,393      | 1,305   | 1,046                 | 1,396 | 960      | 1,164 | 158,460                | 37,500 | 54,300   | 22,920 | 6,780 |
| 1999 | 1,441      | 1,387   | 1,181                 | 1,426 | 1,121    | 1,121 | 217,454                | 24,670 | 46,892   | 37,939 | 8,116 |
| 2000 | 1,391      | 1,321   | 1,075                 | 1,303 | 1,236    | 1,103 | 129,407                | 18,666 | 23,163   | 17,202 | 3,712 |
| 2001 | 1,043      | 1,078   | 1,178                 | 1,309 | 1,235    | 1,090 | 120,039                | 18,980 | 17,839   | 12,957 | 3,513 |
| 2002 | 1,197      | 1,037   | 1,019                 | 1,294 | 986      | 1,030 | 90,023                 | 17,893 | 12,301   | 7,688  | 1,258 |
| 2003 | 924        | 1,092   | 1,344                 | 1,488 | 1,095    | 1,127 | 89,197                 | 21,763 | 15,239   | 11,908 | 958   |
| 2004 | 1,229      | 1,121   | 990                   | 1,298 | 854      | 1,011 | 80,756                 | 13,365 | 17,001   | 10,720 | 708   |
| 2005 | 974        | 965     | 1,019                 | 1,251 | 803      | 1,114 | 75,783                 | 17,109 | 11,452   | 7,574  | 3,422 |
| 2006 | 918        | 956     | 1,153                 | 949   | 730      | 1,068 | 71,494                 | 21,033 | 14,691   | 12,729 | 3,376 |
| 2007 | 1,217      | 1,034   | 1,011                 | 1,278 | 730      | 1,166 | 70,126                 | 15,512 | 10,631   | 8,669  | 4,152 |

|                  |              |              |              |              |              |              |               |               |               |               |              |
|------------------|--------------|--------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|---------------|--------------|
| 2008             | 971          | 950          | 1,168        | 1,220        | 961          | 1,141        | 76,860        | 14,365        | 9,150         | 7,961         | 2,287        |
| 2009             | 915          | 1,022        | 1,173        | 1,338        | 1,049        | 954          | 89,557        | 17,194        | 10,158        | 6,477         | 4,194        |
| 2010             | 964          | 1,040        | 1,081        | 1,095        | 1,773        | 1,024        | 72,969        | 14,491        | 9,133         | 8,760         | 2,609        |
| 2011             | 1,008        | 1,001        | 1,363        | 1,369        | 730          | 979          | 74,916        | 14,463        | 7,026         | 6,387         | 2,601        |
| 2012             | 1,001        | 953          | 1,007        | 1,708        | 952          | 992          | 98,008        | 15,277        | 14,895        | 7,877         | 2,721        |
| 2013             | 1,113        | 1,150        | 1,430        | 973          | 1,209        | 925          | 73,062        | 14,538        | 15,330        | 12,814        | 1,957        |
| 2014             | 1,135        | 1,262        | 1,417        | 973          | 1,399        | 947          | 63,891        | 12,664        | 8,950         | 10,617        | 1,857        |
| 2015             | 1,180        | 1,095        | 1,417        | 2,281        | 1,186        | 956          | 53,746        | 11,771        | 11,406        | 11,041        | 1,962        |
| 2016             | 1,146        | 1,177        | 1,127        | 1,763        | 1,412        | 908          | 53,436        | 11,575        | 10,111        | 12,215        | 3,065        |
| 2017             | 841          | 1,038        | 1,189        | 1,916        | 1,095        | 905          | 58,178        | 11,664        | 6,665         | 9,712         | 952          |
| 2018             | 1,158        | 1,195        | 1,422        | 1,757        | 1,825        | 953          | 58,617        | 8,480         | 4,194         | 7,642         | 373          |
| <b>10 yr avg</b> | <b>1,093</b> | <b>1,046</b> | <b>1,263</b> | <b>1,517</b> | <b>1,263</b> | <b>954</b>   | <b>69,638</b> | <b>13,212</b> | <b>9,787</b>  | <b>9,354</b>  | <b>2,229</b> |
| <b>10 yr SD</b>  | <b>94</b>    | <b>111</b>   | <b>155</b>   | <b>415</b>   | <b>329</b>   | <b>35</b>    | <b>14,329</b> | <b>2,344</b>  | <b>3,308</b>  | <b>2,174</b>  | <b>1,022</b> |
| <b>20 yr avg</b> | <b>1,094</b> | <b>1,088</b> | <b>1,188</b> | <b>1,399</b> | <b>1,119</b> | <b>1,026</b> | <b>85,876</b> | <b>15,774</b> | <b>13,811</b> | <b>11,444</b> | <b>2,690</b> |
| <b>20 yr SD</b>  | <b>120</b>   | <b>156</b>   | <b>152</b>   | <b>330</b>   | <b>303</b>   | <b>82</b>    | <b>35,887</b> | <b>3,847</b>  | <b>8,724</b>  | <b>6,632</b>  | <b>1,674</b> |

| Year | Bottomfish |         | Coral Reef Boat-Based |       |          |       | Coral Reef Shore-Based |         |          |        |       |
|------|------------|---------|-----------------------|-------|----------|-------|------------------------|---------|----------|--------|-------|
|      | # Gears    | # Trips | Spear                 | SCUBA | Gill Net | Troll | H&L                    | Castnet | Gill net | Spear  | H&G   |
| 1982 | 798        | 40      | 949                   | 365   | 0        | 1,506 |                        |         |          |        |       |
| 1983 | 709        | 210     | 669                   | 477   | 0        | 1,428 |                        |         |          |        |       |
| 1984 | 847        | 242     | 1,391                 | 549   | 1,098    | 1,392 | 100,252                | 16,995  | 10,503   | 7,065  | 2,864 |
| 1985 | 883        | 857     | 1,191                 | 791   | 365      | 1,371 | 120,562                | 24,595  | 17,408   | 20,215 | 4,661 |
| 1986 | 794        | 633     | 834                   | 867   | 365      | 1,423 | 91,270                 | 18,289  | 21,959   | 19,236 | 3,847 |
| 1987 | 829        | 852     | 675                   | 863   | 1,241    | 1,489 | 108,016                | 21,759  | 25,008   | 16,672 | 3,320 |
| 1988 | 903        | 1,449   | 832                   | 703   | 366      | 1,479 | 99,458                 | 21,535  | 19,197   | 23,943 | 9,917 |
| 1989 | 869        | 1,338   | 973                   | 897   | 365      | 1,459 | 128,341                | 24,681  | 13,766   | 10,707 | 6,605 |
| 1990 | 883        | 943     | 933                   | 1,074 | 730      | 1,466 | 102,789                | 21,335  | 14,977   | 10,950 | 3,744 |

|                  |            |              |              |              |            |              |               |               |              |              |              |
|------------------|------------|--------------|--------------|--------------|------------|--------------|---------------|---------------|--------------|--------------|--------------|
| 1991             | 843        | 1,125        | 876          | 852          | 597        | 1,392        | 221,109       | 35,446        | 28,876       | 14,600       | 5,678        |
| 1992             | 886        | 945          | 866          | 839          | 471        | 1,447        | 193,008       | 33,219        | 35,056       | 18,287       | 4,073        |
| 1993             | 910        | 1,495        | 836          | 906          | 411        | 1,406        | 195,366       | 35,496        | 23,816       | 19,163       | 9,034        |
| 1994             | 947        | 1,520        | 898          | 947          | 791        | 1,378        | 238,436       | 29,565        | 21,809       | 18,068       | 4,015        |
| 1995             | 1,275      | 2,049        | 854          | 1,082        | 501        | 1,351        | 250,643       | 32,895        | 23,598       | 21,274       | 10,995       |
| 1996             | 1,562      | 1,754        | 880          | 1,075        | 673        | 1,399        | 264,597       | 21,048        | 15,331       | 15,994       | 9,944        |
| 1997             | 1,581      | 1,700        | 944          | 1,068        | 595        | 1,405        | 198,473       | 24,515        | 12,356       | 10,787       | 3,073        |
| 1998             | 1,305      | 2,209        | 798          | 1,113        | 487        | 1,346        | 159,600       | 33,840        | 21,840       | 22,260       | 7,260        |
| 1999             | 1,387      | 2,103        | 909          | 1,137        | 574        | 1,313        | 212,623       | 22,480        | 21,836       | 36,844       | 10,564       |
| 2000             | 1,321      | 1,750        | 919          | 1,053        | 712        | 1,361        | 128,937       | 16,941        | 11,085       | 15,738       | 3,817        |
| 2001             | 1,078      | 1,635        | 1,095        | 1,019        | 786        | 1,365        | 121,362       | 17,702        | 9,079        | 12,501       | 3,969        |
| 2002             | 1,037      | 1,230        | 793          | 995          | 584        | 1,321        | 93,984        | 16,914        | 6,337        | 7,688        | 1,258        |
| 2003             | 1,092      | 1,175        | 1,029        | 1,039        | 426        | 1,306        | 95,584        | 20,896        | 8,030        | 11,954       | 958          |
| 2004             | 1,121      | 1,013        | 969          | 1,198        | 366        | 1,320        | 85,809        | 13,034        | 7,839        | 10,484       | 708          |
| 2005             | 965        | 896          | 791          | 1,043        | 402        | 1,391        | 83,950        | 16,288        | 6,479        | 7,528        | 3,331        |
| 2006             | 956        | 863          | 1,037        | 657          | 365        | 1,365        | 75,783        | 20,349        | 8,623        | 12,182       | 3,376        |
| 2007             | 1,034      | 806          | 870          | 1,278        | 803        | 1,382        | 75,144        | 11,452        | 6,251        | 8,349        | 4,243        |
| 2008             | 950        | 953          | 1,084        | 1,037        | 549        | 1,340        | 75,945        | 13,679        | 4,849        | 7,869        | 2,287        |
| 2009             | 1,022      | 1,110        | 899          | 1,217        | 639        | 1,284        | 96,313        | 16,868        | 6,384        | 6,384        | 4,194        |
| 2010             | 1,040      | 1,316        | 946          | 1,095        | 365        | 1,201        | 78,654        | 13,326        | 5,638        | 8,294        | 2,656        |
| 2011             | 1,001      | 836          | 1,095        | 1,278        | 1,095      | 1,119        | 81,121        | 13,824        | 4,517        | 6,159        | 2,327        |
| 2012             | 953        | 767          | 961          | 1,586        | 366        | 1,099        | 105,408       | 14,369        | 9,548        | 7,877        | 2,721        |
| 2013             | 1,150      | 741          | 1,156        | 730          | 456        | 1,205        | 85,224        | 13,839        | 8,294        | 12,721       | 1,957        |
| 2014             | 1,262      | 702          | 1,353        | 608          | 608        | 1,251        | 69,461        | 12,426        | 5,523        | 10,236       | 1,857        |
| 2015             | 1,095      | 598          | 1,245        | 2,099        | 456        | 1,217        | 57,807        | 11,634        | 7,391        | 10,996       | 1,871        |
| 2016             | 1,177      | 783          | 1,010        | 1,647        | 471        | 1,163        | 60,344        | 11,255        | 7,686        | 12,215       | 3,065        |
| 2017             | 1,038      | 849          | 977          | 1,278        | 365        | 1,234        | 63,367        | 11,283        | 4,475        | 9,760        | 952          |
| 2018             | 1,195      | 697          | 1,153        | 1,460        | 1,825      | 1,178        | 64,768        | 8,014         | 2,796        | 7,642        | 373          |
| <b>10 yr avg</b> | <b>840</b> | <b>1,093</b> | <b>1,080</b> | <b>1,300</b> | <b>665</b> | <b>1,195</b> | <b>76,247</b> | <b>12,684</b> | <b>6,225</b> | <b>9,228</b> | <b>2,197</b> |
| <b>10 yr SD</b>  | <b>204</b> | <b>94</b>    | <b>139</b>   | <b>415</b>   | <b>439</b> | <b>54</b>    | <b>15,165</b> | <b>2,244</b>  | <b>1,934</b> | <b>2,197</b> | <b>1,020</b> |

|                  |              |              |              |              |            |              |               |               |              |               |              |
|------------------|--------------|--------------|--------------|--------------|------------|--------------|---------------|---------------|--------------|---------------|--------------|
| <b>20 yr avg</b> | <b>1,041</b> | <b>1,094</b> | <b>1,015</b> | <b>1,173</b> | <b>611</b> | <b>1,271</b> | <b>90,579</b> | <b>14,829</b> | <b>7,633</b> | <b>11,171</b> | <b>2,824</b> |
| <b>20 yr SD</b>  | <b>386</b>   | <b>120</b>   | <b>140</b>   | <b>338</b>   | <b>335</b> | <b>87</b>    | <b>33,498</b> | <b>3,587</b>  | <b>3,787</b> | <b>6,380</b>  | <b>2,119</b> |

### **1.2.11 Bycatch Estimates**

This section focuses on MSA § 303(a)(11), which requires that all FMPs establish a standardized reporting methodology to assess the amount and type of bycatch occurring in the fishery, and include conservation and management measures that, to the extent practicable, minimize bycatch and bycatch mortality. The MSA § 303(a)(11) standardized reporting methodology is commonly referred to as a “Standardized Bycatch Reporting Methodology” (SBRM) and was added to the MSA by the Sustainable Fisheries Act of 1996 (SFA). The Council implemented omnibus amendments to FMPs in 2003 to address MSA bycatch provisions and establish SBRMs.

The following are recent bycatch estimates for the boat-based non-bottomfishing gear (Table 38), bottomfish fishery (Table 39), and shore-based fisheries with all gears combined (Table 40).

**Calculations:** The number caught is the sum of the total number of individuals found in the raw data including bycatch. The number kept is the total number of individuals in the raw data that are not marked as bycatch. The number released is bycatch caught minus the number of bycatch kept. Percent bycatch is the sum of all bycatch divided by the total catch.

**Table 38. Time series of bycatch estimates in Guam boat-based non-bottomfishing fisheries from 1982-2018**

| <b>Year</b>      | <b># Caught</b> | <b>Kept</b>   | <b>Released</b> | <b>%<br/>Bycatch</b> |
|------------------|-----------------|---------------|-----------------|----------------------|
| 1982             | 5,388           | 5,388         | 0               | 0.0000               |
| 1983             | 3,581           | 3,581         | 0               | 0.0000               |
| 1984             | 5,584           | 5,584         | 0               | 0.0000               |
| 1985             | 8,138           | 8,138         | 0               | 0.0000               |
| 1986             | 4,829           | 4,829         | 0               | 0.0000               |
| 1987             | 4,895           | 4,895         | 0               | 0.0000               |
| 1988             | 8,113           | 8,113         | 0               | 0.0000               |
| 1989             | 12,393          | 12,393        | 0               | 0.0000               |
| 1990             | 7,645           | 7,645         | 0               | 0.0000               |
| 1991             | 9,338           | 9,338         | 0               | 0.0000               |
| 1992             | 7,352           | 7,352         | 0               | 0.0000               |
| 1993             | 9,398           | 9,398         | 0               | 0.0000               |
| 1994             | 9,843           | 9,843         | 0               | 0.0000               |
| 1995             | 17,776          | 17,776        | 0               | 0.0000               |
| 1996             | 20,931          | 20,931        | 0               | 0.0000               |
| 1997             | 19,108          | 19,108        | 0               | 0.0000               |
| 1998             | 16,428          | 16,428        | 0               | 0.0000               |
| 1999             | 19,827          | 19,827        | 0               | 0.0000               |
| 2000             | 23,373          | 23,335        | 38              | 0.0016               |
| 2001             | 10,409          | 10,344        | 65              | 0.0062               |
| 2002             | 5,560           | 5,520         | 40              | 0.0072               |
| 2003             | 8,543           | 8,538         | 5               | 0.0006               |
| 2004             | 5,851           | 5,839         | 12              | 0.0021               |
| 2005             | 4,012           | 4,006         | 6               | 0.0015               |
| 2006             | 7,176           | 7,172         | 4               | 0.0006               |
| 2007             | 5,611           | 5,538         | 73              | 0.0130               |
| 2008             | 9,199           | 9,198         | 1               | 0.0001               |
| 2009             | 11,710          | 11,707        | 3               | 0.0003               |
| 2010             | 8,588           | 8,588         | 0               | 0.0000               |
| 2011             | 21,232          | 21,231        | 1               | 0.0000               |
| 2012             | 12,200          | 12,200        | 0               | 0.0000               |
| 2013             | 11,834          | 11,806        | 28              | 0.0024               |
| 2014             | 8,814           | 8,789         | 25              | 0.0028               |
| 2015             | 8,995           | 8,995         | 0               | 0.0000               |
| 2016             | 11,031          | 11,025        | 6               | 0.0005               |
| 2017             | 8,645           | 8,643         | 2               | 0.0002               |
| 2018             | 9,580           | 9,578         | 2               | 0.0002               |
| <b>10 yr avg</b> | <b>11,263</b>   | <b>11,256</b> | <b>7</b>        | <b>0.0006</b>        |



|                  |               |               |           |               |
|------------------|---------------|---------------|-----------|---------------|
| <b>10 yr SD</b>  | <b>3,589</b>  | <b>3,590</b>  | <b>10</b> | <b>0.0010</b> |
| <b>20 yr avg</b> | <b>10,610</b> | <b>10,594</b> | <b>16</b> | <b>0.0020</b> |
| <b>20 yr SD</b>  | <b>5,078</b>  | <b>5,080</b>  | <b>22</b> | <b>0.0032</b> |

**Table 39. Time series of bycatch estimates in the Guam bottomfish fishery from 1982-2018**

| <b>Year</b> | <b># Caught</b> | <b>Kept</b> | <b>Released</b> | <b>% Bycatch</b> |
|-------------|-----------------|-------------|-----------------|------------------|
| 1982        | 1,597           | 1,597       | 0               | 0.0000           |
| 1983        | 1,507           | 1,507       | 0               | 0.0000           |
| 1984        | 3,347           | 3,347       | 0               | 0.0000           |
| 1985        | 4,840           | 4,840       | 0               | 0.0000           |
| 1986        | 1,624           | 1,624       | 0               | 0.0000           |
| 1987        | 2,519           | 2,519       | 0               | 0.0000           |
| 1988        | 3,002           | 3,002       | 0               | 0.0000           |
| 1989        | 3,562           | 3,562       | 0               | 0.0000           |
| 1990        | 2,870           | 2,870       | 0               | 0.0000           |
| 1991        | 2,783           | 2,783       | 0               | 0.0000           |
| 1992        | 2,527           | 2,527       | 0               | 0.0000           |
| 1993        | 2,893           | 2,893       | 0               | 0.0000           |
| 1994        | 3,730           | 3,730       | 0               | 0.0000           |
| 1995        | 4,985           | 4,985       | 0               | 0.0000           |
| 1996        | 5,244           | 5,244       | 0               | 0.0000           |
| 1997        | 4,342           | 4,342       | 0               | 0.0000           |
| 1998        | 5,138           | 5,138       | 0               | 0.0000           |
| 1999        | 4,938           | 4,938       | 0               | 0.0000           |
| 2000        | 3,905           | 3,373       | 532             | 0.1362           |
| 2001        | 3,896           | 3,273       | 623             | 0.1599           |
| 2002        | 2,504           | 2,151       | 353             | 0.1410           |
| 2003        | 1,888           | 1,697       | 191             | 0.1012           |
| 2004        | 1,804           | 1,682       | 122             | 0.0676           |
| 2005        | 1,706           | 1,640       | 66              | 0.0387           |
| 2006        | 2,188           | 2,043       | 145             | 0.0663           |
| 2007        | 1,372           | 1,233       | 139             | 0.1013           |
| 2008        | 1,657           | 1,536       | 121             | 0.0730           |
| 2009        | 2,851           | 2,774       | 77              | 0.0270           |
| 2010        | 2,588           | 2,559       | 29              | 0.0112           |
| 2011        | 2,128           | 2,083       | 45              | 0.0211           |
| 2012        | 924             | 887         | 37              | 0.0400           |
| 2013        | 1,222           | 1,178       | 44              | 0.0360           |
| 2014        | 2,452           | 2,283       | 169             | 0.0689           |
| 2015        | 1,420           | 1,350       | 70              | 0.0493           |
| 2016        | 1,674           | 1,627       | 47              | 0.0281           |

|                  |              |              |            |               |
|------------------|--------------|--------------|------------|---------------|
| 2017             | 2,313        | 2,287        | 26         | 0.0112        |
| 2018             | 1,521        | 1,492        | 29         | 0.0191        |
| <b>10 yr avg</b> | <b>1,909</b> | <b>1,852</b> | <b>57</b>  | <b>0.0312</b> |
| <b>10 yr SD</b>  | <b>612</b>   | <b>600</b>   | <b>41</b>  | <b>0.0171</b> |
| <b>20 yr avg</b> | <b>2,248</b> | <b>2,104</b> | <b>143</b> | <b>0.0599</b> |
| <b>20 yr SD</b>  | <b>985</b>   | <b>919</b>   | <b>165</b> | <b>0.0454</b> |

Table 40. Time series of bycatch estimates in the Guam shore-based fishery from 1984-2018

| Year | # Caught | Kept   | Released | % Bycatch |
|------|----------|--------|----------|-----------|
| 1984 | 1,845    | 1,845  | 0        | 0.0000    |
| 1985 | 10,200   | 10,200 | 0        | 0.0000    |
| 1986 | 9,172    | 9,169  | 3        | 0.0003    |
| 1987 | 9,860    | 9,860  | 0        | 0.0000    |
| 1988 | 16,199   | 16,199 | 0        | 0.0000    |
| 1989 | 8,802    | 8,802  | 0        | 0.0000    |
| 1990 | 8,817    | 8,817  | 0        | 0.0000    |
| 1991 | 9,880    | 9,880  | 0        | 0.0000    |
| 1992 | 6,753    | 6,753  | 0        | 0.0000    |
| 1993 | 30,916   | 30,916 | 0        | 0.0000    |
| 1994 | 6,013    | 6,013  | 0        | 0.0000    |
| 1995 | 8,360    | 8,360  | 0        | 0.0000    |
| 1996 | 3,385    | 3,385  | 0        | 0.0000    |
| 1997 | 9,233    | 9,216  | 17       | 0.0018    |
| 1998 | 11,589   | 11,580 | 9        | 0.0008    |
| 1999 | 12,592   | 12,530 | 62       | 0.0049    |
| 2000 | 7,861    | 7,831  | 30       | 0.0038    |
| 2001 | 8,653    | 8,593  | 60       | 0.0069    |
| 2002 | 3,122    | 3,114  | 8        | 0.0026    |
| 2003 | 5,364    | 5,345  | 19       | 0.0035    |
| 2004 | 2,655    | 2,611  | 44       | 0.0166    |
| 2005 | 2,684    | 2,654  | 30       | 0.0112    |
| 2006 | 3,928    | 3,851  | 77       | 0.0196    |
| 2007 | 3,361    | 3,238  | 123      | 0.0366    |
| 2008 | 5,359    | 5,282  | 77       | 0.0144    |
| 2009 | 3,254    | 3,160  | 94       | 0.0289    |
| 2010 | 4,321    | 4,222  | 99       | 0.0229    |
| 2011 | 5,262    | 5,187  | 75       | 0.0143    |
| 2012 | 5,590    | 5,559  | 31       | 0.0055    |
| 2013 | 3,300    | 2,893  | 407      | 0.1233    |
| 2014 | 4,732    | 4,622  | 110      | 0.0232    |

|                  |              |              |            |               |
|------------------|--------------|--------------|------------|---------------|
| 2015             | 4,823        | 4,775        | 48         | 0.0100        |
| 2016             | 3,907        | 3,785        | 122        | 0.0312        |
| 2017             | 7,804        | 7,798        | 6          | 0.0008        |
| 2018             | 16,981       | 16,965       | 16         | 0.0009        |
| <b>10 yr avg</b> | <b>5,998</b> | <b>5,897</b> | <b>101</b> | <b>0.0261</b> |
| <b>10 yr SD</b>  | <b>3,868</b> | <b>3,917</b> | <b>109</b> | <b>0.0341</b> |
| <b>20 yr avg</b> | <b>5,778</b> | <b>5,701</b> | <b>77</b>  | <b>0.0191</b> |
| <b>20 yr SD</b>  | <b>3,501</b> | <b>3,526</b> | <b>84</b>  | <b>0.0261</b> |

### 1.2.12 Number of Federal Permit Holders

In Guam, the following Federal permits are required for fishing in the EEZ:

#### 1.2.12.1 Guam Large Vessel Bottomfish Permit

The Code of Federal Regulations (CFR), Title 50, Part 665 requires the following Federal permits for Guam fisheries in the exclusive economic zone (EEZ) under the Mariana FEP:

#### 1.2.12.2 Guam Large Vessel Bottomfish Permit

Regulations require this permit for any large vessel (50 feet or longer in overall length) fishing for, landing, or transshipping bottomfish MUS or bottomfish ecosystem component species (ECS) in the EEZ seaward of Guam.

#### 1.2.12.3 Special Coral Reef Ecosystem Permit

Regulations require the coral reef ecosystem special permit for anyone fishing for coral reef ECS in a low-use marine protected area (MPA), fishing for species on the list of Potentially Harvested Coral Reef Taxa, or using fishing gear not specifically allowed in the regulations. NMFS will make an exception to this permit requirement for any person issued a permit to fish under any fishery ecosystem plan who incidentally catches Guam coral reef ECSS while fishing for bottomfish MUS, crustacean ECS, western Pacific pelagic MUS, precious coral, or seamount groundfish. Regulations require a transshipment permit for any receiving vessel used to land or transship potentially harvested coral reef taxa, or any coral reef ecosystem ECS caught in a low-use MPA.

#### 1.2.12.4 Western Pacific Precious Corals Permit

Regulations require this permit for anyone harvesting or landing black, bamboo, pink, red, or gold corals in the EEZ in the Western Pacific.

#### 1.2.12.5 Western Pacific Crustaceans Permit (Lobster or Deepwater Shrimp)

Regulations require a permit by the owner of a U.S. fishing vessel used to fish for lobster or deepwater shrimp in the EEZ around American Samoa, Guam, Commonwealth of the Northern Mariana Islands (CNMI), Hawaii, and the Pacific Remote Islands Areas (PRIA).

There is no record of special coral reef or precious coral fishery permits issued for the EEZ around Guam since 2007. Table 41 provides the number of permits issued for Guam fisheries

between 2009 and 2018. Historical data are from the PIFSC and 2018 data are from the PIRO Sustainable Fisheries Division permits program as of February 27, 2019.

**Table 41. Number of federal permits holders for the crustacean and bottomfish fisheries of Guam from 2009-2018**

| Guam Fisheries | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|----------------|------|------|------|------|------|------|------|------|------|------|
| Lobster        | 4*   | 0    | 0    | 0    | 0    | 0    | 0    | 1**  | 0    | 1**  |
| Shrimp         | 0    | 2*   | 1*   | 0    | 0    | 0    | 0    | 1    | 0    | 0    |
| Bottomfish     | 2    | 1    | 1    | 4    | 2    | 2    | 1    | 1    | 1    | 1    |

\*Permits apply to multiple areas and may include American Samoa, Guam, CNMI, and PRIA.

\*\*Area 5 CNMI and Guam

## 1.2.13 Status Determination Criteria

### 1.2.13.1 Bottomfish Fishery

Overfishing criteria and control rules are specified and applied to individual species within the multi-species stock whenever possible. When this is not possible, they are based on an indicator species for the multi-species stock. It is important to recognize that individual species would be affected differently based on this type of control rule, and it is important that for any given species fishing, mortality does not currently exceed a level that would result in excessive depletion of that species. No indicator species are being used for the bottomfish multi-species stock complexes and the coral reef species complex. Instead, the control rules are applied to each stock complex as a whole.

The MSY control rule is used as the maximum fishing mortality threshold (MFMT). The MFMT and minimum stock size threshold (MSST) are specified based on recommendations in Restrepo et al. (1998) and both are dependent on the natural mortality rate (M) (Table 42). The value of M used to determine the reference point values are not specified in this document. The latest estimate, published annually in the SAFE report, is used and the value is occasionally re-estimated using the best available information. The range of M among species within a stock complex is taken into consideration when estimating and choosing the M to be used for the purpose of computing the reference point values.

In addition to the thresholds MFMT and MSST, a warning reference point,  $B_{FLAG}$ , is specified at some point above the MSST to provide a trigger for consideration of management action prior to B reaching the threshold. MFMT, MSST, and  $B_{FLAG}$  are specified as indicated in Table 44.

**Table 42. Overfishing threshold specifications for Guam BMUS**

| MFMT   | MSST        | $B_{FLAG}$ |
|--|-------------|------------|
| $F(B) = \frac{F_{MSY} B}{c B_{MSY}} \quad \text{for } B \leq c B_{MSY}$ $F(B) = F_{MSY} \quad \text{for } B > c B_{MSY}$ | $c B_{MSY}$ | $B_{MSY}$  |

$$\text{where } c = \max(1-M, 0.5)$$

Standardized values of fishing effort (E) and catch-per-unit-effort (CPUE) are used as proxies for F and B, respectively, so  $E_{MSY}$ ,  $CPUE_{MSY}$ , and  $CPUE_{FLAG}$  are used as proxies for  $F_{MSY}$ ,  $B_{MSY}$ , and  $B_{FLAG}$ , respectively.

In cases where reliable estimates of  $CPUE_{MSY}$  and  $E_{MSY}$  are not available, they will be estimated from catch and effort time series, standardized for all identifiable biases.  $CPUE_{MSY}$  would be calculated as half of a multi-year average reference CPUE, called  $CPUE_{REF}$ . The multi-year reference window would be objectively positioned in time to maximize the value of  $CPUE_{REF}$ .  $E_{MSY}$  would be calculated using the same approach or, following Restrepo et al. (1998), by setting  $E_{MSY}$  equal to  $E_{AVE}$ , where  $E_{AVE}$  represents the long-term average effort prior to declines in CPUE. When multiple estimates are available, the more precautionary one is used.

Since the MSY control rule specified here applies to multi-species stock complexes, it is important to ensure that no particular species within the complex has a mortality rate that leads to excessive depletion. In order to accomplish this, a secondary set of reference points is specified to evaluate stock status with respect to recruitment overfishing. A secondary “recruitment overfishing” control rule is specified to control fishing mortality with respect to that status. The rule applies only to those component stocks (species) for which adequate data are available. The ratio of a current spawning stock biomass proxy ( $SSB_{Pt}$ ) to a given reference level ( $SSB_{PREF}$ ) is used to determine if individual stocks are experiencing recruitment overfishing.  $SSB_{Pt}$  is CPUE scaled by percent mature fish in the catch. When the ratio  $SSB_{Pt}/SSB_{PREF}$ , or the “SSBP ratio” ( $SSBPR$ ) for any species drops below a certain limit ( $SSBPR_{MIN}$ ), that species is considered to be recruitment overfished and management measures will be implemented to reduce fishing mortality on that species. The rule applies only when the SSBP ratio drops below the  $SSBPR_{MIN}$ , but it will continue to apply until the ratio achieves the “SSBP ratio recovery target” ( $SSBPR_{TARGET}$ ), which is set at a level no less than  $SSBPR_{MIN}$ . These two reference points and their associated recruitment overfishing control rule, which prescribe a target fishing mortality rate ( $F_{RO-REBUILD}$ ) as a function of the SSBP ratio, are specified as indicated in Table 43. Again,  $E_{MSY}$  is used as a proxy for  $F_{MSY}$ .

**Table 43. Rebuilding control rules for Guam BMUS**

| $F_{RO-REBUILD}$   | $SSBPR_{MIN}$ | $SSBPR_{TARGET}$ |
|--|---------------|------------------|
| $F(SSBPR) = 0$ for $SSBPR \leq 0.10$                                   | 0.20          | 0.30             |
| $F(SSBPR) = 0.2 F_{MSY}$ for $0.10 < SSBPR \leq SSBPR_{MIN}$           |               |                  |
| $F(SSBPR) = 0.5 F_{MSY}$ for $SSBPR_{MIN} < SSBPR \leq SSBPR_{TARGET}$ |               |                  |

### 1.2.13.2 Coral Reef Fishery

Available biological and fishery data are poor for all coral reef ecosystem management unit species in the Mariana Islands. There is scant information on the life histories, ecosystem

dynamics, fishery impact, community structure changes, yield potential, and management reference points for many coral reef ecosystem species. Additionally, total fishing effort cannot be adequately partitioned between the various management unit species (MUS) for any fishery or area. Biomass, maximum sustainable yield, and fishing mortality estimates are not available for any single MUS. Once these data are available, fishery managers can establish limits and reference points based on the multi-species coral reef ecosystem as a whole.

When possible, the MSY control rule should be applied to the individual species in a multi-species stock. When this is not possible, MSY may be specified for one or more species; these values can then be used as indicators for the multi-species stock's MSY.

Individual species that are part of a multi-species complex will respond differently to an OY-determined level of fishing effort ( $F_{OY}$ ). Thus, for a species complex that is fished at  $F_{OY}$ , managers still must track individual species' mortality rates in order to prevent species-specific population declines that would lead to depletion.

For the coral reef fishery, the multi-species complex as a whole is used to establish limits and reference points for each area. When possible, available data for a particular species are used to evaluate the status of individual MUS stocks in order to prevent recruitment overfishing. When better data and the appropriate multi-species stock assessment methodologies become available, all stocks will be evaluated independently, without proxy.

#### 1.2.13.2.1 Establishing Reference Point Values

Standardized values of catch per unit effort (CPUE) and effort (E) are used to establish limit and reference point values, which act as proxies for relative biomass and fishing mortality, respectively. Limits and reference points are calculated in terms of  $CPUE_{MSY}$  and  $E_{MSY}$  included in Table 44.

**Table 44. Status determination criteria for the coral reef management unit species using CPUE based proxies**

| Value                | Proxy             | Explanation                                     |
|----------------------|-------------------|---|
| MaxFMT ( $F_{MSY}$ ) | $E_{MSY}$         | $0.91 CPUE_{MSY}$                               |
| $F_{OY}$             | $0.75 E_{MSY}$    | suggested default scaling for target            |
| $B_{MSY}$            | $CPUE_{MSY}$      | operational counterpart                         |
| $B_{OY}$             | $1.3 CPUE_{MSY}$  | simulation results from Mace (1994)             |
| MinSST               | $0.7 CPUE_{MSY}$  | suggested default $(1-M)B_{MSY}$ with $M=0.3^*$ |
| $B_{FLAG}$           | $0.91 CPUE_{MSY}$ | suggested default $(1-M)B_{OY}$ with $M=0.3^*$  |

When reliable estimates of  $E_{MSY}$  and  $CPUE_{MSY}$  are not available, they are generated from time series of catch and effort values, standardized for all identifiable biases using the best available

analytical tools.  $CPUE_{MSY}$  is calculated as one-half a multi-year moving average reference CPUE ( $CPUE_{REF}$ ).

### 1.2.13.3 Current Stock Status

#### 1.2.13.3.1 Bottomfish

Biological and other fishery data are poor for all bottomfish species in the Mariana Archipelago. Generally, data are only available on commercial landings by species and catch-per-unit-effort (CPUE) for the multi-species complexes as a whole. At this time it is not possible to partition these effort measures among the various bottomfish MUS. The most recent stock assessment update (Yau et al, 2015) for the Guam bottomfish management unit species complex (comprised of 17 species of shallow and deep species of snapper, grouper, jacks, and emperors) was based on estimate of total catch, an abundance index derived from the nominal CPUE generated from the creel surveys, and a fishery-independent point estimate of MSY from the Our Living Oceans Report (Humphreys and Moffitt, 1999, Moffitt and Humphreys, 2009). The assessment utilized a state-space surplus production model with explicit process and observation error terms (Meyer and Millar, 1999). Determinations of overfishing and overfished status can then be made by comparing current biomass and harvest rates to MSY level reference points. To date, the Guam BMUS is not subject to overfishing and is not overfished (Table 45).

**Table 45. Stock assessment parameters for the Guam BMUS complex (Yau et al., 2015)**

| Parameter   | Value             | Notes                                       | Status                   |
|-------------|-------------------|---|--------------------------|
| MSY         | $56.13 \pm 7.79$  | Expressed in 1000 lbs. ( $\pm$ std. error)  |                          |
| $H_{2013}$  | 0.123             | Expressed in percentage                     |                          |
| $H_{MSY}$   | $0.352 \pm 0.059$ | Expressed in percentage ( $\pm$ std. error) |                          |
| $H/H_{MSY}$ | 0.356             |   | No overfishing occurring |
| $B_{2013}$  | 264.7             | Expressed in thousand pounds                |                          |
| $B_{MSY}$   | $162.3 \pm 23.8$  | Expressed in 1000 lbs. ( $\pm$ std. error)  |                          |
| $B/B_{MSY}$ | 1.63              |   | Not overfished           |

#### 1.2.13.3.2 Coral reef

The application of the SDCs for the management unit species in the coral reef fisheries is limited due to various challenges. First, the thousands of species included in the coral reef MUS makes the SDC and status determination impractical. Second, the CPUE derived from the creel survey is based on the fishing method and there is no species-specific CPUE information available. In order to allocate the fishing method level CPUE to individual species, the catch data (the value of catch is derived from CPUE hence there is collinearity) will have to be identified to species level and CPUE will be parsed out by species composition. The third challenge is that there is very little species-level identification applied to the creel surveys. There has been no attempt to estimate MSY for the coral reef MUS until the 2007 re-authorization of MSA that requires the Council to specify ACLs for species in the FEPs.

For ACL specification purposes, MSYs in the coral reef fisheries are determined by using the Biomass-Augmented Catch-MSY approach (Sabater and Kleiber, 2014). This method estimates MSY using plausible combination rates of population increase (denoted by  $r$ ) and carrying capacity (denoted by  $k$ ) assumed from the catch time series, resilience characteristics (from FishBase), and biomass from existing underwater census surveys done by the PIFSC. This method was applied to species complexes grouped by taxonomic families. The most recent MSY estimates are found in Table 46. The SSC utilized the MSYs for the coral reef MUS complexes as the OFLs.

**Table 46. Best available MSY estimates for the coral reef MUS in Guam**

| <b>Coral Reef MUS Complex</b>  | <b>MSY (lbs.)</b> |
|--|-------------------|
| <i>Selar crumenophthalmus</i> – atulai or bigeye scad  | 61,300            |
| Acanthuridae – surgeonfish   | 118,000           |
| Carangidae – jacks   | 31,700            |
| Crustaceans – crabs  | 8,600             |
| Holocentridae – squirrelfish   | 13,900            |
| Kyphosidae – chubs/rudderfish  | 10,300            |
| Labridae – wrasses <sup>1</sup>  | 28,500            |
| Lethrinidae – emperors   | 78,000            |
| Lutjanidae – snappers  | 21,800            |
| Mollusks – turbo snail; octopus; giant clams   | 29,000            |
| Mugilidae – mullets  | 26,200            |
| Mullidae – goatfish  | 16,400            |
| Scaridae – parrotfish <sup>2</sup>   | 87,100            |
| Serranidae – groupers  | 28,600            |
| Siganidae – rabbitfish   | 19,700            |
| All Other CREMUS Combined (other coral reef ecosystem finfish, other invertebrates, misc. bottomfish, misc. reef fish, and misc. shallow bottomfish) | 211,300           |
| <i>Cheilinus undulatus</i> – humphead (Napoleon) wrasse  | N.A.              |
| <i>Bolbometopon muricatum</i> – bumphead parrotfish  | N.A.              |
| Carcharhinidae – reef sharks   | 2,900             |

## 1.2.14 Overfishing Limit, Acceptable Biological Catch, and Annual Catch Limits

### 1.2.14.1 Brief Description of the ACL Process

The Council developed a Tiered system of control rules to guide the specification of ACLs and Accountability Measures (AMs) (WPRFMC, 2011). The process starts with the use of the best scientific information available (BSIA) in the form of, but not limited to, stock assessments, published paper, reports, or available data. These information are classified to the different Tiers in the control rule ranging from Tier 1 (most information available typically an assessment) to Tier 5 (catch-only information). The control rules are applied to the BSIA. Tiers 1 to 3 would involve conducting a Risk of Overfishing Analysis (denoted by  $P^*$ ) to quantify the scientific uncertainties around the assessment to specify the Acceptable Biological Catch (ABC). This would lower the ABC from the OFL (MSY-based). A Social, Ecological, Economic, and



Management (SEEM) Uncertainty Analysis is performed to quantify the uncertainties from the SEEM factors. The buffer is used to lower the ACL from the ABC. For Tier 4, which is comprised of stocks with MSY estimates but no active fisheries, the control rule is 91% of MSY. For Tier 5 which has catch-only information, the control rule is a third reduction in the median catch depending on the qualitative evaluation on what the stock status is based on expert opinion. ACL specification can choose from a variety of method including the above-mentioned SEEM analysis or a percentage buffer (% reduction from ABC based on expert opinion) or the use of an Annual Catch Target. Specifications are done on an annual basis but the Council normally specifies a multi-year specification.

The Accountability Measure for the coral reef and bottomfish fisheries in Guam is an overage adjustment. The ACL is downward adjusted with the amount of overage from the ACL based on a three-year running average.

#### 1.2.14.2 Current OFL, ABC, ACL, and Recent Catch

The most recent multiyear specification of OFL, ABC, and ACL for the coral reef fishery was completed in the 160<sup>th</sup> Council meeting on June 25 to 27, 2014. The specification covers fishing year 2015, 2016, 2017, and 2018 for the coral reef MUS complexes. A P\* and SEEM analysis was performed for this multiyear specification (NMFS 2015). For the bottomfish, it was a roll over from the previous specification since an assessment update was not available for fishing year 2015. ACLs were not specified by NMFS for the coral reef ecosystem MUS because NMFS has recently acquired new information that require additional environmental analyses to support the Council's ACL recommendations for these management unit species (50 CFR Part 665).

**Table 47. Guam ACL table with three-year average catch (lbs.)**

| <b>Fishery</b> | <b>MUS</b>                        | <b>OFL</b> | <b>ABC</b> | <b>ACL</b> | <b>Catch</b> |
|----------------|-----------------------------------|------------|------------|------------|--------------|
| Bottomfish     | Bottomfish multi-species complex  | 71,000     | 66,000     | 66,000     | 27,022       |
| Crustacean     | Deepwater shrimp                  | N.A.F.     | 48,488     | 48,488     | N.A.F.       |
|                | Spiny lobster                     | 4,600      | 3,300      | 3,135      | 446          |
|                | Slipper lobster                   | N.A.F.     | 20         | 20         | N.D.         |
|                | Kona crab                         | N.A.F.     | 1,900      | 1,900      | N.A.F.       |
| Precious coral | Black coral                       | 8,250      | 700        | 700        | N.A.F.       |
|                | Precious coral in CNMI expl. area | N.A.F.     | 2,205      | 2,205      | N.A.F.       |

The catch shown in Table 47 takes the average of the recent three years as recommended by the Council at its 160<sup>th</sup> meeting to avoid large fluctuations in catch due to data quality and outliers. "N.A.F." indicates no active fisheries as of date. "N.D." indicates no data.

The ACL for jacks was reduced from 29,300 lbs. in 2015 to 21,201 lbs. for 2016 due to the overage in 2015 of 8,099 lbs. because of the spike in catch in 2013 of 59,468 lbs. NMFS applied the reduction to the ACL by the amount of the overage (82 FR 5517 2017-01-18) based on the Council's accountability measure for this data poor stock.

## 1.2.15 Best Scientific Information Available

### 1.2.15.1 Bottomfish fishery

#### 1.2.15.1.1 Stock assessment benchmark

The benchmark stock assessment for the Territory Bottomfish Management Unit Species complex was developed and finalized in October 2007 (Moffitt et al., 2007). This benchmark utilized a Bayesian statistical framework to estimate parameters of a Schaefer model fit to a time series of annual CPUE statistics. The surplus production model included process error in biomass production dynamics and observation error in the CPUE data. This was an improvement to the previous approach of using index-based proxies for  $B_{MSY}$  and  $F_{MSY}$ . Best available information for the bottomfish stock assessment is as follows:

**Input data:** The CPUE and catch data used were from the Guam off-shore creel survey. The catch and CPUE were expanded on an annual level. CPUE was expressed in line-hours. The data was screened for trips that landed more than 50% BMUS species using the handline gear.

**Model:** state-space model with explicit process and observation error terms (see Meyer and Millar, 1999).

**Fishery independent source for biomass:** point estimate of MSY from the Our Living Oceans Report (Humphreys and Moffitt, 1999; Moffitt and Humphreys, 2009).

#### 1.2.15.1.2 Stock Assessment Updates

Updates to the 2007 benchmark were done in 2012 (Brodziak et al., 2012) and 2015 (Yau et al., 2015). These included a three-year stock projection table used for selecting the level of risk the fishery will be managed under ACLs. Yau et al. (2015) is considered the best scientific information available for the Territory bottomfish MUS complex after undergoing a WPSAR Tier 3 panel review (Franklin et al., 2015). This was the basis for the P\* analysis and SEEM analysis that determined the risk levels to specify ABCs and ACLs.

#### 1.2.15.1.3 Other Information Available

Approximately every five years PIFSC administers a socioeconomic survey to small boat fishermen in Guam. This survey consists of about 60 questions regarding a variety of topics, including fishing experiences, market participation, vessels and gear, demographics and household income, and fishermen perspectives. The survey requests participants to identify which MUS they primarily targeted during the previous 12 months, by percentage of trips. Full reports of these surveys can be found at the PIFSC Socioeconomics webpage (Hospital and Beavers, 2011).

### 1.2.15.2 Coral Reef Fishery

#### 1.2.15.2.1 Stock Assessment Benchmark

No stock assessment has been generated for the coral reef fisheries. The SDCs using index-based proxies were tested for its applicability in the different MUS in the coral reef fisheries (Hawhee, 2007). This analysis was done on a gear level. It paints a dire situation for the shore-based fishery with 43% of the gear/species combination falling below  $B_{flag}$  and 33% below MSST with

most catch and CPUE trends showing a decline over time. The off-shore fisheries were shown to be less dire with 50% of the gear/species combination falling below  $B_{flag}$  and 38% below MSST - but the catch and CPUE trends were increasing over time. The inconsistency in the CPUE and catch trends with the SDC results makes this type of assessment to be unreliable.

The first attempt to use a model-based approach in assessing the coral reef MUS complexes was done in 2014 using a biomass-based population dynamics model (Sabater and Kleiber, 2014). This model was based on the original Martell and Froese (2012) model but was augmented with biomass information to relax the assumption behind carrying capacity. It estimates MSY based on a range of rate of population growth ( $r$ ) and carrying capacity ( $k$ ) values. The best available information for the coral reef stock assessment is as follows:

**Input data:** The catch data was derived from the inshore and off-shore creel surveys. Commercial receipt book information was also used in combination with the creel data. A downward adjustment was done to address for potential overlap due to double reporting.

**Model:** Biomass Augmented Catch MSY approach based on the original catch-MSY model (Martell and Froese, 2012; Sabater and Kleiber, 2014).

**Fishery independent source for biomass:** biomass density from the Rapid Assessment and Monitoring Program of NMFS-CREP was expanded to the hard bottom habitat from 0-30 m (Williams, 2010).

This model had undergone a CIE review in 2014 (Cook, 2014; Haddon, 2014; Jones, 2014). This was the basis for the P\* analysis that determined the risk levels to specify ABCs.

#### **1.2.15.2.2 Stock Assessment Updates**

No updates available for the coral reef MUS complex. However, NMFS-PIFSC is finalizing a length-based model for estimating sustainable yield levels and various biological reference points (Nadon et al., 2015). This can be used on a species level. The Council is also working with a contractor to enhance the BAC-MSY model to incorporate catch, biomass, CPUE, effort, and length-based information in an integrated framework (Martell, 2015).

#### **1.2.15.2.3 Other Information Available**

Approximately every five years PIFSC administers a socioeconomic survey to small boat fishermen in Guam. This survey consists of about 60 questions regarding a variety of topics, including fishing experiences, market participation, vessels and gear, demographics and household income, and fishermen perspectives. The survey requests participants to identify which MUS they primarily targeted during the previous 12 months, by percentage of trips. Full reports of these surveys can be found at the PIFSC Socioeconomics webpage (Hospital and Beavers, 2011).

PIFSC and the Council conducted a workshop with various stakeholders in CNMI to identify factors and quantify uncertainties associated with the social, economic, ecological, and management of the coral reef fisheries (Sievanen and McCaskey, 2014). This was the basis for the SEEM analysis that determined the risk levels to specify ACLs.

### 1.2.16 Harvest Capacity and Extent

The MSA defines the term “optimum,” with respect to the yield from a fishery, as the amount of fish that:

- Will provide the greatest overall benefit to the Nation, particularly with respect to food production and recreational opportunities, and taking into account the protection of marine ecosystems.
- Is prescribed on the basis of the MSY from the fishery, as reduced by any relevant social, economic, or ecological factor.
- In the case of an overfished fishery, provides for rebuilding to a level consistent with producing the MSY in such a fishery [50 CFR §600.310(f)(1)(i)].

Optimum yield in the coral reef and bottomfish fisheries is prescribed based on the MSY from the stock assessment and the best available scientific information. In the process of specifying ACLs, social, economic, and ecological factors were considered and the uncertainties around those factors defined the management uncertainty buffer between the ABC and ACL. OY for the bottomfish and coral reef fish MUS complexes is defined to be the level of harvest equal to the ACL consistent with the goals and objectives of the Fishery Ecosystem Plans and used by the Council to manage the stock.

The Council recognizes that MSY and OY are long-term values whereas the ACLs are yearly snapshots based on the level of fishing mortality at  $F_{MSY}$ . There are situations when the long-term means around MSY are going to be lower than ACLs especially if the stock is known to be productive or relatively pristine or lightly fished. One can have catch levels and catch rates exceeding that of MSY over short-term enough to lower the biomass to a level around the estimated MSY and still not jeopardize the stock. This situation is true for the territory bottomfish multi-species complex.

The harvest extent, in this case, is defined as the level of catch harvested in a fishing year relative to the ACL or OY. The harvest capacity is the level of catch remaining in the annual catch limit that can potentially be used for TALLF. Table 48 summarizes the harvest extent and harvest capacity information for Guam in 2017.

**Table 48. Guam 2018 ACL proportion of harvest extent and the harvest capacity**

| Fishery        | MUS                               | ACL    | Catch  | Harvest extent (%) | Harvest capacity (%) |
|----------------|-----------------------------------|--------|--------|--------------------|----------------------|
| Bottomfish     | Bottomfish multi-species complex  | 66,000 | 27,022 | 40.9               | 59.1                 |
| Crustacean     | Deepwater shrimp                  | 48,488 | N.A.F. | 0                  | 100                  |
|                | Spiny lobster                     | 3,135  | 446    | 14.2               | 85.8                 |
|                | Slipper lobster                   | 20     | N.D.   |                    |                      |
|                | Kona crab                         | 1,900  | N.A.F. | 0                  | 100                  |
| Precious coral | Black coral                       | 700    | N.A.F. | 0                  | 100                  |
|                | Precious coral in CNMI expl. area | 2,205  | N.A.F. | 0                  | 100                  |

## 1.2.17 Other Relevant Ocean-Uses and Fishery-Related Information

### 1.2.17.1 Marine Preserves

Guam has five locally managed Marine Preserves (MPAs): Achang Reef Flat in Merizo, Sasa Bay in Piti, Piti Bombholes in Piti, Tumon Bay in Tumon, and Pati Point in Yigo. A total of 11.8% of Guam's coastline is located within the MPAs.

### 1.2.17.2 Local Environmental Co-Variates

In early 2010, the U.S. military began exercises in an area south and southeast of Guam designated W-517. W-517 is a special use airspace (SUA) (approximately 14,000 nm<sup>2</sup>) that overlays deep open ocean approximately 50 miles south-southwest of Guam. Exercises in W-517 generally involve live fire and/or pyrotechnics. When W-517 is in use, a notice to mariners (NTM) is issued, and vessels attempting to use the area are advised to be cautious of objects in the water and other small vessels. This discourages access to virtually all banks south of Guam, including Galvez, Santa Rosa, White Tuna, and other popular fishing areas. From 1982-2015, DAWR surveys recorded more than 2930 trolling and bottom fishing trips to these southern banks, an average of more than 83 trips per year. The number of NTM in 2016 was 64, equaling 123 closure days. There were 109 closure days in 2015, certainly impacted the number of available fishing days south of Guam.

## 1.2.18 Administrative and Regulatory Actions

This summary describes management actions NMFS has taken for CNMI fisheries since the April 2018 Joint FEP Plan Team meeting.

June 14, 2018. Final rule. 5-Year Extension of Moratorium on Harvest of Gold Corals. This final rule extends the region-wide moratorium on the harvest of gold corals in the U.S. Pacific Islands through June 30, 2023. NOAA Fisheries intends this final rule to prevent overfishing and to stimulate research on gold corals.

February 8, 2019. Final rule. Reclassifying Management Unit Species to Ecosystem Component Species. This final rule reclassifies certain management unit species in the Pacific Islands as ecosystem component species. The rule also updates the scientific and local names of certain species. The intent of this final rule is to prioritize conservation and management efforts and to improve efficiency of fishery management in the region. This rule is effective March 11, 2019.

## 1.2.19 References

- Cook, R., 2014. Report on the Review of the Biomass Augmented Catch-MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report submitted to the Center for Independent Experts.
- Fricke, H. and D. Meischner. 1985. Depth limits of Bermudan scleractinian corals: a submersible survey. *Marine Biology*, 88(2), pp.175-187.

- Green, A.L. 1997. An assessment of the status of the coral reef resources, and their patterns of use, in the U.S. Pacific Islands. Report submitted to the Western Pacific Regional Fishery Management Council. Honolulu, HI 96813. 278 p.
- Haddon, M. 2014. Center for Independent Experts (CIE) Peer Review of the Biomass Augmented Catch-MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report submitted to the Center for Independent Experts.
- Hart, S.R., Coetzee, M., Workman, R.K., Blusztajn, J., Johnson, K.T.M., Sinton, J.M., Steinberger, B., and J.W. Hawkins. 2004. Genesis of the Western Samoa seamount province: age, geochemical fingerprint and tectonics. *Earth and Planetary Science Letters*, 227(1), pp. 37-56.
- Hensley, R.A. and T.S. Sherwood. 1993. An overview of Guam's inshore fisheries. *Marine Fisheries Review*, 55(2), pp. 129-138.
- Humphreys, R. and R. Moffitt. 1999. Unit 17-Western Pacific Bottomfish and Armorhead Fisheries. DOC, NOAA, NMFS Our Living Oceans--Report on the Status of US Living Marine Resources, pp. 189-192.
- Jones, C., 2014. Biomass Augmented Catch--MSY Model for Pacific Island Coral Reef Ecosystem Resources. Report submitted to the Center for Independent Experts.
- Kahng, S.E. and J.E. Maragos. 2006. The deepest, zooxanthellate scleractinian corals in the world?. *Coral Reefs*, 25(2), pp. 254-254.
- Konter, J.G. and M.G. Jackson. 2012. Large volumes of rejuvenated volcanism in Samoa: Evidence supporting a tectonic influence on late-stage volcanism. *Geochemistry, Geophysics, Geosystems*, 13(6).
- Lang, J.C., 1974. Biological Zonation at the Base of a Reef: Observations from the submersible Nekton Gamma have led to surprising revelations about the deep fore-reef and island slope at Discovery Bay, Jamaica. *American Scientist*, 62(3), pp. 272-281.
- Martell, S. and R. Froese. 2013. A simple method for estimating MSY from catch and resilience. *Fish and Fisheries*, 14(4), pp. 504-514.
- Meyer, R. and R.B. Millar. 1999. Bayesian stock assessment using a state-space implementation of the delay difference model. *Canadian Journal of Fisheries and Aquatic Sciences*, 56(1), pp. 37-52.
- Myers, R.F., 1997. Assessment of coral reef resources of Guam with emphasis on waters of federal jurisdiction. Report prepared for the Western Pacific Regional Fishery Management Council. Honolulu, HI 96813.
- Moffitt, R.B., Brodziak, J., and T. Flores. 2007. Status of the Bottomfish Resources of American Samoa, Guam, and Commonwealth of the Northern Mariana Islands, 2005. NOAA Pacific Islands Fisheries Science Center Administrative Report H-07-04.

- Nadon, M.O., Ault, J.S., Williams, I.D., Smith, S.G. and G.T. DiNardo. 2015. Length-based assessment of coral reef fish populations in the main and northwestern Hawaiian Islands. *PloS one*, 10(8), p.e0133960.
- Neall, V.E. and S.A. Trewick. 2008. The age and origin of the Pacific islands: a geological overview. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1508), pp. 3293-3308.
- NMFS. 2015. Specification of Annual Catch Limits and Accountability Measures for Pacific Island Coral Reef Ecosystem Fisheries in Fishing Years 2015 through 2018. Honolulu, HI 96813. RIN 0648-XD558. 228 pp.
- Oram, R., Flores Jr., T., Tibbatts, B. Gutierrez, J. Gesner, J.P., Wusstig, S., Regis., A., Hamm, D., Quach, M., and P Tao. 2011a. Guam Boat-Based Creel Survey Documentation. NOAA, National Marine Fishery Service, Pacific Island Fishery Science Center, Administrative Report.
- Oram, R., Flores Jr., T., Tibbatts, B. Gutierrez, J. Gesner, J.P., Wusstig, S., Regis., A., Hamm, D., Quach, M., and P. Tao. 2011b. Guam Shore-Based Creel Survey Documentation. NOAA, National Marine Fishery Service, Pacific Island Fishery Science Center, Administrative Report.
- Restrepo, V.R., Thompson, G.G., Mace, P.M., Gabriel, W.L., Low, L.L., MacCall, A.D., Methot, R.D., Powers, J.E., Taylor, B.L., Wade, P.R., and J.F. Witzig. 1998. Technical guidance on the use of precautionary approaches to implementing National Standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. NOAA Technical Memorandum NMFS-F/SPO-31, pp. 1-54.
- Sabater, M. and P. Kleiber. 2014. Augmented catch-MSY approach to fishery management in coral-associated fisheries. *Interrelationships between Corals and Fisheries*. CRC Press, Boca Raton, FL, pp. 199-218.
- Williams, I. 2010. U.S. Pacific Reef Fish Estimates Based on Visual Survey Data. NOAA Pacific Islands Fisheries Science Center Internal Report IR-10-024.
- WPRFMC. 2011. Omnibus Amendment for the Western Pacific Region to Establish a Process for Specifying Annual Catch Limits and Accountability Measures. Western Pacific Regional Fishery Management Council. Honolulu, HI 96813.
- Yau, A., Nadon, M., Richards, B., Brodziak, J., and E. Fletcher. 2016. Stock assessment updates of the Bottomfish Management Unit species of American Samoa, the Commonwealth of the Northern Mariana Islands, and Guam in 2015 using data through 2013. U.S. Dept. of Commerce, NOAA Technical Memorandum, NMFS-PIFSC-51, 54 p.

This page was intentionally left blank.



## 2 ECOSYSTEM CONSIDERATIONS

### 2.1 CORAL REEF ECOSYSTEM PARAMETERS

#### 2.1.1 Regional Reef Fish Biomass

**Description:** ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data between 2010 and 2018.

**Rationale:** Reef fish biomass has been widely used as an indicator of relative ecosystem status, and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime.

**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** American Samoa, Guam, CNMI, MHI, NWHI, and PRIAs

**Spatial Scale:** Regional

**Data Source:** Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods are described in detail elsewhere ([http://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_15-07.pdf](http://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_15-07.pdf)), but in brief involve teams of divers conducting stationary point count cylinder (SPC) surveys within a target domain of <30 meter hard-bottom habitat at each island, stratified by depth zone and, for larger islands, by section of coastline. For consistency among islands, only data from forereef habitats are used. At each SPC, divers record the number, size, and species of all fishes within or passing through paired 15 meter-diameter cylinders over the course of a standard count procedure.

Fish sizes and abundance are converted to biomass using standard length-to-weight conversion parameters, taken largely from FishBase (<http://www.fishbase.org>), and converted to biomass per unit area by dividing by the area sampled per survey. Site-level data were pooled into island-scale values by first calculating mean and variance within strata, and then calculating weighted island-scale mean and variance using the formulas given in Smith et al. (2011), with strata weighted by their respective sizes.





Figure 1. Mean fish biomass (g/m<sup>2</sup> ± standard error) of CREMUS from 2010-2018 by latitude; figure continued on next page



### 2.1.2 CNMI Reef Fish Biomass

**Description:** ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data between 2010 and 2018. Anatahan, Guguan, and Sarigan have been grouped.

**Rationale:** Reef fish biomass has been widely used as an indicator of relative ecosystem status, and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime.

**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1).





Figure 2. Mean fish biomass ( $\text{g/m}^2 \pm$  standard error) of CREMUS from 2010-2018 by latitude with mean estimates overlaid in red; figure continued on next page



### **2.1.3 CNMI Archipelagic Mean Fish Size**

**Description:** ‘Mean fish size’ is mean size of reef fishes > 10 cm TL derived from visual survey data between 2010 and 2018. Anatahan, Guguan, and Sarigan have been grouped.

**Rationale:** Mean size is important as it is widely used as an indicator of fishing pressure. A fishery can sometimes preferentially target large individuals, and can also the number of fishes reaching older (and larger) size classes. Large fishes contribute disproportionately to community fecundity and can have important ecological roles; for example, excavating bites by large parrotfishes probably have a longer lasting impact on reef benthos than bites by smaller fishes.

**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Fishes smaller than 10 cm TL are excluded so that the fish assemblage measured more closely reflects fishes that are potentially fished, and so that mean sizes are not overly influenced by variability in space and time of recent recruitment.





Figure 3. Mean fish size (cm, TL ± standard error) of CNMI CREMUS from 2010-2018 by latitude with mean estimates overlaid in red; figure continued on next page





### 2.1.4 CNMI Reef Fish Population Estimates

**Description:** ‘Reef fish population estimates’ are calculated by multiplying mean biomass per unit area by estimated hardbottom area in a consistent habitat across all islands (specifically, the area of hardbottom forereef habitat in < 30 meters of water).

**Rationale:** Reef fish population estimate data have utility in understanding the size of populations from which fishery harvests are extracted.

**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** Data used to generate mean size estimates come from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Those estimates are converted to population estimates by multiplying biomass (g/m<sup>2</sup>) per island by the estimated area of hardbottom habitat < 30 meters deep at the island, which is the survey domain for the monitoring program that biomass data comes from. Measures of estimated habitat area per island are derived from GIS bathymetry and NOAA Coral Reef Ecosystems Program habitat maps.

Many reef fish taxa are present in other habitats than is surveyed by the program, and some taxa likely have the majority of their populations in deeper water. Additionally, fish counts have the potential to be biased by the nature of fish response to divers. Curious fishes, particularly in locations where divers are not perceived as a threat, will tend to be overestimated by visual survey, while skittish fishes will tend to be undercounted. It is also likely that numbers of jacks and sharks in some locations, such as the NWHI are overestimated by visual survey. Nevertheless, the data shown here are consistently gathered across space and time.

**Table 49. Reef fish estimates for CNMI CREMUS in 0-30 m hard bottom habitat**

| Island              | Total area of reef (Ha) | N   | Estimated population biomass (metric tons) in survey domain of < 30 m hardbottom |            |               |               |            |          |
|---------------------|-------------------------|-----|--|------------|---------------|---------------|------------|----------|
|                     |                         |     | Acanthuridae   | Carangidae | Carcharhinids | Holocentridae | Kyphosidae | Labridae |
| Farallon de Pajaros | 138.5                   | 39  | 13.9   | 11.0       | 5.8           | 6.9           | 7.8        | 1.3      |
| Maug                | 313.9                   | 108 | 26.2   | 5.1        | 7.3           | 5.6           | 10.7       | 3.2      |
| Asuncion            | 248.6                   | 60  | 29.7   | 10.6       | 8.9           | 8.2           | 9.7        | 2.2      |
| Agrihan             | 850.6                   | 39  | 115.8  | 101.0      | 182.3         | 14.1          | 16.6       | 10.6     |

| Pagan               | 1,512.9                        | 112        | 159.4  | 41.0              | 43.2                 | 18.4                 | 13.8              | 15.2             |
|---------------------|--------------------------------|------------|--|-------------------|----------------------|----------------------|-------------------|------------------|
| AGS                 | 743.9                          | 84         | 68.8   | 58.2              | 51.5                 | 18.8                 | 13.4              | 7.1              |
| Saipan              | 3,539.0                        | 115        | 107.6  | 0.1               | 6.8                  | 11.8                 | 0.4               | 20.8             |
| Tinian              | 1,414.2                        | 62         | 44.3   | 4.4               | -                    | 5.3                  | 3.5               | 8.4              |
| Aguijan             | 405.6                          | 40         | 18.9   | 0.1               | 4.8                  | 2.0                  | 0.2               | 2.0              |
| Rota                | 1,331.4                        | 80         | 39.0   | 1.4               | 5.7                  | 3.1                  | 1.7               | 8.4              |
| <b>TOTAL</b>        | <b>10,498.5</b>                | <b>739</b> | <b>600.9</b>   | <b>205.2</b>      | <b>257.8</b>         | <b>91.6</b>          | <b>74.2</b>       | <b>77.4</b>      |
| <b>Island</b>       | <b>Total Area of reef (Ha)</b> | <b>N</b>   | <b>Lethrinidae</b>   | <b>Lutjanidae</b> | <b>Mullidae</b>      | <b>Scaridae</b>      | <b>Serranidae</b> | <b>Siganidae</b> |
| Farallon de Pajaros | 138.5                          | 39         | 0.7  | 22.1              | 0.8                  | 3.0                  | 4.3               | -                |
| Maug                | 313.9                          | 108        | 2.8  | 27.7              | 1.6                  | 16.5                 | 6.8               | -                |
| Asuncion            | 248.6                          | 60         | 1.4  | 31.1              | 2.0                  | 13.0                 | 8.3               | -                |
| Agrihan             | 850.6                          | 39         | 8.8  | 113.0             | 5.2                  | 72.4                 | 36.6              | -                |
| Pagan               | 1,512.9                        | 112        | 23.9   | 165.7             | 8.9                  | 94.3                 | 35.0              | -                |
| AGS                 | 743.9                          | 84         | 10.4   | 56.6              | 3.3                  | 40.8                 | 27.1              | -                |
| Saipan              | 3,539.0                        | 115        | 9.8  | 49.9              | 6.2                  | 72.2                 | 13.7              | 0.9              |
| Tinian              | 1,414.2                        | 62         | 16.0   | 16.4              | 3.1                  | 28.0                 | 8.8               | 0.3              |
| Aguijan             | 405.6                          | 40         | 2.0  | 9.9               | 1.0                  | 8.4                  | 2.8               | -                |
| Rota                | 1,331.4                        | 80         | 14.2   | 10.8              | 3.4                  | 47.8                 | 4.9               | 0.2              |
| <b>TOTAL</b>        | <b>10,498.5</b>                | <b>739</b> | <b>89.3</b>  | <b>478.1</b>      | <b>34.2</b>          | <b>381.0</b>         | <b>137.9</b>      | <b>1.4</b>       |
| <b>Island</b>       | <b>Total area of reef (Ha)</b> | <b>N</b>   | <b>Estimated population biomass (metric tons) in survey domain of &lt; 30 m hardbottom</b> |                   |                      |                      |                   |                  |
|                     |                                |            | <b>Acanthuridae</b>  | <b>Carangidae</b> | <b>Carcharhinids</b> | <b>Holocentridae</b> | <b>Kyphosidae</b> | <b>Labridae</b>  |
| Farallon de Pajaros | 138.5                          | 23         | 15.0   | 8.8               | 8.7                  | 7.6                  | 9.4               | 1.1              |
| Maug                | 313.9                          | 70         | 26.4   | 5.4               | 6.8                  | 6.3                  | 7.5               | 2.9              |
| Asuncion            | 248.6                          | 41         | 28.0   | 7.7               | 12.0                 | 7.9                  | 9.5               | 2.3              |
| Agrihan             | 850.6                          | 20         | 131.9  | 36.0              | 116.4                | 11.9                 | 5.8               | 8.5              |
| Pagan               | 1,512.9                        | 72         | 156.3  | 34.2              | 39.6                 | 19.0                 | 13.0              | 15.1             |

| AGS                 | 743.9                          | 57         | 85.0               | 73.6              | 48.0            | 19.7            | 11.0              | 8.5              |
|---------------------|--------------------------------|------------|--------------------|-------------------|-----------------|-----------------|-------------------|------------------|
| Saipan              | 4,846.6                        | 78         | 168.5              | 0.3               | -               | 17.3            | 0.7               | 31.2             |
| Tinian              | 1,414.2                        | 38         | 51.4               | 5.9               | -               | 4.4             | 4.2               | 10.5             |
| Aguijan             | 405.6                          | 23         | 22.4               | -                 | 7.2             | 1.1             | 0.3               | 2.2              |
| Rota                | 1,331.4                        | 52         | 45.4               | 2.1               | -               | 2.7             | 2.5               | 8.1              |
| <b>TOTAL</b>        | <b>11,806.1</b>                | <b>474</b> | <b>689.4</b>       | <b>164.1</b>      | <b>186.0</b>    | <b>95.5</b>     | <b>63.5</b>       | <b>88.8</b>      |
| <b>Island</b>       | <b>Total Area of reef (Ha)</b> | <b>N</b>   | <b>Lethrinidae</b> | <b>Lutjanidae</b> | <b>Mullidae</b> | <b>Scaridae</b> | <b>Serranidae</b> | <b>Siganidae</b> |
| Farallon de Pajaros | 138.5                          | 23         | 0.7                | 23.4              | 0.6             | 3.7             | 3.8               | -                |
| Maug                | 313.9                          | 70         | 2.8                | 27.0              | 1.6             | 15.4            | 6.5               | -                |
| Asuncion            | 248.6                          | 41         | 1.2                | 27.5              | 1.8             | 13.0            | 7.0               | -                |
| Agrihan             | 850.6                          | 20         | 4.7                | 142.1             | 5.4             | 80.1            | 45.6              | -                |
| Pagan               | 1,512.9                        | 72         | 22.9               | 149.6             | 5.2             | 85.3            | 33.3              | -                |
| AGS                 | 743.9                          | 57         | 14.3               | 63.5              | 3.2             | 48.6            | 28.7              | -                |
| Saipan              | 4,846.6                        | 78         | 14.9               | 94.4              | 8.4             | 93.1            | 17.8              | 1.8              |
| Tinian              | 1,414.2                        | 38         | 19.9               | 11.7              | 2.6             | 27.1            | 8.7               | 0.1              |
| Aguijan             | 405.6                          | 23         | 2.9                | 10.7              | 1.2             | 9.4             | 2.6               | -                |
| Rota                | 1,331.4                        | 52         | 16.9               | 11.9              | 2.2             | 56.6            | 5.6               | 0.3              |
| <b>TOTAL</b>        | <b>11,806.1</b>                | <b>474</b> | <b>102.1</b>       | <b>508.8</b>      | <b>30.5</b>     | <b>405.3</b>    | <b>140.4</b>      | <b>2.3</b>       |

Notes: (1) No *Bolbometopon muricatum* were observed during these surveys in CNMI.

(2) *Cheilinus undulatus* were recorded at Tinian (0.7 t).

(3) N is the number of site surveyed.

(4) 'AGS' is a combined value for Alamagan, Guguan, and Sarigan.

### 2.1.5 Guam Reef Fish Biomass

**Description:** ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data (details of survey program below) between 2009 and 2015.

**Rationale:** Reef fish biomass has been widely used as an indicator of relative ecosystem status, and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime.


**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1).



**Figure 4. Mean fish biomass ( $g/m^2 \pm$  standard error) of Guam CREMUS from the years 2010-2018 with mean estimates overlaid in red**

### **2.1.6 Guam Archipelagic Mean Size**

**Description:** 'Mean fish size' is mean size of reef fishes > 10 cm TL derived from visual survey data between 2010 and 2018.

**Rationale:** Mean size is important as it is widely used as an indicator of fishing pressure. A fishery can sometimes preferentially target large individuals, and can also the number of fishes reaching older (and larger) size classes. Large fishes contribute disproportionately to community fecundity and can have important ecological roles; for example, excavating bites by large parrotfishes probably have a longer lasting impact on reef benthos than bites by smaller fishes.


**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Fishes smaller than 10 cm TL are excluded so that the fish assemblage measured more closely reflects fishes that are potentially fished, and so that mean sizes are not overly influenced by variability in space and time of recent recruitment.



**Figure 5. Mean fish size (cm, TL ± standard error) of Guam CREMUS from 2010-2018 with mean estimates overlaid in red**



### 2.1.7 Guam Reef Fish Population Estimates

**Description:** ‘Reef fish population estimates’ are calculated by multiplying mean biomass per unit area by estimated hardbottom area in a consistent habitat across all islands (specifically, the area of hardbottom forereef habitat in < 30 meters of water).

**Rationale:** Reef fish population estimates data have utility in understanding the size of populations from which fishery harvests are extracted.

**Data Category:** Fishery independent

**Timeframe:** Triennial

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** Data used to generate mean size estimates come from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program ([http://www.pifsc.noaa.gov/cred/pacific\\_ramp.php](http://www.pifsc.noaa.gov/cred/pacific_ramp.php)). Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Those estimates are converted to population estimates by multiplying biomass (g/m<sup>2</sup>) per island by the estimated area of hardbottom habitat <30 meters deep at the island, which is the survey domain for the monitoring program that biomass data comes from. Measures of estimated habitat area per island are derived from GIS bathymetry and NOAA CREP habitat maps.

Many reef fish taxa are present in other habitats than is surveyed by the program, and some taxa likely have the majority of their populations in deeper water. Additionally, fish counts have the potential to be biased by the nature of fish response to divers. Curious fishes, particularly in locations where divers are not perceived as a threat, will tend to be overestimated by visual survey, while skittish fishes will tend to be undercounted. It is also likely that numbers of jacks and sharks in some locations, such as the NWHI are overestimated by visual survey.

Nevertheless, the data shown here are consistently gathered across space and time. Nevertheless, in spite of these issues, the data shown here are consistently gathered across space and time.

**Table 50. Reef fish population estimates for Guam CREMUS**

| Island | Total area of reef (Ha) | N   | Estimated population biomass (metric tons) in survey domain of < 30 m hardbottom |            |               |               |            |           |
|--------|-------------------------|-----|--|------------|---------------|---------------|------------|-----------|
|        |                         |     | Acanthuridae   | Carangidae | Carcharhinids | Holocentridae | Kyphosidae | Labridae  |
| Guam   | 5,177.4                 | 304 | 186.1  | 142.6      | 39.8          | 21.8          | 4.2        | 34.6      |
|        |                         |     | Lethrinidae  | Lutjanidae | Mullidae      | Scaridae      | Serranidae | Siganidae |
| Guam   | 5,177.4                 | 304 | 27.7   | 46.2       | 11.0          | 181.3         | 31.9       | 2.10      |

Notes:

(1) N is the number of sites surveyed.

(2) No *Bolbometopon muricatum* were observed during these surveys in Guam.

(3) *Cheilinus undulatus* were recorded in Guam (43.2 t).

## 2.2 LIFE HISTORY AND LENGTH-DERIVED PARAMETERS

The annual SAFE report will serve as the repository of available life history information for the Western Pacific region. Life history data particularly age, growth, reproduction and mortality information inform the stock assessment on fish productivity and population dynamics. Some assessments particularly for data poor stocks like coral reefs utilize information from other areas that introduces errors and uncertainties in the population estimates. An archipelago specific life history parameter ensures accuracy in the input parameters used in the assessment.

The NMFS Biosampling Program allows for significant collection of life history samples like otoliths and gonads from priority species in the bottomfish and coral reef fisheries. A significant number of samples are also collected during research cruises. These life history samples, once processed and data extracted, will contribute to the body of scientific information for the two data-poor fisheries in the region. The life history information available from the region will be monitored by the Fishery Ecosystem Plan Team and will be tracked through this section of the report.

This section will be divided into two fisheries: 1) coral reef; and 2) bottomfish. Within each fishery, the available life history information will be described under the age, growth, and reproductive maturity section. The section labelled fish length-derived parameters summarizes available information derived from sampling the fish catch or the market. Monitoring length information provides insight on the state of the fish stock where the change in length can be used as an indicator of population level mortality. Length-weight conversion coefficients provide area-specific values to convert length from fishery dependent and fishery independent data collection to weight or biomass.

### 2.2.1 CNMI Coral Reef Ecosystem – Reef Fish Life History

#### 2.2.1.1 Age & Growth and Reproductive Maturity

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely-cut, thin sections of sagittal otoliths. Validated age determination is based on an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally-based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. The relation between age and fish length is evaluated by fitting this data to a von Bertalanffy growth function based on statistical analyses. The resulting von Bertalanffy growth function predicts the pattern of growth over time for that particular species. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ) which together characterize the shape of the length-at-age growth relationship.

Length-at-reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved then subsequently cut into five micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard

cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of mature samples for a given length interval are assembled for each sex and these data are fitted to a three- or four-parameter logistic function to determine the best fit of these data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region) - such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes - standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a three or four-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age & growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted logistic function provides a direct estimate of the age at which 50% of fish of a particular species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

### **Parameter definitions:**

**$T_{max}$  (maximum age)** – The maximum observed age revealed from an otolith-based age determination study.  $T_{max}$  values can be derived from ages determined by annuli counts of sagittal otolith sections and/or bomb radiocarbon ( $^{14}\text{C}$ ) analysis of otolith core material. Units are years.

**$L_{\infty}$  (asymptotic length)** – One of three coefficients of the von Bertalanffy growth function (VBGF) that measures the mean maximum length at which the growth curve plateaus and no longer increases in length with increasing age. This coefficient reflects the mean maximum length and not the observed maximum length. Units are centimeters.

**$k$  (growth coefficient)** – One of three coefficients of the VBGF that measures the shape and steepness by which the initial portion of the growth function approaches its mean maximum length ( $L_{\infty}$ ).

**$t_0$  (hypothetical age at length zero)** – One of three coefficients of the VBGF whose measure is highly influenced by the other two VBGF coefficients ( $k$  and  $L_{\infty}$ ) and typically assumes a negative value when specimens representing early growth phases (0+ to 1+ ages) are not available for age determination. Units are years.

**$M$  (natural mortality)** – This is a measure of mortality rate for a fish stock not under the influence of fishing pressure and is considered to be directly related to stock productivity (i.e., high  $M$  indicates high productivity and low  $M$  indicates low stock productivity).  $M$  can be

derived through use of various equations that link  $M$  to  $T_{max}$  and two VBGF coefficients ( $k$  and  $L_{\infty}$ ) or by calculating the value of the slope from a regression fit to a declining catch curve (regression of the natural logarithm of abundance versus age class) derived from fishing an unfished or lightly fished population.

**$A_{50}$  (age at 50% maturity)** – Age at which 50% of the sampled stock under study has attained reproductive maturity. This parameter is best determined based on studies that concurrently determine both age (otolith-based age data) and reproductive maturity status (logistic function fitted to percent mature by age class with maturity determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A_{50}$  is to use an existing  $L_{50}$  estimate to find the corresponding age ( $A_{50}$ ) from an existing VBGF curve. Units are years.

**$A\Delta_{50}$  (age of sex switching)** – Age at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal. This parameter is best determined based on studies that concurrently determines both age (otolith-based age data) and reproductive sex reversal status (logistic function fitted to percent sex reversal by age class with sex reversal determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A\Delta_{50}$  is to use an existing  $L\Delta_{50}$  estimate to find the corresponding age ( $A\Delta_{50}$ ) from the VBGF curve. Units are years.

**$L_{50}$  (length at which 50% of a fish species are capable of spawning)** – Length (usually in terms of fork length) at which 50% of the females of a sampled stock under study has attained reproductive maturity; this is the length associated with  $A_{50}$  estimates. This parameter is derived using a logistic function to fit the percent mature data by length class with maturity status best determined via microscopic analyses of gonad histology preparations).  $L_{50}$  information is typically more available than  $A_{50}$  since  $L_{50}$  estimates do not require knowledge of age & growth. Units are centimeters.

**$L\Delta_{50}$  (length of sex switching)** – Length (usually in terms of fork length) at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal; this is the length associated with  $A\Delta_{50}$  estimates. This parameter is derived using a logistic function to fit the percent sex reversal data by length class with sex reversal status best determined via microscopic analyses of gonad histology preparations).  $L\Delta_{50}$  information is typically more available than  $A\Delta_{50}$  since  $L\Delta_{50}$  estimates do not require knowledge of age & growth. Units are centimeters.

**Rationale:** These nine life-history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. Currently, the assessment of coral reef fish resources in the CNMI is data-limited. Knowledge of these life-history parameters support current efforts to characterize the resilience of these resources, provide important biological inputs for future stock assessment efforts, and enhance our understanding of the species' likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** CNMI

**Spatial Scale:** Archipelagic

**Data Source:** Sources of data are directly derived from research cruises sampling and market samples collected by the CNMI contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 51 for specific details on data sources by species.

**Table 51. Available age, growth, and reproductive maturity information for coral reef species targeted for otolith and gonad sampling in CNMI**

| Species                             | Age, growth, and reproductive maturity parameters |                   |                  |                                    |                   |                               |                |                                 |                | Reference                          |
|-------------------------------------|---|-------------------|------------------|------------------------------------|-------------------|-------------------------------|----------------|---------------------------------|----------------|------------------------------------|
|                                     | $T_{max}$   | $L_{\infty}$      | $k$              | $t_0$                              | $M$               | $A_{50}$                      | $A\Delta_{50}$ | $L_{50}$                        | $L\Delta_{50}$ |                                    |
| <i>Calotomus carolinus</i>          |   |                   |                  |                                    |                   |                               |                |                                 |                |                                    |
| <i>Chlorurus spilurus</i>           |   |                   |                  |                                    |                   |                               |                |                                 |                |                                    |
| <i>Lethrinus atkinsoni</i>          |   |                   |                  |                                    |                   |                               |                |                                 |                |                                    |
| <i>Lethrinus obsoletus</i>          | 13 <sup>d</sup>                                   | 25.1 <sup>d</sup> | 0.6 <sup>d</sup> | 3.0 (L <sub>0</sub> ) <sup>d</sup> | 0.32 <sup>d</sup> | 3.8 (f), 2.8 (m) <sup>d</sup> | <sup>a</sup>   | 22.9 (f), 19.9 (m) <sup>d</sup> |                | <sup>d</sup> Taylor et. al. (2017) |
| <i>Mulloidichthys flavolineatus</i> | X <sup>a</sup>                                    | X <sup>a</sup>    | X <sup>a</sup>   | X <sup>a</sup>                     | X <sup>a</sup>    | X <sup>a</sup>                |                | X <sup>a</sup>                  |                | Reed et al., in prep.              |
| <i>Naso unicornis</i>               |   |                   |                  |                                    |                   |                               | NA             | 238 <sup>b</sup>                | NA             |                                    |
| <i>Parupeneus barberinus</i>        | X <sup>a</sup>                                    | X <sup>a</sup>    | X <sup>a</sup>   | X <sup>a</sup>                     | X <sup>a</sup>    | X <sup>a</sup>                | NA             | X <sup>a</sup>                  |                | Reed et al., in prep.              |
| <i>Sargocentron tere</i>            |   |                   |                  |                                    |                   |                               | NA             |                                 | NA             |                                    |
| <i>Siganus argenteus</i>            | 7 <sup>d</sup>                                    | 274 <sup>d</sup>  | 0.9 <sup>d</sup> | -0.3 <sup>d</sup>                  | 0.56 <sup>d</sup> | 1.3 <sup>d</sup>              | NA             | 218 <sup>d</sup>                | NA             | <sup>d</sup> Taylor et. al. (2016) |

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (including in press).

Parameter estimates are for females unless otherwise noted (F=females, M=males). Parameters  $T_{max}$ ,  $t_0$ ,  $A_{50}$ , and  $A\Delta_{50}$  are in units of years;  $L_{\infty}$ ,  $L_{50}$ , and  $L\Delta_{50}$  are in units of mm fork length (FL);  $k$  in units of year<sup>-1</sup>; X=parameter estimate too preliminary or Y=published age and growth parameter estimates based on DGI numerical integration technique and likely to be inaccurate;

NA=not applicable. Superscript letters indicate status of parameter estimate (see footnotes below table). Published or in press publications (<sup>d</sup>) are denoted in “Reference” column.

### 2.2.1.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Biosampling Program started in 2009. This program has two components: first is the Field/Market Sampling Program, and the second is the Life History Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program are:

- Broad scale look at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification;
- Develop accurate local length-weight curves.

In CNMI, the Biosampling is focused on the commercial coral reef spear fishery with occasional sampling of the bottomfish fishery occurring locally and less frequently at the northern islands. Sampling is conducted in partnership with the fish vendors. The Market Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information. Specific for CNMI, the program collects Daily Vendor Logs for reef fish that includes basic catch and effort information.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** NMFS Biosampling Program

**Parameter definitions:**

$L_{max}$  – **maximum fish length** is the longest fish per species recorded in the Biosampling Program from the commercial spear fishery. This value is derived from measuring the fork length of individual samples for species occurring in the spear fishery.

$L_{bar}$  – **mean length** is the average value of all lengths recorded from the commercial spear fishery. This can be influenced by gear selectivity since the commercial spear fishery has a typical size target based on customer demand. This can also be influenced by size regulations.

$n$  – **sample size** is the total number of samples accumulated for each species recorded in the commercial spear fishery.

$N_{L-W}$  – **sample size for L-W regression** is the number of samples used to generate the  $a$  &  $b$  coefficients.


*a* & *b* – **length-weight coefficients** are the coefficients derived from the regression line fitted to all length- and weight-measured per species in the commercial spear fishery. These values are used to convert length information to weight. Values are influenced by the life history characteristics of the species, geographic location, population status, and nature of the fisheries from which the species are harvested.

**Rationale:** Length-derived information is being used as an indicator of population status particularly for data poor stocks like coral reef fish. Average length ( $L_{bar}$ ) was used as a principal stock assessment indicator variable for exploited reef fish population (Nadon et al., 2015). Average length was also shown to be correlated with population size (Kerr and Dickle, 2001). Maximum length ( $L_{max}$ ), typically coupled with maximum age, is typically used as a proxy for fish longevity which has implications on the productivity and susceptibility of a species to fishing pressure. The length-weight coefficients (*a* & *b* values) are used to convert length to weight for fishery-dependent and fishery-independent data collection where length are typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the CNMI coral reef and bottomfish fisheries.


**Table 52. Available length-derived information for various coral reef species in CNMI**

| Species                             | Length-derived parameters |           |          |            |          |          | Reference |
|-------------------------------------|---------------------------|-----------|----------|------------|----------|----------|-----------|
|                                     | $L_{max}$                 | $L_{bar}$ | <i>N</i> | <i>L-W</i> | <i>a</i> | <i>b</i> |           |
| <i>Naso lituratus</i>               | 30.1                      | 20.26     | 17,478   | 3,813      | 0.0167   | 3.1022   |           |
| <i>Acanthurus lineatus</i>          | 23.5                      | 18.33     | 15,772   | 4,901      | 0.0383   | 2.8718   |           |
| <i>Siganus argenteus</i>            | 34.1                      | 20.82     | 11,867   | 3,662      | 0.0133   | 3.1007   |           |
| <i>Mulloidichthys flavolineatus</i> | 31.4                      | 18.08     | 9,596    | 2,357      | 0.0137   | 3.0547   |           |
| <i>Naso unicornis</i>               | 53.6                      | 29.62     | 8,323    | 4,349      | 0.0266   | 2.9115   |           |
| <i>Siganus spinus</i>               | 25.6                      | 16.64     | 7,685    | 1,078      | 0.0118   | 3.1459   |           |
| <i>Parupeneus barberinus</i>        | 37.3                      | 21.73     | 7,597    | 2,706      | 0.0175   | 3.0119   |           |
| <i>Selar crumenophthalmus</i>       | 26.5                      | 19.08     | 4922     | 2654       | 0.0051   | 3.3958   |           |
| <i>Scarus ghobban</i>               | 38.1                      | 24.07     | 4,964    | 1,502      | 0.0124   | 3.1271   |           |
| <i>Lethrinus atkinsoni</i>          | 35.1                      | 21.06     | 4,306    | 2,095      | 0.0163   | 3.0971   |           |
| <i>Lethrinus obsoletus</i>          | 29.0                      | 21.10     | 3,673    | 1,472      | 0.0171   | 3.0313   |           |
| <i>Mulloidichthys vanicolensis</i>  | 28.0                      | 18.94     | 3,233    | 701        | 0.0103   | 3.1948   |           |
| <i>Scarus rubroviolaceus</i>        | 52.6                      | 34.49     | 3,141    | 1,791      | 0.0087   | 3.2447   |           |
| <i>Chlorurus sordidus</i>           | 30.8                      | 22.33     | 3,346    | 956        | 0.0173   | 3.0795   |           |
| <i>Siganus punctatus</i>            | 34.8                      | 20.82     | 2,798    | 833        | 0.0129   | 3.1911   |           |
| <i>Sargocentron spiniferum</i>      | 34.6                      | 20.31     | 2,589    | 684        | 0.0245   | 2.9780   |           |
| <i>Myripristis murdjan</i>          | 22.3                      | 16.84     | 2,488    | 823        | 0.1699   | 2.3426   |           |
| <i>Scarus psittacus</i>             | 28.9                      | 21.24     | 2,466    | 771        | 0.0212   | 2.9928   |           |
| <i>Acanthurus nigricauda</i>        | 26.3                      | 20.07     | 2,354    | 799        | 0.0217   | 3.0583   |           |
| <i>Cheilinus trilobatus</i>         | 35.2                      | 24.06     | 2,223    | 1,196      | 0.0470   | 2.7156   |           |
| <i>Hipposcarus longiceps</i>        | 52.0                      | 29.10     | 2,194    | 615        | 0.0149   | 3.0624   |           |
| <i>Panulirus penicillatus</i>       | 17.0                      | 9.05      | 2,043    | 1,119      | 1.4849   | 2.6925   |           |
| <i>Leptoscarus vaigiensis</i>       | 35.2                      | 26.31     | 1,982    | 807        | 0.0234   | 2.8648   |           |
| <i>Calotomus carolinus</i>          | 31.0                      | 24.21     | 1,734    | 662        | 0.0156   | 3.1012   |           |









| Species                     | Length-derived parameters |           |       |       |        |        | Reference |
|-----------------------------|---------------------------|-----------|-------|-------|--------|--------|-----------|
|                             | $L_{max}$                 | $L_{bar}$ | $N$   | $L-W$ | $a$    | $b$    |           |
| <i>Myripristis violacea</i> | 20.6                      | 15.54     | 1,796 | 514   | 0.1361 | 2.4356 |           |




**Figure 6. Average length over time of representative CNMI CREMUS from the Biosampling Program; figure continues for various species onto the next two pages**









## 2.2.2 CNMI Bottomfish Ecosystem – Bottomfish Life History

### 2.2.2.1 Age & Growth and Reproductive Maturity

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely-cut, thin sections of sagittal otoliths. Validated age determination is based on an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally-based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. The relation between age and fish length is evaluated by fitting this data to a von Bertalanffy growth function based on statistical analyses. The resulting von Bertalanffy growth function predicts the pattern of growth over time for that particular species. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ) which together characterize the shape of the length-at-age growth relationship.

Length-at-reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved then subsequently cut into five micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of mature samples for a given length interval are assembled for each sex and these data are fitted to a three- or four-parameter logistic function to determine the best fit of these data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region), such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes, standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a three or four-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age & growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted logistic function provides a direct estimate of the age at which 50% of fish of a particular species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** Sources of data are directly derived from research cruises sampling and market samples collected by the CNMI contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 53 for specific details on data sources by species.

**Parameter definitions:**

**$T_{max}$  (maximum age)** – The maximum observed age revealed from an otolith-based age determination study.  $T_{max}$  values can be derived from ages determined by annuli counts of sagittal otolith sections and/or bomb radiocarbon ( $^{14}\text{C}$ ) analysis of otolith core material. Units are years.

**$L_{\infty}$  (asymptotic length)** – One of three coefficients of the von Bertalanffy growth function (VBGF) that measures the mean maximum length at which the growth curve plateaus and no longer increases in length with increasing age. This coefficient reflects the mean maximum length and not the observed maximum length. Units are centimeters

**$k$  (growth coefficient)** – One of three coefficients of the VBGF that measures the shape and steepness by which the initial portion of the growth function approaches its mean maximum length ( $L_{\infty}$ ).

**$t_0$  (hypothetical age at length zero)** – One of three coefficients of the VBGF whose measure is highly influenced by the other two VBGF coefficients ( $k$  and  $L_{\infty}$ ) and typically assumes a negative value when specimens representing early growth phases (0+ to 1+ ages) are not available for age determination. Units are years.

**$M$  (natural mortality)** – this is a measure of mortality rate for a fish stock not under the influence of fishing pressure and is considered to be directly related to stock productivity (i.e., high  $M$  indicates high productivity and low  $M$  indicates low stock productivity).  $M$  can be derived through use of various equations that link  $M$  to  $T_{max}$  and two VBGF coefficients ( $k$  and  $L_{\infty}$ ) or by calculating the value of the slope from a regression fit to a declining catch curve (regression of the natural logarithm of abundance versus age class) derived from fishing an unfished or lightly fished population.

**$A_{50}$  (age at 50% maturity)** – Age at which 50% of the sampled stock under study has attained reproductive maturity. This parameter is best determined based on studies that concurrently determine both age (otolith-based age data) and reproductive maturity status (logistic function fitted to percent mature by age class with maturity determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A_{50}$  is to use an existing  $L_{50}$  estimate to find the corresponding age ( $A_{50}$ ) from an existing VBGF curve. Units are years.

**$A\Delta_{50}$  (age of sex switching)** – Age at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal. This parameter is best determined based on studies that concurrently determines both age (otolith-based age data) and reproductive sex reversal status (logistic function fitted to percent sex reversal by age class with sex reversal determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A\Delta_{50}$  is to use an existing  $L\Delta_{50}$  estimate to find the corresponding age ( $A\Delta_{50}$ ) from the VBGF curve. Units are years.

**$L_{50}$  (length at which 50% of a fish species are capable of spawning)** – Length (usually in terms of fork length) at which 50% of the females of a sampled stock under study has attained reproductive maturity; this is the length associated with  $A_{50}$  estimates. This parameter is derived using a logistic function to fit the percent mature data by length class with maturity status best determined via microscopic analyses of gonad histology preparations).  $L_{50}$  information is typically more available than  $A_{50}$  since  $L_{50}$  estimates do not require knowledge of age & growth. Units are in centimeters.

**$L\Delta_{50}$  (length of sex switching)** – Length (usually in terms of fork length) at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex

reversal; this is the length associated with  $A\Delta_{50}$  estimates. This parameter is derived using a logistic function to fit the percent sex reversal data by length class with sex reversal status best determined via microscopic analyses of gonad histology preparations).  $L\Delta_{50}$  information is typically more available than  $A\Delta_{50}$  since  $L\Delta_{50}$  estimates do not require knowledge of age & growth. Units are in centimeters.

**Rationale:** These nine life-history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. Currently, the assessment of coral reef fish resources in CNMI is data-limited. Knowledge of these life-history parameters support current efforts to characterize the resilience of these resources, provide important biological inputs for future stock assessment efforts, and enhance our understanding of the species' likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Table 53. Available age, growth, and reproductive maturity information for bottomfish species targeted for otoliths and gonads sampling in CNMI**

| Species                            | Age, growth, and reproductive maturity parameters |                |                |                |                |          |                |          |                | Reference                   |
|------------------------------------|---|----------------|----------------|----------------|----------------|----------|----------------|----------|----------------|-----------------------------|
|                                    | $T_{max}$   | $L_{\infty}$   | $k$            | $t_0$          | $M$            | $A_{50}$ | $A\Delta_{50}$ | $L_{50}$ | $L\Delta_{50}$ |                             |
| <i>Aphareus rutilans</i>           | Y   | Y              | Y              | Y              |                |          | NA             |          | NA             | Y-Ralston & Williams (1988) |
| <i>Aprion virescens</i>            |   |                |                |                |                |          | NA             |          | NA             |                             |
| <i>Etelis carbunculus</i>          |   |                |                |                |                |          | NA             |          | NA             |                             |
| <i>Etelis coruscans</i>            | Y   | Y              | Y              | Y              |                |          | NA             |          | NA             | Y-Ralston & Williams (1988) |
| <i>Monotaxis grandoculis</i>       |   |                |                |                |                |          |                |          |                |                             |
| <i>Pristipomoides auricilla</i>    | X <sup>c</sup>                                    | X <sup>c</sup> | X <sup>c</sup> | X <sup>c</sup> | X <sup>c</sup> |          | NA             |          | NA             | O'Malley et al. (in review) |
| <i>Pristipomoides filamentosus</i> |   |                |                |                |                |          | NA             |          | NA             |                             |
| <i>Pristipomoides flavipinnis</i>  |   |                |                |                |                |          | NA             |          | NA             |                             |
| <i>Pristipomoides sieboldii</i>    | Y   | Y              | Y              | Y              |                |          | NA             |          | NA             | Y-Ralston & Williams (1988) |
| <i>Pristipomoides zonatus</i>      | X <sup>a</sup>                                    | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> |          | NA             |          | NA             | LHP (in prep)               |
| <i>Variola louti</i>               |   |                |                |                |                |          |                |          |                |                             |

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (including in press).

Parameter estimates are for females unless otherwise noted (F=females, M=males). Parameters  $T_{max}$ ,  $t_0$ ,  $A_{50}$ , and  $A\Delta_{50}$  are in units of years;  $L_{\infty}$ ,  $L_{50}$ , and  $L\Delta_{50}$  are in units of mm fork length (FL);  $k$  in units of year<sup>-1</sup>; X=parameter estimate too preliminary or Y=published age and growth

parameter estimates based on DGI numerical integration technique and likely to be inaccurate; NA=not applicable. Superscript letters indicate status of parameter estimate (see footnotes below table). Published or in press publications (<sup>d</sup>) are denoted in “Reference” column.

#### 2.2.2.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Biosampling Program started in 2009. This program has two components: first is the Field/Market Sampling Program and the second is the Life History Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program are:

- Broad scale look at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification;
- Develop accurate local length-weight curves.

In CNMI, the Biosampling is focused on the commercial coral reef spear fishery with occasional sampling of the bottomfish fishery occurring locally and less frequently at the northern islands. Sampling is conducted in partnership with the fish vendors. The Market Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information. Specific for CNMI, the program collects Daily Vendor Logs for bottomfish that includes basic catch and effort information.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** CNMI

**Spatial Scale:** Island

**Data Source:** NMFS Biosampling Program

**Parameter definitions:**

$L_{max}$  – **maximum fish length** is the longest fish per species recorded in the Biosampling Program from the commercial bottomfish fishery. This value is derived from measuring the fork length of individual samples for species occurring in the spear fishery.

$L_{bar}$  – **mean length** is the average value of all lengths recorded from the commercial spear fishery. This can be influenced by gear selectivity since the commercial bottomfish fishery has a typical size target based on customer demand. This can also be influenced by size regulations.

$n$  – **sample size** is the total number of samples accumulated for each species recorded in the commercial bottomfish fishery.








$N_{L-W}$  – **sample size for L-W regression** is the number of samples used to generate the  $a$  &  $b$  coefficients.

*a & b – length-weight coefficients* are the coefficients derived from the regression line fitted to all length and weight measured per species in the commercial bottomfish fishery. These values are used to convert length information to weight. Values are influenced by the life history characteristics of the species, geographic location, population status, and nature of the fisheries from which the species are harvested from.

**Rationale:** Length-derived information is being used as an indicator of population status particularly for data poor stocks like coral reef fish. Average length ( $L_{bar}$ ) was used as a principal stock assessment indicator variable for exploited reef fish population (Nadon et al., 2015). Average length was also shown to be correlated with population size (Kerr and Dickle, 2001). Maximum length ( $L_{max}$ ), typically coupled with maximum age, is typically used as a proxy for fish longevity which has implications on the productivity and susceptibility of a species to fishing pressure. The length-weight coefficients (*a & b* values) are used to convert length to weight for fishery-dependent and fishery-independent data collection where length are typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the bottomfish fisheries.

**Table 54. Available length-derived information for various bottomfish species in CNMI**

| Species                               | Length-derived parameters |           |       |       |        |        | Reference |
|---------------------------------------|---------------------------|-----------|-------|-------|--------|--------|-----------|
|                                       | $L_{max}$                 | $L_{bar}$ | $N$   | $L-W$ | $a$    | $b$    |           |
| <i>Lethrinus rubrioperculatus</i>     | 38.0                      | 28.01     | 1,353 | 1,021 | 0.0185 | 2.9897 |           |
| <i>Etelis carbunculus</i>             | 53.5                      | 30.18     | 685   | 685   | 0.0150 | 3.0430 |           |
| <i>Pristipomoides auricilla</i>       | 39.5                      | 28.59     | 465   | 465   | 0.0189 | 3.0060 |           |
| <i>Pristipomoides zonatus</i>         | 45.4                      | 32.99     | 371   | 370   | 0.0180 | 3.0411 |           |
| <i>Etelis coruscans</i>               | 96.4                      | 72.50     | 325   | 325   | 0.0716 | 2.6147 |           |
| <i>Lutjanus kasmira</i>               | 32.5                      | 24.84     | 258   | 258   | 0.0087 | 3.2307 |           |
| <i>Pristipomoides flavipinnis</i>     | 51.5                      | 37.05     | 168   | 168   | 0.0133 | 3.0762 |           |
| <i>Pristipomoides argyrogrammicus</i> | 31.6                      | 24.44     | 150   | 150   | 0.0174 | 3.0464 |           |
| <i>Pristipomoides filamentosus</i>    | 58.5                      | 39.97     | 123   | 123   | 0.0773 | 2.5914 |           |
| <i>Caranx lugubris</i>                | 82.5                      | 46.07     | 122   | 122   | 0.0309 | 2.8768 |           |





**Figure 7. Average length over time of CNMI BMUS derived from the Biosampling Program; continued from previous page**

### **2.2.3 Guam Coral Reef Ecosystem – Reef Fish Life History**

#### **2.2.3.1 Age & Growth and Reproductive Maturity**

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely-cut, thin sections of sagittal otoliths. Validated age determination, particularly for long-lived ( $\geq 30$  years) fish, is based on an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally-based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. The relation between age and fish length is evaluated by fitting this data to a von Bertalanffy growth function based on statistical analyses. The resulting von Bertalanffy growth function predicts the pattern of growth over time for that particular species. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ) which together characterize the shape of the length-at-age growth relationship.

Length at reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved then subsequently cut into five-micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of mature samples for a given length interval are assembled for each sex and these data are fitted to a three- or four-parameter logistic function to determine the best fit of these data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region), such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes, standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a 3- or 4-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age & growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted

logistic function provides a direct estimate of the age at which 50% of fish of a particular species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** Sources of data are directly derived from research cruises sampling and market samples collected by the Guam-contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 55 for specific details on data sources by species.

**Parameter definitions:**

**$T_{max}$  (maximum age)** – The maximum observed age revealed from an otolith-based age determination study.  $T_{max}$  values can be derived from ages determined by annuli counts of sagittal otolith sections and/or bomb radiocarbon ( $^{14}\text{C}$ ) analysis of otolith core material. Units are years.

**$L_{\infty}$  (asymptotic length)** – One of three coefficients of the von Bertalanffy growth function (VBGF) that measures the mean maximum length at which the growth curve plateaus and no longer increases in length with increasing age. This coefficient reflects the mean maximum length and not the observed maximum length. Units are centimeters.

**$k$  (growth coefficient)** – One of three coefficients of the VBGF that measures the shape and steepness by which the initial portion of the growth function approaches its mean maximum length ( $L_{\infty}$ ).

**$t_0$  (hypothetical age at length zero)** – One of three coefficients of the VBGF whose measure is highly influenced by the other two VBGF coefficients ( $k$  and  $L_{\infty}$ ) and typically assumes a negative value when specimens representing early growth phases (0+ to 1+ ages) are not available for age determination. Units are years.

**$M$  (natural mortality)** – This is a measure of mortality rate for a fish stock not under the influence of fishing pressure and is considered to be directly related to stock productivity (i.e., high  $M$  indicates high productivity and low  $M$  indicates low stock productivity).  $M$  can be derived through use of various equations that link  $M$  to  $T_{max}$  and  $k$ , or in some instances, by calculating the value of the slope from a regression fit to a declining catch curve (regression of the natural logarithm of abundance versus age class) derived from fishing an unfished or lightly fished population.

**$A_{50}$  (age at 50% maturity)** – Age at which 50% of the sampled stock under study has attained reproductive maturity. This parameter is best determined based on studies that concurrently

determine both age (otolith-based age data) and reproductive maturity status (logistic function fitted to percent mature by age class with maturity determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A_{50}$  is to use an existing  $L_{50}$  estimate to find the corresponding age ( $A_{50}$ ) from an existing VBGF curve. Units are years.

**$A\Delta_{50}$  (age of sex switching)** – Age at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal. This parameter is best determined based on studies that concurrently determines both age (otolith-based age data) and reproductive sex reversal status (logistic function fitted to percent sex reversal by age class with sex reversal determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A\Delta_{50}$  is to use an existing  $L\Delta_{50}$  estimate to find the corresponding age ( $A\Delta_{50}$ ) from the VBGF curve. Units are years.

**$L_{50}$  (length at which 50% of a fish species are capable of spawning)** – Length (usually in terms of fork length) at which 50% of the females of a sampled stock under study has attained reproductive maturity; this is the length associated with  $A_{50}$  estimates. This parameter is derived using a logistic function to fit the percent mature data by length class with maturity status best determined via microscopic analyses of gonad histology preparations).  $L_{50}$  information is typically more available than  $A_{50}$  since  $L_{50}$  estimates do not require knowledge of age & growth. Units are centimeters

**$L\Delta_{50}$  (length of sex switching)** – Length (usually in terms of fork length) at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal; this is the length associated with  $A\Delta_{50}$  estimates. This parameter is derived using a logistic function to fit the percent sex reversal data by length class with sex reversal status best determined via microscopic analyses of gonad histology preparations.  $L\Delta_{50}$  information is typically more available than  $A\Delta_{50}$  since  $L\Delta_{50}$  estimates do not require knowledge of age & growth. Units are centimeters.

**Rationale:** These nine life history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. Currently, the assessment of coral reef fish resources in Guam is data-limited. Knowledge of these life history parameters support current efforts to characterize the resilience of these resources and also provide important biological inputs for future stock assessment efforts and enhance our understanding of the species' likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Table 55. Available age, growth, and reproductive maturity information for coral reef species targeted for otoliths and gonads sampling in Guam**

| Species          | Age, growth, and reproductive maturity parameters |                   |                   |                     |                   |                   |                   | Reference        |
|------------------|---|-------------------|-------------------|---------------------|-------------------|-------------------|-------------------|------------------|
|                  | $T_{max}$   | $L_{\infty}$      | $k$               | $t_0$               | $A_{50}$          | $L_{50}$          | $L\Delta_{50}$    |                  |
| <i>Calatomus</i> | 3 <sup>d</sup>                                    | 26.3 <sup>d</sup> | 0.91 <sup>d</sup> | -0.065 <sup>d</sup> | 1.14 <sup>d</sup> | 16.8 <sup>d</sup> | 21.3 <sup>d</sup> | Taylor and Choat |

|                                  |                 |                                    |                                    |                                      |                                  |                                    |                   |                         |
|----------------------------------|-----------------|------------------------------------|------------------------------------|--------------------------------------|----------------------------------|------------------------------------|-------------------|-------------------------|
| <i>carolinus</i>                 |                 |                                    |                                    |                                      |                                  |                                    |                   | (2014)                  |
| <i>Oxycheilinus unifasciatus</i> |                 |                                    |                                    |                                      |                                  |                                    |                   |                         |
| <i>Chlorurus frontalis</i>       | 11 <sup>d</sup> | 37.2 <sup>d</sup>                  | 0.71 <sup>d</sup>                  | -0.058 <sup>d</sup>                  | 1.55 <sup>d</sup>                | 24.0 <sup>d</sup>                  | 34.3 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Chlorurus microrhinos</i>     | 11 <sup>d</sup> | 45.7 <sup>d</sup>                  | 0.34 <sup>d</sup>                  | -0.097 <sup>d</sup>                  | 3.7 <sup>d</sup>                 | 30.8 <sup>d</sup>                  | 37.8 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Chlorurus spilurus</i>        | 9 <sup>d</sup>  | 21.8 <sup>d</sup>                  | 0.95 <sup>d</sup>                  | -0.075 <sup>d</sup>                  | 1.3 <sup>d</sup>                 | 14.4 <sup>d</sup>                  | 20.7 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Hipposcarus longiceps</i>     | 10 <sup>d</sup> | 39.6 (f),<br>46.6 (m) <sup>d</sup> | 0.97 (f),<br>0.67 (m) <sup>d</sup> | -0.04 (f),<br>-0.05 (m) <sup>d</sup> |                                  | 40.1 <sup>d</sup>                  |                   | Taylor and Cruz (2017)  |
| <i>Naso lituratus</i>            | 13 <sup>d</sup> | 20.4 <sup>d</sup>                  | 0.93 <sup>d</sup>                  | -0.030 <sup>d</sup>                  | 2.4 (m) <sup>d</sup>             | 14.5 (f),<br>17.8 (m) <sup>d</sup> |                   | Taylor et al. (2014)    |
| <i>Naso unicornis</i>            | 23 <sup>d</sup> | 49.3 <sup>d</sup>                  | 0.22 <sup>d</sup>                  | -0.048 <sup>d</sup>                  | 4.0 (f),<br>3.2 (m) <sup>d</sup> | 29.2 (f),<br>27.1 (m) <sup>d</sup> |                   | Taylor et al. (2014)    |
| <i>Scarus altipinnis</i>         | 14 <sup>d</sup> | 33.9 <sup>d</sup>                  | 0.66 <sup>d</sup>                  | -0.69 <sup>d</sup>                   | 2.89 <sup>d</sup>                | 25.1 <sup>d</sup>                  | 33.7 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Scarus forsteni</i>           | 12 <sup>d</sup> | 28.1 <sup>d</sup>                  | 0.88 <sup>d</sup>                  | -0.62 <sup>d</sup>                   | 1.79 <sup>d</sup>                | 21.6 <sup>d</sup>                  | 27.1 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Scarus psittacus</i>          | 6 <sup>d</sup>  | 20.7 <sup>d</sup>                  | 0.91 <sup>d</sup>                  | -0.083 <sup>d</sup>                  | 1.36 <sup>d</sup>                | 10.3 <sup>d</sup>                  | 19.3 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Scarus rubroviolaceus</i>     | 6 <sup>d</sup>  | 37.6 <sup>d</sup>                  | 0.66 <sup>d</sup>                  | -0.062 <sup>d</sup>                  | 1.91 <sup>d</sup>                | 27.1 <sup>d</sup>                  | 32.9 <sup>d</sup> | Taylor and Choat (2014) |
| <i>Scarus schlegeli</i>          | 8 <sup>d</sup>  | 25.2 <sup>d</sup>                  | 1.03 <sup>d</sup>                  | -0.06 <sup>d</sup>                   | 1.99 <sup>d</sup>                | 19.7 <sup>d</sup>                  | 22.0 <sup>d</sup> | Taylor and Choat (2014) |

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (including in press).

Parameter estimates are for females unless otherwise noted (F=females, M=males). Parameters  $T_{max}$ ,  $t_0$ ,  $A_{50}$ , and  $A\Delta_{50}$  are in units of years;  $L_{\infty}$ ,  $L_{50}$ , and  $L\Delta_{50}$  are in units of mm fork length (FL);  $k$  in units of year<sup>-1</sup>; X=parameter estimate too preliminary or Y=published age and growth parameter estimates based on DGI numerical integration technique and likely to be inaccurate; NA=not applicable. Superscript letters indicate status of parameter estimate (see footnotes below table). Published or in press publications (<sup>d</sup>) are denoted in “Reference” column.

### 2.2.3.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Biosampling Program started in 2009. This program has two components: first is the Field/Market Sampling Program and the second is the

Life History Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program are:

- Broad scale look at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification;
- Develop accurate local length-weight curves.

In the Guam, the Biosampling is focused on the commercial coral reef spear fishery with occasional sampling of the bottomfish fishery occurring locally and less frequently at the banks. Sampling is conducted in direct partnership with the spear fisherman. The Market Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** NMFS Biosampling Program

**Parameter definition:**

*L<sub>max</sub>* – **maximum fish length** is the longest fish per species recorded in the Biosampling Program from the commercial spear fishery. This value is derived from measuring the fork length of individual samples for species occurring in the spear fishery.

*L<sub>bar</sub>* – **mean length** is the average value of all lengths recorded from the commercial spear fishery. This can be influenced by gear selectivity since the commercial spear fishery has a typical size target based on customer demand. This can also be influenced by size regulations.

*n* – **sample size** is the total number of samples accumulated for each species recorded in the commercial spear fishery.









*N<sub>L-W</sub>* – **sample size for L-W regression** is the number of samples used to generate the *a* & *b* coefficients.









*a* & *b* – **length-weight coefficients** are the coefficients derived from the regression line fitted to all length and weight measured per species in the commercial spear fishery. These values are used to convert length information to weight. Values are influenced by the life history characteristics of the species, geographic location, population status, and nature of the fisheries from which the species are harvested

**Rationale:** Length-derived information is being used as an indicator of population status particularly for data-poor stocks like coral reef fish. Average length ( $L_{bar}$ ) was used as a principal stock assessment indicator variable for exploited reef fish population (Nadon et al., 2015). Average length was also shown to be correlated with population size (Kerr and Dickle, 2001). Maximum length ( $L_{max}$ ), typically coupled with maximum age, is typically used as a proxy for fish longevity which has implications on the productivity and susceptibility of a species to fishing pressure. The length-weight coefficients ( $a$  &  $b$  values) are used to convert length to weight for fishery dependent and fishery independent data collection where length are typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the CNMI coral reef and bottomfish fisheries.

**Table 56. Available length derived information for various coral reef species in Guam**

| Species                      | Length-derived parameters |           |        |        |        | Reference                            |
|------------------------------|---------------------------|-----------|--------|--------|--------|--------------------------------------|
|                              | $L_{max}$                 | $L_{bar}$ | $n$    | $a$    | $b$    |                                      |
| <i>Naso unicornis</i>        | 57.2                      | 38.02     | 15,461 | 0.0278 | 2.9135 | 2010-2015 Guam Bio-Sampling Database |
| <i>Naso lituratus</i>        | 29.6                      | 21.35     | 16,702 | 0.0223 | 3.0264 |                                      |
| <i>Acanthurus lineatus</i>   | 28.9                      | 19.04     | 4,325  | 0.0473 | 2.8110 |                                      |
| <i>Scarus altipinnis</i>     | 46.4                      | 31.16     | 3,913  | 0.0207 | 3.0040 |                                      |
| <i>Myripristis bendti</i>    | 29.4                      | 18.63     | 3,903  | 0.0858 | 2.5911 |                                      |
| <i>Siganus argenteus</i>     | 34.5                      | 21.71     | 3,653  | 0.0163 | 3.0428 |                                      |
| <i>Acanthurus nigricauda</i> | 29.1                      | 21.40     | 3,500  | 0.0511 | 2.7811 |                                      |
| <i>Hipposcarus longiceps</i> | 51.4                      | 37.30     | 3,149  | 0.0172 | 3.0320 |                                      |
| <i>Scarus schlegeli</i>      | 36.2                      | 25.19     | 2,787  | 0.0205 | 3.0033 |                                      |
| <i>Siganus punctatus</i>     | 32.0                      | 23.97     | 2,619  | 0.0199 | 3.0690 |                                      |
| <i>Monotaxis grandoculis</i> | 48.9                      | 29.17     | 2,388  | 0.0440 | 2.8384 |                                      |
| <i>Scarus rubroviolaceus</i> | 47.8                      | 31.91     | 2,192  | 0.0114 | 3.1812 |                                      |
| <i>Lethrinus obsoletus</i>   | 34.7                      | 22.15     | 2,273  | 0.0169 | 3.0471 |                                      |
| <i>Scarus forsteni</i>       | 39.1                      | 28.13     | 1,801  | 0.0149 | 3.1169 |                                      |
| <i>Lutjanus gibbus</i>       | 43.5                      | 29.99     | 1,687  | 0.0195 | 3.0274 |                                      |
| <i>Parupeneus insularis</i>  | 28.5                      | 21.89     | 1,560  | 0.0178 | 3.0865 |                                      |
| <i>Siganus spinus</i>        | 27.5                      | 16.53     | 1,670  | 0.0353 | 2.7886 |                                      |
| <i>Lethrinus atkinsoni</i>   | 33.7                      | 21.93     | 1,644  | 0.0215 | 3.0217 |                                      |
| <i>Chlorurus microrhinus</i> | 50.5                      | 32.54     | 1,527  | 0.0187 | 3.0520 |                                      |
| <i>Chlorurus sordidus</i>    | 33.1                      | 22.39     | 1,234  | 0.0208 | 3.0293 |                                      |
| <i>Kyphosus cinerascens</i>  | 50.7                      | 29.94     | 1,146  | 0.0323 | 2.9267 |                                      |









**Figure 8. Average length over time of representative Guam CREMUS derived from the Biosampling Program; continued from previous two pages**

## 2.2.4 Guam Bottomfish Ecosystem – Bottomfish Life History

### 2.2.4.1 Age & Growth and Reproductive Maturity

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely-cut thin sections of sagittal otoliths. Validated age determination is based on an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally-based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. The relation between age and fish length is evaluated by fitting this data to a von Bertalanffy growth function based on statistical analyses. The resulting von Bertalanffy

growth function predicts the pattern of growth over time for that particular species. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ) which together characterize the shape of the length-at-age growth relationship.

Length-at-reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved then subsequently cut into five micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of mature samples for a given length interval are assembled for each sex and these data are fitted to a three- or four-parameter logistic function to determine the best fit of these data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region), such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes, standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a three- or four-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age & growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted logistic function provides a direct estimate of the age at which 50% of fish of a particular species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** Sources of data are directly derived from research cruises sampling and market samples collected by the Guam-contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 57 for specific details on data sources by species.

**Parameter definitions:**

**$T_{max}$  (maximum age)** – The maximum observed age revealed from an otolith-based age determination study.  $T_{max}$  values can be derived from ages determined by annuli counts of sagittal otolith sections and/or bomb radiocarbon ( $^{14}\text{C}$ ) analysis of otolith core material. Units are years.

**$L_{\infty}$  (asymptotic length)** – One of three coefficients of the von Bertalanffy growth function (VBGF) that measures the mean maximum length at which the growth curve plateaus and no longer increases in length with increasing age. This coefficient reflects the mean maximum length and not the observed maximum length. Units are centimeters.

**$k$  (growth coefficient)** – One of three coefficients of the VBGF that measures the shape and steepness by which the initial portion of the growth function approaches its mean maximum length ( $L_{\infty}$ ).

**$t_0$  (hypothetical age at length zero)** – One of three coefficients of the VBGF whose measure is highly influenced by the other two VBGF coefficients ( $k$  and  $L_{\infty}$ ) and typically assumes a negative value when specimens representing early growth phases (0+ to 1+ ages) are not available for age determination. Units are years.

**$M$  (natural mortality)** – this is a measure of mortality rate for a fish stock not under the influence of fishing pressure and is considered to be directly related to stock productivity (i.e., high  $M$  indicates high productivity and low  $M$  indicates low stock productivity).  $M$  can be derived through use of various equations that link  $M$  to  $T_{max}$  and two VBGF coefficients ( $k$  and  $L_{\infty}$ ) or by calculating the value of the slope from a regression fit to a declining catch curve (regression of the natural logarithm of abundance versus age class) derived from fishing an unfished or lightly fished population.

**$A_{50}$  (age at 50% maturity)** – Age at which 50% of the sampled stock under study has attained reproductive maturity. This parameter is best determined based on studies that concurrently determine both age (otolith-based age data) and reproductive maturity status (logistic function fitted to percent mature by age class with maturity determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A_{50}$  is to use an existing  $L_{50}$  estimate to find the corresponding age ( $A_{50}$ ) from an existing VBGF curve. Units are years.

**$A\Delta_{50}$  (age of sex switching)** – Age at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal. This parameter is best determined based on studies that concurrently determines both age (otolith-based age data) and reproductive sex reversal status (logistic function fitted to percent sex reversal by age class with sex reversal determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A\Delta_{50}$  is to use an existing  $L\Delta_{50}$  estimate to find the corresponding age ( $A\Delta_{50}$ ) from the VBGF curve. Units are years.

**$L_{50}$  (length at which 50% of a fish species are capable of spawning)** – Length (usually in terms of fork length) at which 50% of the females of a sampled stock under study has attained reproductive maturity; this is the length associated with  $A_{50}$  estimates. This parameter is derived using a logistic function to fit the percent mature data by length class with maturity status best determined via microscopic analyses of gonad histology preparations).  $L_{50}$  information is

typically more available than  $A_{50}$  since  $L_{50}$  estimates do not require knowledge of age & growth. Units are centimeters.

**$L\Delta_{50}$  (length of sex switching)** – Length (usually in terms of fork length) at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal; this is the length associated with  $A\Delta_{50}$  estimates. This parameter is derived using a logistic function to fit the percent sex reversal data by length class with sex reversal status best determined via microscopic analyses of gonad histology preparations.  $L\Delta_{50}$  information is typically more available than  $A\Delta_{50}$  since  $L\Delta_{50}$  estimates do not require knowledge of age & growth. Units are centimeters.

**Rationale:** These nine life history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. Currently, the assessment of coral reef fish resources in Guam is data-limited. Knowledge of these life history parameters support current efforts to characterize the resilience of these resources and also provide important biological inputs for future stock assessment efforts and enhance our understanding of the species' likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Table 57. Available age, growth, and reproductive maturity information for bottomfish species targeted for life history sampling (otoliths and gonads) in Guam**

| Species                            | Age, growth, and reproductive maturity parameters |                |                |                |                |          |                |                  |                | Reference                    |
|------------------------------------|---|----------------|----------------|----------------|----------------|----------|----------------|------------------|----------------|------------------------------|
|                                    | $T_{max}$   | $L_{\infty}$   | $k$            | $t_0$          | $M$            | $A_{50}$ | $A\Delta_{50}$ | $L_{50}$         | $L\Delta_{50}$ |                              |
| <i>Aphareus rutilans</i>           |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Aprion virescens</i>            |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Etelis carbunculus</i>          |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Etelis coruscans</i>            |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Monotaxis grandoculis</i>       | X <sup>a</sup>                                    | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> |          |                |                  |                | Cruz et al. (in prep.)       |
| <i>Pristipomoides auricilla</i>    | X <sup>c</sup>                                    | X <sup>c</sup> | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> |          | NA             |                  | NA             | O'Malley et al. (in review)  |
| <i>Pristipomoides filamentosus</i> | X <sup>a</sup>                                    | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> |          | NA             |                  | NA             | Villagomez et al. (in prep.) |
| <i>Pristipomoides flavipinnis</i>  |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Pristipomoides sieboldii</i>    |   |                |                |                |                |          | NA             |                  | NA             |                              |
| <i>Pristipomoides zonatus</i>      | X <sup>a</sup>                                    | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> | X <sup>a</sup> |          | NA             |                  | NA             | LHP (in prep.)               |
| <i>Variola louti</i>               |   |                |                |                |                |          |                | 220 <sup>b</sup> | X <sup>a</sup> |                              |

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (+ in press).

Parameter estimates are for females unless otherwise noted (F=females, M=males). Parameters  $T_{max}$ ,  $t_0$ ,  $A_{50}$ , and  $A\Delta_{50}$  are in units of years;  $L_{\infty}$ ,  $L_{50}$ , and  $L\Delta_{50}$  are in units of mm fork length (FL);  $k$  in units of year<sup>-1</sup>; X=parameter estimate too preliminary or Y=published age and growth parameter estimates based on DGI numerical integration technique and likely to be inaccurate; NA=not applicable. Superscript letters indicate status of parameter estimate (see footnotes below table). Published or in press publications (<sup>d</sup>) are denoted in “Reference” column.

#### 2.2.4.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Biosampling Program started in 2009. This program has two components: first is the Field/Market Sampling Program and the second is the Life History Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program are:

- Broad scale look at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification;
- Develop accurate local length-weight curves.

In Guam, the Biosampling is focused on the commercial fishery. Sampling is conducted in partnership with the Guam Fisherman’s Cooperative Association (GFCA). The Market Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information. More specific fishery information such as gear information, species composition and total catch information is recorded through the log book system implemented by GFCA and transcribed into the database maintained by the Western Pacific Fishery Information Network.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** Guam

**Spatial Scale:** Island

**Data Source:** NMFS Biosampling Program

**Parameter definition:**

$L_{max}$  – *maximum fish length* is the longest fish per species recorded in the Biosampling Program from the commercial spear fishery. This value is derived from measuring the fork length of individual samples for species occurring in the spear fishery. Units are centimeters.

$L_{bar}$  – *mean length* is the average value of all lengths recorded from the commercial spear fishery. This can be influenced by gear selectivity since the commercial spear fishery has a

typical size target based on customer demand. This can also be influenced by size regulations. Units are centimeters.

*n* – **sample size** is the total number of samples accumulated for each species recorded in the commercial spear fishery.


*N<sub>L-W</sub>* – **sample size for L-W regression** is the number of samples used to generate the a & b coefficients

*a* & *b* – **length-weight coefficients** are the coefficients derived from the regression line fitted to all length and weight measured per species in the commercial spear fishery. These values are used to convert length information to weight. Values are influenced by the life history characteristics of the species, geographic location, population status, and nature of the fisheries from which the species are harvested from.

**Rationale:** Length-derived information is being used as an indicator of population status particularly for data-poor stocks like coral reef fish. Average length (*L<sub>bar</sub>*) was used as a principal stock assessment indicator variable for exploited reef fish population (Nadon et al., 2015). Average length was also shown to be correlated with population size (Kerr and Dickle, 2001). Maximum length (*L<sub>max</sub>*), typically coupled with maximum age, is typically used as a proxy for fish longevity which has implications on the productivity and susceptibility of a species to fishing pressure. The length-weight coefficients (*a* & *b* values) are used to convert length to weight for fishery dependent and fishery independent data collection where length are typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the CNMI coral reef and bottomfish fisheries.

**Table 58. Available length derived information for various bottomfish species in Guam**

| Species                           | Length-derived parameters |                        |          |          |          | Reference                           |
|-----------------------------------|---------------------------|------------------------|----------|----------|----------|-------------------------------------|
|                                   | <i>L<sub>max</sub></i>    | <i>L<sub>bar</sub></i> | <i>n</i> | <i>a</i> | <i>b</i> |                                     |
| <i>Lethrinus rubrioperculatus</i> | 46.6                      | 27.10                  | 3374     | 0.0248   | 2.9158   | 2010-2015 Guam Biosampling Database |
| <i>Epinephelus fasciatus</i>      | 35.8                      | 24.01                  | 3033     | 0.0141   | 3.0303   |                                     |
| <i>Pristipomoides auricilla</i>   | 39.0                      | 28.18                  | 1732     | 0.0152   | 3.0742   |                                     |



**Figure 9. Average length over time of representative Guam BMUS derived from the Biosampling Program**

## 2.2.5 References

- Kerr, S.R. and L.M. Dickie. 2001. *The biomass spectrum: a predator-prey theory of aquatic production*. Columbia University Press.
- Nadon, M.O., Ault, J.S., Williams, I.D., Smith, S.G., and G.T. DiNardo. 2015. Length-based assessment of coral reef fish populations in the Main and Northwestern Hawaiian Islands. *PLoS One*, 10(8), p.e0133960.
- Ralston S.V. and H.A. Williams. 1988. Depth distributions, growth, and mortality of deep slope fishes from the Mariana Archipelago. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-SWFC-113, 47 pp.
- Taylor, B.M. and J.H. Choat. 2014. Comparative demography of commercially important parrotfish species from Micronesia. *J. Fish Biol.* 84, pp. 383-402.
- Taylor, B.M., Rhodes, K.L., Marshall, A., and J.L. McIlwain. 2014. Age-based demographic and reproductive assessment of orangespine *Naso lituratus* and bluespine *Naso unicornis* unicornfishes. *J. Fish Biol.* 85, pp. 901–916. <https://doi:10.1111/jfb.12479>.
- Taylor, B.M., Gourley J., and M.S. Trianni. 2016. Age, growth, reproductive biology and spawning periodicity of the fork-tail rabbitfish (*Siganus argenteus*) from the Mariana Islands. *Marine & Freshwater Research*. <https://doi.org/10.1071/MF16169>.

Taylor B.M. and E. Cruz. 2017. Age-based and reproductive biology of the Pacific Longnose Parrotfish *Hipposcarus longiceps* from Guam. *PeerJ*, 5:e4079.  
<https://doi.org/10.7717/peerj.4079>.

Taylor B.M., Oyafuso Z.S., and M.S. Trianni. 2017. Life history of the orange-striped emperor *Lethrinus obsoletus* from the Mariana Islands. *Ichth. Res.*, 4, pp. 423–432  
<https://doi.org/10.1007/s10228-017-0573-8>.



## 2.3 SOCIOECONOMICS

This section outlines the pertinent economic, social, and community information available for assessing the successes and impacts of management measures or the achievements of the Fishery Ecosystem Plan for the Marianas Archipelago (Western Pacific Regional Fishery Management Council, 2016). It meets the objective “Support Fishing Communities” adopted at the 165<sup>th</sup> Council meeting; specifically, it identifies the various social and economic groups within the region’s fishing communities and their interconnections. The section begins with an overview of the socioeconomic context for the region, and then provides a summary of relevant studies and data for CNMI and Guam, followed by summaries of relevant studies and data for each fishery in CNMI and Guam.

In 1996, the Magnuson-Stevens Fishery Conservation and Management Act’s National Standard 8 (NS8) specified that conservation and management measures take into account the importance of fishery resources to fishing communities, to provide for their sustained participation in fisheries and to minimize adverse economic impacts, provided that these considerations do not compromise the achievement of conservation. Unlike other regions of the U.S., the settlement of the Western Pacific region was intimately tied to the sea (Figure 10), which is reflected in local culture, customs, and traditions.



Figure 10. Settlement of the Pacific Islands, courtesy of Wikimedia Commons, [https://commons.wikimedia.org/wiki/File:Polynesian\\_Migration.svg](https://commons.wikimedia.org/wiki/File:Polynesian_Migration.svg).

Polynesian voyagers relied on the ocean and marine resources on their long voyages in search of new islands, as well as in sustaining established island communities. Today, the population of the region also represents many Asian cultures from Pacific Rim countries, which reflect similar importance of marine resources. Thus, fishing and seafood are integral local community ways of life. This is reflected in the amount of seafood eaten in the region relative to the rest of the United States, as well as the language, customs, ceremonies, and community events. Because fishing is such an integral part of the culture, it is difficult to discern commercial from non-commercial fishing as most trips involving multiple motivations and multiple uses of the fish caught. While economics are an important consideration, fishermen report other motivations, such as customary exchange, as being equally important. Due to changing economies and westernization, recruitment of younger fishermen has become a concern for the sustainability of fishing and fishing traditions in the region.

The Marianas Archipelago consists of the Commonwealth of the Northern Mariana Islands (CNMI) at the northern end and Guam, the southernmost island. These are typically treated as two jurisdictions, which will be presented separately in the rest of this section despite being grouped under one FEP.

### **2.3.1 Response to Previous Council Recommendations**

At its 173<sup>rd</sup> meeting held in Wailea, HI, the Council recommended NMFS to address data gaps and research needs for ESA-listed shark species, such as

- improving data collection for oceanic whitetip shark capture data in non-longline pelagic fisheries; and
- conducting outreach to fishermen to improve species identification for shark species to facilitate improved accurate catch data reporting.

At the 174<sup>th</sup> Council meetings in Saipan and Guam, PIFSC staff met with members of the Marianas fishing community to discuss their concerns related to shark depredation for both insular and pelagic fisheries across the Marianas Archipelago and consider possible research opportunities.

Current research by PIFSC social scientists is working towards identifying patterns in oceanic whitetip shark interactions, handling practices, and perceptions within West Hawaii's non-longline pelagic fisheries, based on interviews with the fishing community (Oct. 2017 - Sept. 2018). The goal of this research is to identify opportunities and barriers to engaging fishermen in science- and management-based efforts (like shark identification, catch reporting, and mortality reduction). Research results are expected by May 2019.

PIFSC received funding in early 2019 to extent this West Hawaii research to both insular and pelagic fisheries in the Marianas. Research will be conducted in 2019-2020 with the goal to engage the Marianas fishing community to better understand the nature of shark interactions and explore mitigation techniques aligned with community needs and values.

## 2.3.2 CNMI

### 2.3.2.1 Introduction

An overview of CNMI history, culture, geography, and relationship with the U.S. is described in the Fishery Ecosystem Plan for the Mariana Archipelago (Western Pacific Regional Fishery Management Council, 2016). Over the past decade, a number of studies have synthesized more specifics about the role of fishing and marine resources across CNMI, as well as information about the people who engage in the fisheries or use fishery resources.

The ancestors of the indigenous Chamorro first arrived in the Marianas around 3,500 years ago and relied on seafood as their principal source of protein (Allen and Amesbury, 2012, and Grace McCaskey, 2014). Similar to other archipelagos in the Western Pacific, fish and marine resources have played a central role in shaping the social, cultural, and economic fabric of the CNMI that continues today. They fished for both reef and pelagic species, collected mollusks and other invertebrates, and caught sea turtles. The occupation of CNMI by foreign nations dramatically changed the island's ecosystems, reshaped communities, and disrupted fishing traditions. In the 17<sup>th</sup> and 18<sup>th</sup> centuries, Spanish colonizers destroyed the Chamorro's seagoing canoes, suppressed offshore fishing practices, and relocated populations from their traditional home. The CNMI was briefly occupied by Germany from 1899 to the beginning of WWII. During WWII, the CNMI was occupied by the Japanese military, and then was captured by the United States. Throughout this time, fishing remained an important activity. Later immigrants to the islands from East and Southeast Asia also possessed a strong fishing tradition. Today, only Saipan, Rota, and Tinian are permanently inhabited, with 90% of the population living on the island of Saipan. Although the CNMI has transitioned to a tourism-based economy, fishing still plays an important cultural role and serves as a reliable source of local food (Ayers, 2018).

### 2.3.2.2 People who Fish

Allen and Amesbury (2012) summarized results of studies that demonstrated the sociocultural importance of fishing to Saipan residents. In a 2005 study, most of the active or commercial fishermen who responded to the survey had fished for more than 10 years. They most often participated in snorkel spearfishing at night (participated in by 73% of the fishermen) and snorkel spear fishing during daytime (58% of the fishermen), followed by hook-and-line less than 100 ft. deep (36%), trolling (21%), cast net (talaya; 14%), hook-and-line more than 100 ft. deep (9%), trapping (octopus, crabs, etc.; 19%), and foraging the reef (8%); 18% said they participated in one or more other techniques. Less than a third (~30%) said they owned a boat. The primary reasons for fishing were social, cultural, and nutrition; in addition to reporting that they enjoy the activity itself (32%), many said they needed the fish to feed their family (23%), give to family and friends to strengthen social bonds (13%), that their family has always fished (12%), and that it strengthens bonds with their children/family (6%). Only 4% said they needed the money from the fish they sold. Other motivations included strengthening the bond with their fellow fishermen, fishing to catch fish for festivals and parties, and seasonal fishing for manahak, ti'ao, and i'e (2% each).

The fishermen reported fishing an average of 71 days per year, with 26% going once every two to three days, and 24% fishing once every two weeks. Those surveyed also reported a decrease in the amount of time they have spent fishing in the past decade, fishing 93 days per year on

average. Saipan reef fish were the most frequently harvested species (caught by 54% of the fishermen), followed by shallow-water bottomfish (23%) and reef invertebrates such as octopus, shellfish and crabs (14%).

As in other parts of the region, much of the fisher's catch in the CNMI was consumed by themselves and their immediate family (70%), with another 20% consumed by extended family and friends. Only 8% of the catch was sold. There were 18 respondents that identified themselves as commercial fishermen. They reported a median monthly income of \$200 from fishing, with average monthly income of just over \$1,000. Costs exceeded sales for almost every income category for fishermen, suggesting that fishing is not a business for most, but that catch is simply sold to cover some of the cost.

While fish remain an important part of the local diet and an integral part of the people's history and culture, adaptation to and integration with a more westernized lifestyle appears to have changed people's dietary preferences on Saipan. Nearly half (45%) of the survey respondents reported eating "somewhat less fish" than they did a decade ago, although the majority still ate fish between one and three times a week. The majority also purchased their fish from a store or restaurant (40%), while 31% purchased fish from roadside vendors. Less common was acquiring fish from an extended relative/friend (13%) or their own catch (11%). Most of the fish consumed came from the U.S. mainland (41%), with other important sources coming from Saipan's coral reefs (31%), deepwater or pelagic fish caught off of Saipan (23%), or fish imported from other Pacific islands (e.g. Chuuk ; 10%).

Few other surveys have been conducted on fishing in the CNMI. A household survey conducted in 2012 found that 37% of households had at least one individual that self-identified as a fisherman (Kotowicz and Allen, 2015). Respondents from fishing households tended to be younger, possess lower education levels, and have a higher rate of unemployment than respondents from non-fishing households.

While proportionally few residents own a boat, more than 400 vessels were registered in the CNMI small boat fleet between 2010 and 2011 (Allen and Amesbury, 2012). More than 200 of the vessels were active and operating in CNMI waters at that time, and more than 100 of the vessels were involved in fishing activities. The active small boat fleet targeted tunas, other small pelagics (through trolling), and bottomfish; with the increase in gas prices, however, pelagic fishing has waned. When caught, these fish are marketed locally, given away to family and friends, or used for ceremonial purposes such as parties, culturally significant fiestas, and the patron saint's days for each village.

On Saipan, fisheries managers estimated the active small boat fleet at approximately 100 vessels from 2010 to 2011. Full-time commercial fishing is primarily conducted by ethnic nonindigenous minorities, namely Filipino residents that fish primarily as independent owners and/or operators and recent immigrants from the Federated States of Micronesia that fish for income. Chamorro and Carolinians, in contrast, primarily fish for recreational and subsistence purposes, typically only selling catch to recoup costs. A few vessel owner operators are considered "pescadors", a term used to refer to fishermen who provide fish for important community and familial events. Pescadors customarily provide 100-200 lbs. of reef fish for cooked dishes and pelagic species for kelaguen (a raw fish dish) used in community and family

celebrations. The system of seafood distribution underwent significant changes from approximately the turn of the century with the establishment of large seafood vendors. In contrast to individual fishermen/vendors who only market their own catch, large vendors typically own and operate a number of vessels and purchase catch from independent fishermen to sell. This trend has reportedly caused prices to decline. In addition, increases in fuel prices, low market prices for fish, and downturns in the domestic economy have led to a general decline in participation in this fishery since 2000 in numbers of fishermen, trips, landings, and seafood purchasers. The Saipan Fishermen's Association (SFA) is a nonprofit organization established in 1985 that holds annual fishing derbies and participated in community involvement projects, such as beach cleanup.

On Tinian, estimates of fleet size range from 15 to 20 vessels in 2010-2011. An estimated one to three fishermen fished consistently with the primary intent of selling fish. Respondents suggested that fishing and eating of fish was more habitual, rather than geared toward a particular event. Increasing fuel prices have reportedly led to the decline in number of active fishermen, and fishermen frequently have sold fish to cover fuel costs. Three restaurants and two stores in Tinian purchase fish, although fishermen have also resorted to selling house-to-house; the fishermen commonly have an established clientele. A few charter boats serve tourist clientele, however they do not land much catch, and even trolling trips serve more as photo opportunities. Charter boats are reportedly owned by non-local residents and target tourists by their country of origin (e.g. Japan, China, or Korea).

On Rota, fishermen target pelagic species when in season and bottomfish the rest of the year. Like on the other islands, the number and activity of fishermen have declined as a result of increased fuel prices. Family members will often make requests for certain kinds of fish, but they will also contribute money to purchase fuel for a fishing trip. In addition, fishermen will often check demand with local restaurants. In 2010 and 2011, fishermen sold catch to three separate restaurants or to neighbors and friends within the community (door-to-door or from a cooler on the roadside). One general store sold fish caught by a family member, who fished specifically to sell to that store. Rota holds a fishing derby in celebration of San Francisco, saint of the island.

A survey of the small boat fleet was also conducted in 2011 (Hospital and Beavers, 2014). Respondents were 41 years old and had been boat fishing for 15 years on average, providing evidence of a deep tradition of boat fishing in the CNMI. They were more likely to identify themselves as Chamorro relative to the general population of the CNMI, although they were equally likely to have been born in the CNMI. In general, fishermen were more educated than the general population and of comparable affluence. Pelagic trolling was the most popular gear type, followed by deepwater bottomfishing, shallow-water bottomfishing, and spearfishing. Most fishermen (71%) reported fishing adjacent to a Fish Aggregating Device (FAD) at some point in the past 12 months, and did so on nearly 22% of their fishing trips. A high degree of seasonal fishing effort was reported across most fishing fleet subgroups, though fishermen on Tinian and Rota were more likely to fish year-round than those on Saipan.

A majority of fishermen (74%) reported selling at least a portion of their catch in the past year. However, less than half of survey respondents (43%) indicated that they could always sell any fish that they wanted. A significant percentage of fish caught was consumed at home (28%) or given away to relatives, friends, or for cultural events (38%); this reflects the strong family and

social connections associated with fishing in the CNMI. Approximately 29% of fish catch was sold, with the remaining catch either released (2%) or exchanged for goods and services (3%). Even fishermen who regularly sold fish still retained approximately 22% of their catch for home consumption, participation in traditional fish-sharing networks, and customary exchange. Additionally, 91% of survey respondents considered the bottomfish they catch to be an important source of food, and 93% considered the reef fish to be similarly important. These findings validate the significance of fishing in building and maintaining social networks, perpetuating fishing traditions, and providing fish to local communities as a source of food security.

Fishing in the CNMI is a social activity; only 3% of fishermen reported to fish alone, but 70% reported that their boat is used without them on occasion. In addition, the majority of fishermen (57%) agreed that, as a fisherman, they are respected by the greater community. Nearly a third of respondents were neutral (27%) regarding this sentiment, while some were hesitant to express an opinion or simply did not know (13%). The study found that very few fishers (3%) felt that they were not respected by the community.

The designation of the Marianas Trench Marine National Monument (the Monument) in 2009 has resulted in concerns about loss of fishing access (Richmond and Kotowicz, 2015; Kotowicz and Richmond, 2013; Kotowicz and Allen, 2015; and Kotowicz *et al.*, 2017). Despite long distance, high cost, and inconvenience, travel to the areas now protected by the Monument were rare but culturally significant events, and fishing was an essential component. While CNMI residents generally supported designation of the monument, awareness was low regarding specific impacts (Kotowicz *et al.*, 2017). In addition, fishing households showed higher awareness of the Monument, but were less likely to strongly support it.

Overall, the CNMI small boat fisheries are a mix of subsistence, cultural, recreational, and quasi-commercial fishermen whose fishing behaviors provide evidence of the importance of fishing to the people of the CNMI. For nearly all fishery participants, the social and cultural motivations for fishing far outweigh economic prospects. Nearly all fishermen supplement their income with other jobs and are predominantly subsistence fishermen.


### **2.3.2.3 CNMI Bottomfish**

Bottomfish was one of the gear types included in the 2011 Small Boat Survey (Hospital and Beavers, 2014). Overall fisher demographics and catch disposition were summarized in the previous section. Approximately 68% of respondents reported fishing for deepwater bottomfish and 65% for shallow-water bottomfish; additionally, 41% identified deepwater bottomfish as their primary target, and 49% identified shallow-water bottomfish as their primary target. Approximately 37% of trips included some form of bottomfishing. In general, deepwater bottomfishing appeared to be associated with more commercially-motivated fishermen. Fishers who primarily targeted bottomfish sold over half of their catch (52%) to friends, neighbors, and co-workers. Some self-identified primarily as subsistence fishers (58% selected this category) and recreational expense fishers (41%), although respondents spanned all response categories (full-time commercial, part-time commercial, recreational expense, purely recreational, subsistence, and cultural). Nearly half identified multiple motivations (49%).


### 2.3.2.3.1 Commercial Participation, Landings, Revenue, Prices

This section will describe trends in commercial pounds sold, revenues and prices, for the CNMI bottomfish fishery. Figure 11 presents the trends of commercial pounds sold and revenues of bottomfish fishery (BMUS only) during 2009-2018 and Figure 12 presents the trend of fish price of bottomfish sold for the same period. Supporting data for Figure 11 and Figure 12 are shown in Table 59. The table also includes and the percentage of pounds sold to the total pounds landed of the bottomfish fishery. Both nominal and adjusted values are included. As shown in Figure 11, the commercial landings of CNMI bottomfish were quite stable except 2015 and 2018. Fish price was in an increasing trend from up to 2015. Price dropped in 2016, but increased in the recent two years.

Please notice that the data for pounds caught and pounds sold are collected by two different data collection methods. The data of pounds sold were collected through “Commercial Sales Receipt Books” Program, while the data of pounds caught were collected through “Boat-based Creel Survey” and “Shore-based Creel Survey” ([https://www.pifsc.noaa.gov/wpacfin/cnmi/Pages/cnmi\\_coll\\_3.php](https://www.pifsc.noaa.gov/wpacfin/cnmi/Pages/cnmi_coll_3.php)). Both data series are generated from an expansion algorithm built on a non-census data collection program respectively, and the survey coverage rates of two data collection methods may change independently in individual years. Therefore, the two time series may not move coherently to each other. For example, the low percentage of pounds sold compared to pounds caught could be due to the low coverage of dealer participations in the Commercial Receipt Books Program, or vice versa. In 2014, the ratio of pounds sold to pound caught of BMUS was particularly high, 229%, while the total pounds sold in 2014 were similar to the figures in the previous years and the estimated pounds caught was particularly low for 2014. It seems that the data quality for the pounds landed estimation for 2014 had some issues.



**Figure 11. The commercial landings and revenues of BMUS, for the CNMI bottomfish fishery, 2009-2018 (Adjusted to 2018 dollars)**



**Figure 12. The prices of BMUS for the CNMI bottomfish fishery, 2009-2018**

**Table 59. Commercial landings and revenue information of CNMI bottomfish fishery, 2009-2018\***

| Year | Estimated pounds caught (lb) | Estimated pounds sold (lb) | Estimated revenue (\$) | Estimated revenue (\$ adjusted) | % of pounds sold | Fish price (\$) | Fish price (\$ adjusted) | CPI adjustor |
|------|------------------------------|----------------------------|------------------------|---------------------------------|------------------|-----------------|--------------------------|--------------|
| 2009 | 69,587                       | 20,419                     | 63,822                 | 67,205                          | 29%              | 3.13            | 3.29                     | 1.05         |
| 2010 | 58,610                       | 14,730                     | 50,947                 | 50,845                          | 25%              | 3.46            | 3.45                     | 1.00         |
| 2011 | 29,600                       | 16,931                     | 52,377                 | 51,120                          | 57%              | 3.09            | 3.02                     | 0.98         |
| 2012 | 136,853                      | 11,747                     | 45,636                 | 44,039                          | 9%               | 3.89            | 3.75                     | 0.97         |
| 2013 | 24,065                       | 17,769                     | 71,323                 | 70,610                          | 74%              | 4.01            | 3.97                     | 0.99         |
| 2014 | 8,450                        | 19,333                     | 93,474                 | 91,511                          | 229%             | 4.83            | 4.73                     | 0.98         |
| 2015 | 11,121                       | 4,196                      | 22,177                 | 22,643                          | 38%              | 5.28            | 5.4                      | 1.02         |
| 2016 | 49,367                       | 18,726                     | 76,938                 | 76,938                          | 38%              | 4.11            | 4.11                     | 1            |
| 2017 | 46,290                       | 9,623                      | 45,706                 | 45,706                          | 21%              | 4.75            | 4.75                     | 1            |
| 2018 | 858                          | 4,612                      | 21,994                 | 21,994                          | 538%             | 4.77            | 4.77                     | 1            |

Data source: PIFSC WPacFIN.


### 2.3.2.3.2 Costs of Fishing

Since 2009, PIFSC economists have maintained a continuous economic data collection program for small boat fisheries in Saipan through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN) (Chan and Pan 2019). The economic data collection program gathers fishing expenditure data for boat-based reef fish, bottomfish, and pelagic fishing trips on an ongoing basis. Data for fishing trip expenses include; gallons of fuel used, price per gallon of



fuel, cost of ice used, cost of bait & chum used, cost of fishing gear lost, and the engine type of the boat. These economic data are collected from same subset of fishing trips as the boat-based creel survey carried out by the local fisheries management agencies and WPacFIN. These data are currently under PIFSC editorial review and future versions of this report will include a time-series of Saipan boat-based trip costs by target species and/or gear. Metadata for these data are available online (PIFSC Socioeconomics Program, 2016). Island-specific (Saipan, Tinian, and Rota) trip cost estimates for bottomfish fishing trips are available only for 2011 in Hospital and Beavers (2014). Other relevant cost information in Hospital and Beavers (2014) include estimates of annual fishing expenditures (fixed costs) and levels of investment in the fishery.

The trip cost data presented in this section were collected through the continuous economic data collection program on Saipan through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN). Figure 13 shows the trend of average trip costs for CNMI bottomfish trips during 2009–2017 (adjusted to 2017 dollars), as 2018 data were not available due to limited observations for trip costs. Supporting data of Figure 13 are presented in Table 60. The trip costs seem to have substantial interannual variability. The average costs for a bottomfish trip was \$38 in 2017.



**Figure 13. Average costs for CNMI bottomfish trips, 2009–2017 (adjusted to 2017 dollars)**

**Table 60. Average trip costs for CNMI bottomfish trips, 2009–2017\*, adjusted to 2017 dollars**

| Year | Total trip costs (\$) | Total trip costs (\$ adjusted) | Fuel cost (\$) | Fuel cost (\$ adjusted) | Ice cost (\$) | Ice cost (\$ adjusted) | Gear lost cost (\$) | Gear lost cost (\$ adjusted) | Bait & chum cost (\$) | Bait & chum cost (\$ adjusted) | CPI adjustor |
|------|-----------------------|--------------------------------|----------------|-------------------------|---------------|------------------------|---------------------|------------------------------|-----------------------|--------------------------------|--------------|
| 2009 | 37                    | 39                             | 30             | 32                      | 4             | 4                      | 0.13                | 0.1                          | 3                     | 3                              | 1.053        |
| 2010 | 20                    | 20                             | 17             | 17                      | 2             | 2                      | 0.00                | 0.0                          | 0                     | 0                              | 0.998        |
| 2011 | 19                    | 19                             | 16             | 16                      | 2             | 2                      | 0.10                | 0.1                          | 1                     | 1                              | 0.976        |
| 2012 | 61                    | 59                             | 51             | 49                      | 8             | 8                      | 0.00                | 0.0                          | 2                     | 2                              | 0.965        |
| 2013 | 63                    | 62                             | 57             | 56                      | 3             | 3                      | 0.59                | 0.6                          | 2                     | 2                              | 0.990        |
| 2014 | 22                    | 22                             | 20             | 19                      | 3             | 3                      | 0.00                | 0.0                          | 0                     | 0                              | 0.979        |
| 2015 | 35                    | 36                             | 32             | 33                      | 3             | 3                      | 0.00                | 0.0                          | 0                     | 0                              | 1.021        |
| 2016 | 65                    | 65                             | 57             | 57                      | 8             | 8                      | 0.00                | 0.0                          | 0                     | 0                              | 1.000        |
| 2017 | 38                    | 38                             | 32             | 32                      | 5             | 5                      | 0.00                | 0.0                          | 1                     | 1                              | 1.000        |

\*No cost update for 2018 due to no data collected in 2018.

Data source: PIFSC Continuous Cost Data Collection Program (Chan and Pan 2019).

#### 2.3.2.4 CNMI Reef Fish

Coral reef fish were also included in the 2011 small boat survey (Hospital and Beavers, 2014). Unsurprisingly, fishermen targeting reef fish, on average, were slightly younger than others, likely due to the physical requirements of reef fishing. Approximately 54% of respondents reported atulai fishing, 50% reported spearfishing, and 12% reported net fishing. Atulai was identified as the primary choice by 46% of fishermen, while 38% indicated spearfishing was preferable, and 14% net fishing as their primary gear type. Fishers who primarily targeted reef fish sold almost half of their catch (45%) to friends, neighbors, and co-workers. They self-identified primarily as subsistence fishers (44%) and cultural fishers (38%), although respondents spanned all response categories (full-time commercial, part-time commercial, recreational expense, purely recreational, subsistence, and cultural). Over one-third identified multiple motivations (38%).

In addition to playing an important role in subsistence and cultural fishing, coral reef ecosystems of Saipan only have been estimated at a value of \$61 million, 70% of which is accounted for by tourism (Grace McCaskey, 2014).


##### 2.3.2.4.1 Commercial Participation, Landings, Revenue, Prices

This section will describe trends in commercial pounds sold, revenues and prices, for the CNMI coral reef fish fishery. Figure 14 presents the trends of commercial pounds sold and revenues of coral reef fish fishery during 2009-2018 and Figure 15 presents the trend of fish price of coral reef fish sold during 2009-2018. As showing in Figure 14, the coral reef fish fishery (pounds sold and revenue) in CNMI in a declining trend, except in 2016. Fish price was pretty flat around \$2.76 in nominal value (Figure 15). Supporting data for Figure 14 and Figure 15 are shown in Table 61.


Table 61 also includes % of pounds sold to the total pounds caught of the coral reef fish fishery. Both nominal and adjusted values are included in the table. Compared to the pounds sold to

pounds caught, it seems that the reported pounds sold were higher than pounds caught in most of the years, in average across years, pounds sold were 40% greater than pounds landed. Reef fish price was steady in general during the period, but slightly decreased in recent two years.

Please notice that the data for pounds caught and pounds sold are collected by two different data collection methods. The data of pounds sold were collected through “Commercial Sales Receipt Books” Program, while the data of pounds caught were collected through “Boat-based Creel Survey” and “Shore-based Creel Survey” ([https://www.pifsc.noaa.gov/wpacfin/cnmi/Pages/cnmi\\_coll\\_3.php](https://www.pifsc.noaa.gov/wpacfin/cnmi/Pages/cnmi_coll_3.php)). Both data series are generated from an expansion algorithm built on a non-census data collection program respectively, and the survey coverage rates of two data collection methods may change independently in individual years. Therefore, the two time series may not move coherently to each other. For example, the low percentage of pounds sold compared to pounds caught could be due to the low coverage of dealer participations in the Commercial Receipt Books Program, or vice versa.



**Figure 14. The pounds sold and revenues for the CNMI reef fish fishery, 2009-2018 (adjusted to 2018 dollars)**



Note: Missing 2018 and 2017 CNMI inflation rate, assuming 0%

**Figure 15. The fish prices of CREMUS for the CNMI reef fishery, 2009-2018**

**Table 61. Commercial landings and revenue information of CNMI coral reef fishery, 2009-2018**

| Year | Estimated pounds caught (lb) | Estimated pounds sold (lb) | Estimated revenue (\$) | Estimated revenue (\$ adjusted) | % of pounds sold | Fish price (\$) | Fish price (\$ adjusted) | CPI adjustor |
|------|------------------------------|----------------------------|------------------------|---------------------------------|------------------|-----------------|--------------------------|--------------|
| 2009 | 107,543                      | 125,403                    | 330,957                | 348,498                         | 1.166073         | 2.64            | 2.78                     | 1.05         |
| 2010 | 72929                        | 89566                      | 233020                 | 232554                          | 1.228126         | 2.6             | 2.6                      | 1.00         |
| 2011 | 79269                        | 95080                      | 248352                 | 242392                          | 1.19946          | 2.61            | 2.55                     | 0.98         |
| 2012 | 68476                        | 68160                      | 181411                 | 175062                          | 0.995385         | 2.66            | 2.57                     | 0.97         |
| 2013 | 94541                        | 77117                      | 220965                 | 218755                          | 0.815699         | 2.87            | 2.84                     | 0.99         |
| 2014 | 24022                        | 74988                      | 246724                 | 241543                          | 3.121638         | 3.29            | 3.22                     | 0.98         |
| 2015 | 30882                        | 42343                      | 121056                 | 123598                          | 1.371122         | 2.86            | 2.92                     | 1.02         |
| 2016 | 58525                        | 75949                      | 235358                 | 235358                          | 1.297719         | 3.1             | 3.1                      | 1            |
| 2017 | 22735                        | 37573                      | 109610                 | 109610                          | 1.65265          | 2.92            | 2.92                     | 1            |
| 2018 | 26126                        | 29006                      | 82547                  | 82547                           | 1.110235         | 2.85            | 2.85                     | 1            |


Data source: PIFSC WPacFIN

#### 2.3.2.4.2 Costs of Fishing

Since 2009, the PIFSC Socioeconomics Program has maintained a continuous economic data collection program small boat fisheries in Saipan through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN) (Chan and Pan 2019). The economic data

collection program gathers fishing expenditure data for boat-based reef fish, bottomfish, and pelagic fishing trips on an ongoing basis. Data for fishing trip expenses include: gallons of fuel used, price per gallon of fuel, cost of ice used, cost of bait and chum used, cost of fishing gear lost, and the engine type of the boat. These economic data are collected from same subset of fishing trips as the boat-based creel survey carried out by the local fisheries management agencies and WPacFIN. These data are currently under PIFSC editorial review and future versions of this report will include time-series of Saipan boat-based trip costs by target species and/or gear. Meta-data for these time series are available online (PIFSC Socioeconomics Program, 2016). Island-specific trip cost estimates for reef fishing trips are available only in 2011 from Hospital and Beavers (2014). Other relevant cost information from Hospital and Beavers (2014) includes estimates of annual fishing expenditures (fixed costs) and levels of investment in the fishery.

The trip cost data presented in this section were collected through the continuous economic data collection program in collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN). Figure 16 shows the trend of average trip costs for CNMI coral reef fish fishing trips during 2009–2017 (adjusted to 2017 dollars), as the 2018 cost data were not available due to limited observations. Only four years are available with data during the 10 year period due to limited observations in fishing cost data collections. Supporting data for Figure 16 are listed in Table 62.



\*Trip cost data are not presented since the observations for those years were less than 3.

**Figure 16. Average costs for CNMI spearfishing trips, 2009–2017 (adjusted to 2017 dollars)**

**Table 62. Average costs for CNMI bottomfish trips, 2009–2017 (adjusted to 2017 dollars)**

| Year  | Total trip costs (\$) | Total trip costs (\$ (adjusted)) | Fuel cost (\$) | Fuel cost (\$ (adjusted)) | Ice cost (\$) | Ice cost (\$ (adjusted)) | Gear lost cost (\$) | Gear lost cost (\$ (adjusted)) | Bait & chum cost (\$) | Bait & chum cost (\$ (adjusted)) | CPI adjustor |
|-------|-----------------------|----------------------------------|----------------|---------------------------|---------------|--------------------------|---------------------|--------------------------------|-----------------------|----------------------------------|--------------|
| 2009  | 14                    | 15                               | 12             | 12                        | 2             | 2                        | 0                   | 0                              | 0                     | 0                                | 1.053        |
| 2010  | 0                     | 0                                | -              | -                         | -             | -                        | -                   | -                              | -                     | -                                | 0.998        |
| 2011  | 9                     | 9                                | 7              | 7                         | 2             | 2                        | 0                   | 0                              | 0                     | 0                                | 0.976        |
| 2012* | 10                    | 9                                | 8              | 8                         | 1             | 1                        | 0                   | 0                              | 0                     | 0                                | 0.965        |
| 2013  | 8                     | 7                                | 6              | 6                         | 1             | 1                        | 0                   | 0                              | 0                     | 0                                | 0.990        |
| 2014  | -                     | -                                |                |                           |               |                          |                     |                                |                       |                                  | 0.979        |
| 2015  | -                     | -                                |                |                           |               |                          |                     |                                |                       |                                  | 1.021        |
| 2016  | -                     | -                                |                |                           |               |                          |                     |                                |                       |                                  | 1            |
| 2017  | -                     | -                                |                |                           |               |                          |                     |                                |                       |                                  | 1            |
| 2018  | -                     | -                                |                |                           |               |                          |                     |                                |                       |                                  | 1            |

\*Trip cost data are not presented since the observations for those years were less than 3.

<sup>1</sup>Data source: PIFSC Continuous Cost Data Collection Program (Chan and Pan 2019).

### 2.3.2.5 CNMI Crustaceans

*There are currently no socioeconomics data specific to the crustacean fishery. Future reports will include new information as resources allow.*

### 2.3.2.6 CNMI Precious Corals

*There are currently no socioeconomic data specific to this fishery. Future reports will include new information as resources allow.*

## 2.3.3 Guam

### 2.3.3.1 Introduction

An overview of Guam’s history, culture, geography, and relationship with the U.S. is described in the Fishery Ecosystem Plan for the Mariana Archipelago (Western Pacific Regional Fishery Management Council, 2016b). Guam is the largest and southernmost island of the Mariana Archipelago, and is also the largest and most heavily populated island in Micronesia. Over the past decade, a number of studies have synthesized more details about the role of fishing and marine resources for residents of Guam, as well as information about the people who engage in the fisheries and/or utilize fishery resources.

The ancestors of the indigenous Chamorro first arrived in the Marianas around 3,500 years ago, and were expert fishermen and seafarers, relying on seafood as their principal source of protein (Allen and Bartram, 2008; Grace McCaskey, 2014; Hospital and Beavers, 2012). They fished on the high seas in large sailing canoes (proas) and used numerous methods to catch reef and bottomfish from boats. Similar to other archipelagos in the Western Pacific, fish and marine resources have played a central role in shaping the social, cultural, and economic fabric of Guam that continues today. Chamorro fished for both reef and pelagic species, collected mollusks and other invertebrates, and caught sea turtles.

The occupation of Guam by foreign nations dramatically changed the island's ecosystems, reshaped communities, and disrupted fishing traditions. In the 17<sup>th</sup> and 18<sup>th</sup> centuries, Spanish colonizers destroyed the Chamorro' seagoing canoes, suppressed offshore fishing practices, and relocated populations from their traditional home. Following the Spanish-American War in 1898, the U.S. Navy took control of Guam until it was occupied by Japan from 1941-1944. Guam became a U.S. territory in 1950, and the U.S. military is currently in the process of building up an even greater presence on the island. Throughout this time, fishing has remained an important activity, although by the time Guam became an American territory, the indigenous inhabitants had lost many of their seafaring skills, fishing skills, and even the native names of many of the offshore species. Later immigrants to the islands from East and Southeast Asia also possessed a strong fishing tradition. In 2000, 37% of Guam's population that identified as a single ethnicity were Chamorro, followed by 32% Asian (about 80% of whom were Filipino), 17% other Pacific Islander, 7% white, and 1% black. Despite rapid socioeconomic change, households still reflect the traditional pattern of extended families with multigenerational clustering of relatives, especially in Guam's southern villages. Social occasions such as neighborhood parties, wedding and baptismal parties, wakes and funerals, and especially village fiestas that follow the religious celebrations of village patron saints all require large quantities of fish and other traditional foods, reflecting the role of fish in maintaining social ties and cultural identities. Sometimes fish are also sold to earn money to buy gifts for friends and relatives on important Catholic religious occasions such as novenas, births and christenings, and other holidays.

Since the late 1970s, Guam's most important role in commercial fisheries activity has been as a major regional fish transshipment center and resupply base for domestic and foreign tuna fishing fleets. Services provided include fueling, provisioning, unloading, air and sea transshipment, net and vessel repair, crew repatriation, medical care, and warehousing. Among Guam's advantages as a home port are: well-developed and highly efficient port facilities in Apra Harbor, an availability of relatively low-cost vessel fuel, a well-established marine supply/repair industry, and recreational amenities for crew shore leave. In addition, the territory is exempt from the Nicholson Act, which prohibits foreign ships from landing their catches in U.S. ports. Initially, the majority of vessels calling in Apra Harbor to discharge frozen tuna for transshipment were Japanese purse seine boats and carrier vessels. In the late 1980s, Guam became an important port for Japanese and Taiwanese longline fleets, but port calls have steadily declined and the transshipment volume has declined accordingly. By the early 1990s, an air transshipment operation had also been established on Guam. Fresh tuna was flown into Guam from the Federated States of Micronesia and elsewhere on air cargo planes and out of Guam to the Japanese market on wide-body passenger planes. Further, vessels from Japan and Taiwan also landed directly into Guam, where their fish were packed and transshipped by air to Japan. A second air transshipment operation began in the mid-1990s that was transporting fish to Europe that did not meet Japanese sashimi market standards, but this has since ceased. Moreover, the entire transshipment industry has contracted markedly with only a few operators still making transshipments to Japan. Annual volumes of tuna transshipped of between 2007 and 2011 averaged about 3,400 mt, with a 2012 estimate of 2,222 mt, compared to over 12,000 mt at the peak of operations between 1995 and 2001. As early as 2006, it was noted that the Port of Guam had lost much of its competitive advantage compared to alternative transshipment locations in the western Pacific and elsewhere, a trend that may not be reversible.

Otherwise, commercial fisheries have a relatively minor contribution to Guam's economy; the social and cultural importance of fisheries in Guam dwarfs their commercial value. Nearly all Guam domestic fishermen hold jobs outside the fishery, with fishing typically supplementing family subsistence. High value is placed on sharing one's fish catch with relatives and friends, and this social obligation extends to part-time and full-time commercial fishermen alike. A survey of Guam households in 2005 found that nearly one-quarter (24%) of fish consumed were caught by the respondent or an immediate family member, and an additional 14% were caught by a friend or extended family member (Allen and Bartram, 2008). However, a little more than half (51%) of the fish consumed were purchased at a store or restaurant, and 9% were purchased at a flea market or from a roadside stand. The same study found that annual seafood consumption in Guam is estimated to be about 60 lbs. per capita, with approximately 43% imported from the U.S.

The westernization of Guam, particularly since World War II, has not only resulted in a transition from a subsistence to wage-based economy, but has also contributed to dramatic changes in eating patterns, including lower seafood consumption. Indeed, recent years have seen steady declines in the market demand for fresh local fish across Guam (Hospital and Beavers, 2012). While some families continue to supplement their diet by fishing and farming, no existing communities are completely dependent on local fishing as a source of food. A household survey conducted in 2016 found that only 29% of respondents participate in fishing (National Coral Reef Monitoring Program, 2016a).

Allen and Bartram (2008) reviewed the history of shoreline and inshore fishing on Guam. They noted that the number of people engaged in shore fishing in the 1970s was surprisingly large, given that about 90% of the food consumed on the island was imported. A study conducted in 1975 found that 65% of households reported some participation in fishing, which was presumably shore-fishing as a result of the low level of boat ownership at the time. Creel surveys conducted by the Guam DAWR indicated that CPUE in Guam's shore-based fisheries for reef fish (pole, spear, cast net, surround net, and gill net) declined sharply in the 1980s and had not recovered by 2008. Offshore (boat-based) catches of reef-associated fish were relatively constant between 1992 and 2008, whereas inshore catches that accounted for the majority of the reef fish harvest during the 1990s comprised a minority of the total harvest by 2008. Much of the traditional harvest targets seasonal runs of juvenile rabbitfish, goatfish, bigeye scad (atulai, *Selar crumenophthalmus*), and jacks (i'e, family Carangidae). A study in 2007 estimated that Guam's coral reef resources were valued at close to \$127 million annually, primarily driven by the island's important tourism industry (Grace McCaskey, 2014). Nearly 1.2 million people visited Guam in 2010, many of them attracted by reef-related activities, such as snorkeling and scuba diving.

As recently as the early 1970s, relatively few people from Guam fished offshore because boats and deep-sea fishing equipment were prohibitively expensive (Allen and Bartram, 2008). During the economic boom from the late-1980s through most of the 1990s, Guam developed a small boat fishery that conducted trolling and bottomfishing mostly within 30 miles of shore.

The Guam Fishermen's Cooperative Association (GFCA) plays an important role in preserving important fishing traditions. It began operations in 1976 and was incorporated in 1977. In 2006, its membership included 164 full- and part-time fishermen from every district on Guam, and it



processed and marketed approximately 80% of the local commercial catch. In addition, it plays a role in fisheries data collection, marine education and training, and fisheries conservation and management. The GFCA strives to provide benefits not just to fishermen but to residents throughout Guam, benefitting the broader Guam community. It utilizes a Hazard Analysis and Critical Control Point (HACCP) system to ensure safe seafood, and tests fish for potential toxins or whenever requested by the Guam Department of Health and Sanitation. It has also become a focal point for community activities, such as the Guam Marianas International Fishing Derby, cooking competitions, the Guam Fishermen's Festival, dissemination of educational materials on marine resources, vessel safety, seafood preparation, public meetings on resource management issues, and communications via radio base to relay information and coordinate rescues. It also has adopted a policy of purchasing local origin products that benefits 40 small businesses on Guam, regularly donates seafood for village functions and charitable activities, and provides assistance to victims of periodic typhoons with emergency supplies of ice and fuel. In addition, the GFCA has become a voice for Guam fishermen in the policy arena to ensure that concerns of fishermen are incorporated into relevant issues, including the military buildup and loss of fishing grounds due to establishment of Marine Preserve Areas.

Fishing in Guam continues to be important not only in contributing to the subsistence needs of the Chamorro and other residents, but also in preserving their histories and identities. Knowledge of how fish are distributed and consumed locally is crucial to understanding the social and cultural significance of fishing on Guam.

#### **2.3.3.2 People who Fish**

Few studies have been conducted on fishing in Guam in general. A household survey conducted in 2012 found that 35% of respondents said that they or someone else in their household was a fisherman (Kotowicz and Allen, 2015). Respondents from fishing households tended to have lower education levels and have a higher rate of unemployment than respondents from non-fishing households.

As described in Allen and Bartram (2008), in 1999, a detailed study of the inshore fishing behaviors and spatial patterns was conducted for the three largest resident fishing cultures on Guam: Chamorro, Micronesian, and Filipino. At that time, Chamorro comprised about 75% of the fishing parties encountered, while Micronesians constituted about 17% and Filipinos about 7%. A number of contemporary reef fishing methods on Guam were observed, including gleaning, hand line, rod and reel, talaya (cast net), tekken (gill net), chenchulu (surround net), and spearfishing. Explicit rules governing permanent marine ownership were not observed, but Chamorro fishermen maintained a strong identification with village and municipal space. This village relationship included the reef during the early part of the 20th century but that has since largely disappeared. Instead, a system of "pliant tenure" (a vestige of traditional marine tenure) was recognized; while any reef area is publicly accessible, fishermen act according to a system of temporary ownership or pliant tenure of reef area. These rules were understood and incorporated by Chamorro and immigrant fishers alike. Respondents voiced concern about the loss of fishing grounds through designation of marine reserves and tourist watercraft activities. They viewed reduced coastal access as threatening the perpetuation of cultural identity and practice by reducing ability to teach and practice traditions such as communal harvests and distribution of the catches, which reinforce family cohesion and communal identity. These

practices have been further jeopardized by the build-up of U.S. military personnel and families in recent years.

In the mid-1980s Guam fisheries were characterized as including (1) a small number of true commercial fishermen, (2) subsistence/recreational fishermen who regularly sell part of their catch, (3) a large number of subsistence fishermen who rarely sell any of their catch, and (4) a substantial number of recreational fishermen. Approximately 60% of catch was non-commercial, with fish sales primarily used to generate revenue to pay for fuel costs. A similar pattern continues in recent years.

In 2011, a survey was conducted of the small boat fleet, which included questions about trolling, bottomfishing, and reef fishing. On average, fishermen responding to the survey were 44 years old and reported to have been boat fishing for an average of 20 years. Respondents were also more educated and more affluent than the general population. The majority of respondents described themselves as Chamorro (72%), followed by white (23%) with relatively small proportions of Filipinos (6%), Micronesians (6%), other ethnicities (5%), and Carolinians (1%) represented. There was considerable evidence of co-ownership and sharing of fishing vessels. In addition, fishermen reported the use of multiple gear types, with pelagic trolling as the most popular gear type followed by shallow-water bottomfish fishing and deepwater bottomfish fishing. Almost all (96%) fishermen reported fishing at a Fish Aggregating Device (FAD) during the past year and on nearly half (53%) of their fishing trips. Fishing for bottomfish and reef fish was highly seasonal compared to pelagics. Whereas over half of the survey respondents (54%) fished all year for pelagics, only 16% fished year-round for bottomfish and reef fish.

Approximately 70% of fishermen reported selling at least a portion of their catch, and 82% could always sell all the fish that they wanted to sell. However, nearly 30% reported that they had not sold any fish in the past year, and nobody reported selling all the fish they caught. Instead, cost recovery was cited as the primary motivation for the sale of fish, with fish sales contributing very little to personal income for the majority of respondents (59%). In fact, 64% of fishermen reporting the sale of fish earned fishing revenues of less than \$1,000, which would not cover overall trip expenditures for a year. Sale of pelagic fish contributes to nearly 67% of fishing income, with 20% from bottomfish revenues and the rest from reef fish.

While respondents sold approximately 24% of their total catch, 29% was consumed at home, while 42% was given away. The remaining catch was either released (2%) or exchanged for goods and services (3%). This diversity of catch disposition extends to fishermen who regularly sell fish, as they still retain approximately 30% of their catch for home consumption and participation in traditional fish-sharing networks and customary exchange. Additionally, 78% consider the pelagic fish they catch to be an important source of food, 79% for bottomfish, and 85% for reef fish. These findings validate the importance of fishing in terms of building and maintaining social and community networks, perpetuating fishing traditions, and providing food security to local communities.

Like with CNMI, fishing on Guam is a social activity. Only 7% of fishermen reported fishing alone, and 45% reported that their boat is used without them on occasion. In addition, 61% reported to be a member of a fishing club, association, or group. The majority of fishermen

(60%) also agreed that as a fisherman, they are respected by the Guam community. Very few felt that they were not respected by the community.

There was also an open-ended portion of the survey that asked for comments. The two most prevalent themes were that of a rising population and rising fuel costs. Many believed that the expanding population would increase the demand for fish and number of fishermen, yet at the same time, others noted that fuel costs and economic considerations could restrict fishing. In addition, there was concern about the designation of Marianas Trench Marine National Monument, especially since respondents felt that the Marine Preserve Areas established in 1997 had already displaced them from their traditional fishing grounds. Military exercises also affected fishing trips. Other studies have also documented concerns about fishing access related to the designation of the Monument (Richmond and Kotowicz, 2015; Kotowicz and Richmond 2013; and Kotowicz and Allen, 2015). Despite long distance, high cost, and inconvenience, travel to the areas now protected by the Monument were rare but culturally significant events of which fishing was an essential component.

Similar to CNMI, Guam's small boat fisheries are a complex mix of subsistence, cultural, recreational, and quasi-commercial fishermen whose fishing behaviors provide evidence of the importance of fishing to the island of the Guam. For nearly all fishery participants, the social and cultural motivations for fishing far outweigh any economic prospects. Nearly all fishermen supplement their income with other jobs and are predominantly subsistence fishermen, selling occasionally to recover trip expenses.

### **2.3.3.3 Guam Bottomfish**

Allen and Bartram (2008) reviewed the history of the bottomfish fishery on Guam, which consists of both shallow- and deep water aspects. They noted that during the 1980s and 1990s, bottomfish fishing was a highly seasonal, small-scale, commercial, subsistence, and recreational fishery. The majority of the participants operated vessels less than 25 ft. long and targeted the shallow-water bottomfish complex because of the lower expenditure and relative ease of fishing close to shore. The commercially-oriented vessels tended to be longer than 25 ft., concentrating effort on the deepwater bottomfish complex. Both deepwater and shallow-water bottomfish are also important target species of the charter fishing fleet, and charter trips accounted for about 15–20% of all Guam bottomfishing trips from 1995 through 2000. In 1998, the charter fleet attracted approximately 3% of visitors to Guam and consisted of a dozen core boats.


Bottomfish was one of the gear types included in the 2011 small boat survey (Hospital and Beavers, 2014). Overall fisher demographics and catch disposition were summarized in the previous section. Approximately 57% of respondents reported fishing for deepwater bottomfish and 59% for shallow-water bottomfish, with 52% identifying deepwater bottomfish as their primary target and 49% identifying shallow-water bottomfish as their primary target. Fishers who primarily targeted bottomfish allocated their catch mainly through the Guam Fisherman's Cooperative Association (55%), or to friends, neighbors, and co-workers (41%). For the most part, they self-identified as recreational expense fishers (40%), cultural fishers (35%), subsistence fishers (35%), purely recreational fishers (30%), though respondents spanned all response categories except full-time commercial (i.e., part-time commercial, recreational expense, purely recreational, subsistence, and cultural). Over half of the respondents identified multiple motivations (54%).

### 2.3.3.3.1 Commercial Participation, Landings, Revenue, Prices


This section describes trends in commercial pounds sold, revenues and prices, for the Guam bottomfish fishery. Figure 17 presents the trends of commercial pounds sold and revenues of bottomfish fishery during 2009-2018 and Figure 18 presents the trend of fish price of bottomfish sold during 2009-2018 (for BMUS only). Supporting data for Figure 17 and Figure 18 are shown in Table 63. Table 63 also includes the percentage of pounds sold to the total pounds caught of the bottomfish fishery. Both nominal and adjusted values are included in the table.

As showing in Figure 17, the bottomfish fishery (commercial landings and revenue) in Guam was in a declining trend from 2009 to 2016 and it went up in 2017. The commercial landings of 2018 were lower than 2017, but higher than that in the period of 2013-2015. Fish price was pretty steady (Figure 18). Compared to total pounds landed, the commercial landings of BMUS were only small portion. On an average of the recent ten years (2009-2018), the pounds sold were only 15% of total pounds caught. In 2018, the percentage pounds sold was 11% of total pounds caught. Bottomfish price were steady in general, but variations showed in the recent three years.

Please notice that the data for pounds caught and pounds sold are collected by two different data collection methods. The data of pounds sold were collected through Commercial Sales Receipt Books Program, while the data of pounds caught were collected through Boat-based Creel Survey and Shore-based Creel Survey ([https://www.pifsc.noaa.gov/wpacfin/guam/dawr/Pages/gdawr\\_coll\\_3.php](https://www.pifsc.noaa.gov/wpacfin/guam/dawr/Pages/gdawr_coll_3.php)). Both data series are generated from an expansion algorithm built on a non-census data collection program respectively, and the survey coverage rates of two data collection methods may change independently in individual years. Therefore, the two time series may not move coherently to each other. For example, the low percentage of pounds sold compared to pounds caught could be due to the low coverage of dealer participations in the Commercial Receipt Books Program, or vice versa.



**Figure 17. The pounds sold and revenues for the Guam bottomfish fishery, 2009-2018 (adjusted to 2018 dollars)**



**Figure 18. The prices of BMUS for the Guam bottomfish fishery, 2009-2018**

**Table 63. Commercial landings, revenue, and price information of Guam bottomfish fishery, 2009-2018**


| Year | Estimated pounds caught (lb) | Estimated pounds sold (lb) | Estimated revenue (\$) | Estimated revenue (\$ adjusted) | % of pounds sold | Fish price (\$) | Fish price (\$ adjusted) | CPI adjustor |
|------|------------------------------|----------------------------|------------------------|---------------------------------|------------------|-----------------|--------------------------|--------------|
| 2009 | 39,248                       | 10,232                     | 40,284                 | 49,146                          | 26%              | 3.94            | 4.80                     | 1.22         |
| 2010 | 27,044                       | 5,890                      | 25,008                 | 29,760                          | 22%              | 4.25            | 5.05                     | 1.19         |
| 2011 | 59,023                       | 9,941                      | 42,153                 | 48,476                          | 17%              | 4.24            | 4.88                     | 1.15         |
| 2012 | 22,182                       | 5,133                      | 21,740                 | 24,131                          | 23%              | 4.24            | 4.70                     | 1.11         |
| 2013 | 30,869                       | 2,809                      | 11,575                 | 12,848                          | 9%               | 4.12            | 4.57                     | 1.11         |
| 2014 | 24,865                       | 1,650                      | 7,112                  | 7,823                           | 7%               | 4.31            | 4.74                     | 1.10         |
| 2015 | 13,837                       | 804                        | 3,263                  | 3,622                           | 6%               | 4.06            | 4.50                     | 1.11         |
| 2016 | 26,893                       | 1,620                      | 7,485                  | 7,859                           | 6%               | 4.62            | 4.85                     | 1.05         |
| 2017 | 19,143                       | 5,094                      | 20,557                 | 21,174                          | 27%              | 4.04            | 4.16                     | 1.03         |
| 2018 | 32,751                       | 3,557                      | 17,022                 | 17,022                          | 11%              | 4.79            | 4.79                     | 1            |

Data source: PIFSC Continuous Cost Data Collection Program (Chan and Pan 2019).

#### 2.3.3.3.2 Costs of Fishing

Since 2011, PIFSC economists have maintained a continuous economic data collection program for small boat fishing on Guam through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN) (Chan and Pan 2019). The economic data collection gathers fishing expenditure data for boat-based reef fish, bottomfish, and pelagic fishing trips on an ongoing basis. Data for fishing trip expenses include; gallons of fuel used, price per gallon of fuel, cost of ice used, cost of bait & chum used, cost of fishing gear lost, and the engine type of the boat. These economic data are collected from same subset of fishing trips as the boat-based creel survey carried out by the local fisheries management agencies and WPacFIN. Metadata for these data are available online (PIFSC Socioeconomics Program, 2016). Guam trip cost estimates from 2011 for bottomfish fishing trips and other relevant cost information (such as estimates of annual fixed costs) are available in a one-time survey (Hospital and Beavers, 2012).

The time series of trip costs of Guam bottomfish fishing presented in Figure 19 are based on a continuous economic data collection program maintained by the PIFSC Socioeconomics Program through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN). The fishing costs of bottomfish were in a declining trend from 2012-2016, and then went up substantially in 2017. The trip costs of 2018 was lower than 2017, but it was still high compared to the most of the previous years. Supporting data for are presented in Table 64.



\* The number of boats (respondents) was fewer than 3; due to confidentiality concerns, responses are not presented.

**Figure 19. Average trip costs for Guam bottomfish fishing trips from 2009–2018 (adjusted to 2018 dollars)**

**Table 64. Average trip costs for Guam bottomfish fishing trips from 2009–2018**

| Year  | Total trip costs (\$) | Total trip costs (\$ (adjusted)) | Fuel cost (\$) | Fuel cost (\$ (adjusted)) | Ice cost (\$) | Ice cost (\$ (adjusted)) | Gear lost cost (\$) | Gear lost cost (\$ (adjusted)) | Bait & chum cost (\$) | Bait & chum cost (\$ (adjusted)) | CPI adjustor |
|-------|-----------------------|----------------------------------|----------------|---------------------------|---------------|--------------------------|---------------------|--------------------------------|-----------------------|----------------------------------|--------------|
| 2011* | -                     | -                                | -              | -                         | -             | -                        | -                   | -                              | -                     | -                                | 1.149        |
| 2012  | 70                    | 78                               | 33             | 37                        | 11            | 13                       | 18                  | 20                             | 8                     | 8                                | 1.114        |
| 2013  | 61                    | 68                               | 34             | 38                        | 6             | 7                        | 9                   | 10                             | 12                    | 13                               | 1.114        |
| 2014  | 51                    | 56                               | 24             | 27                        | 7             | 8                        | 9                   | 10                             | 10                    | 11                               | 1.105        |
| 2015  | 40                    | 45                               | 17             | 18                        | 6             | 7                        | 7                   | 8                              | 10                    | 11                               | 1.115        |
| 2016  | 36                    | 38                               | 15             | 16                        | 6             | 7                        | 8                   | 9                              | 6                     | 6                                | 1.052        |
| 2017  | 72                    | 74                               | 35             | 36                        | 12            | 12                       | 7                   | 7                              | 18                    | 19                               | 1.026        |
| 2018  | 57                    | 57                               | 27             | 27                        | 12            | 12                       | 10                  | 10                             | 8                     | 8                                | 1            |

\* The number of boats (respondents) was fewer than 3; due to confidentiality concerns, responses are not presented. Data source: PIFSC Continuous Cost Data Collection Program (Chan and Pan 2019).

#### 2.3.3.4 Guam Reef Fish

Coral reef fish were also included in the 2011 small boat survey (Hospital and Beavers, 2014). Approximately 33% of respondents reported atulai fishing, 32% spearfishing, and 8% net fishing. Atulai was identified as the primary target by 31%, 20% indicated spearfishing, and 4% indicated net fishing as their primary gear type. Fishers who primarily targeted reef fish sold their catch mainly through the Guam Fisherman’s Cooperative Association (37%) or to friends, neighbors, and co-workers (51%). For the most part, respondents self-identified as subsistence fishers (46%), purely recreational fishers (46%), cultural fishers (38.5%), and recreational

expense fishers (31%) although respondents spanned all response categories except full-time commercial (i.e., part-time commercial, recreational expense, purely recreational, subsistence, and cultural). Over half of respondents identified multiple motivations (54%).


#### **2.3.3.4.1 Commercial Participation, Landings, Revenue, Prices**

This section describes trends in commercial pounds sold, revenues and prices, for the Guam coral reef fish fishery. Figure 20 presents the trends of commercial landings (pounds sold) and revenues of coral reef fish fishery during 2009-2018 and Figure 21 presents the trend of fish price of coral reef fish sold during 2009-2018. Supporting data for Figure 20 and Figure 21 are shown in Table 65. Table 65 also includes the percentage of pounds sold to the total pounds caught of the CREMUS. Both nominal and adjusted values are also included in the table.


As showing in Figure 20, the coral reef fish fishery (pounds sold) and revenue in Guam was in a declining trend except a soar in both commercial landings and revenue appeared in 2017. The pounds sold in 2018 were lower 2017. On an average of the ten years period (2009-2018), the total commercial landings were 61% of total pounds caught. In 2018, the percentage pounds sold was 55% of total pounds landed. Fish price had been steady up to 2015, but it was lower than the average in recent three years (2016, 2017, and 2018), as Figure 21 shows.

Please notice that the data for pounds caught and pounds sold are collected by two different data collection methods. The data of pounds sold were collected through Commercial Sales Receipt Books Program, while the data of pounds caught were collected through Boat-based Creel Survey and Shore-based Creel Survey ([https://www.pifsc.noaa.gov/wpacfin/guam/dawr/Pages/gdawr\\_coll\\_3.php](https://www.pifsc.noaa.gov/wpacfin/guam/dawr/Pages/gdawr_coll_3.php)). Both data series are generated from an expansion algorithm built on a non-census data collection program respectively, and the survey coverage rates of two data collection methods may change independently in individual years. Therefore, the two time series may not move coherently to each other. For example, the low percentage of pounds sold compared to pounds caught could be due to the low coverage of dealer participations in the Commercial Receipt Books Program, or vice versa.





**Figure 20. The commercial landings and revenues, for the Guam reef fish fishery, 2009-2018**



**Figure 21. The prices of CREMUS for the Guam reef fish fishery, 2009-2018**


**Table 65. Commercial landings and revenue information of Guam coral reef fish fishery, 2009-2018**

| Year | Estimated pounds caught (lb) | Estimated pounds sold (lb) | Estimated revenue (\$) | Estimated revenue (\$ adjusted) | % of pounds sold | Fish price (\$) | Fish price (\$ adjusted) | CPI adjustor |
|------|------------------------------|----------------------------|------------------------|---------------------------------|------------------|-----------------|--------------------------|--------------|
| 2009 | 538,372                      | 179,262                    | 547,105                | 667,468                         | 33%              | 3.05            | 3.72                     | 1.22         |
| 2010 | 156,573                      | 164,367                    | 501,950                | 597,320                         | 105%             | 3.05            | 3.63                     | 1.19         |
| 2011 | 338,417                      | 140,075                    | 424,802                | 488,522                         | 41%              | 3.03            | 3.49                     | 1.15         |
| 2012 | 237,127                      | 96,349                     | 297,952                | 330,727                         | 41%              | 3.09            | 3.43                     | 1.11         |
| 2013 | 268,885                      | 72,008                     | 227,451                | 252,471                         | 27%              | 3.16            | 3.51                     | 1.11         |
| 2014 | 190,951                      | 87,801                     | 283,255                | 311,580                         | 46%              | 3.23            | 3.55                     | 1.10         |
| 2015 | 203,241                      | 58,762                     | 193,723                | 215,033                         | 29%              | 3.30            | 3.66                     | 1.11         |
| 2016 | 154,873                      | 73,252                     | 214,653                | 225,386                         | 47%              | 2.93            | 3.08                     | 1.05         |
| 2017 | 147,475                      | 273,372                    | 715,623                | 737,092                         | 185%             | 2.62            | 2.70                     | 1.03         |
| 2018 | 243,108                      | 133,941                    | 392,548                | 392,548                         | 55%              | 2.93            | 2.93                     | 1            |

Data source: PIFSC WPacFIN

#### 2.3.3.4.2 Costs of Fishing

The trip costs presented in Figure 22 are based on a continuous economic data collection program for small boat fisheries in Guam maintained by the PIFSC economists through collaboration with the PIFSC Western Pacific Fisheries Information Network (WPacFIN) (Chan and Pan 2019). Due to limited observations, only four years’ data were presented in the charts. The fishing costs of coral reef fish fishing trips were in an increasing trend since 2014 to 2017. Supporting data for Figure 22 are presented in Table 66.



\* The number of boats (respondents) was fewer than 3; due to confidentiality concerns, responses are not presented.

**Figure 22. Average trip costs for Guam spear/snorkel trips from 2011–2018 (adjusted to 2018 dollars)**

**Table 66. Average trip costs for Guam spear/snorkel fish trips from 2011–2018 (adjusted to 2018 dollars)**

| Year  | Total trip costs (\$) | Total trip costs (\$)<br>(adjusted) | Fuel cost (\$) | Fuel cost (\$)<br>(adjusted) | Ice cost (\$) | Ice cost (\$)<br>(adjusted) | Gear losted cost (\$) | Gear losted cost (\$)<br>(adjusted) | Bait & chum cost (\$) | Bait & chum cost (\$)<br>(adjusted) | CPI adjustor |
|-------|-----------------------|-------------------------------------|----------------|------------------------------|---------------|-----------------------------|-----------------------|-------------------------------------|-----------------------|-------------------------------------|--------------|
| 2011* | -                     | -                                   |                |                              |               |                             |                       |                                     |                       |                                     | 1.149        |
| 2012* | -                     | -                                   |                |                              |               |                             |                       |                                     |                       |                                     | 1.114        |
| 2013* | -                     | -                                   |                |                              |               |                             |                       |                                     |                       |                                     | 1.114        |
| 2014  | 28                    | 31                                  | 18             | 20                           | 10            | 11                          | 0                     | 0                                   | 0                     | 0                                   | 1.105        |
| 2015  | 33                    | 37                                  | 26             | 29                           | 7             | 7                           | 0                     | 0                                   | 0                     | 0                                   | 1.115        |
| 2016  | 28                    | 30                                  | 21             | 22                           | 7             | 7                           | 0                     | 0                                   | 0                     | 0                                   | 1.052        |
| 2017  | 45                    | 46                                  | 26             | 26                           | 12            | 12                          | 7                     | 7                                   | 0                     | 0                                   | 1.026        |
| 2018  | -                     | -                                   |                |                              |               |                             |                       |                                     |                       |                                     | 1            |

\* The number of boats (respondents) was fewer than 3; due to confidentiality concerns, responses are not presented. Data source: PIFSC Continuous Cost Data Collection Program (Chan and Pan 2019).

### 2.3.3.5 Guam Crustaceans

*There are currently no socioeconomic data specific to this fishery. Future reports will include new information as resources allow.*

### 2.3.3.6 Guam Precious Corals

*There are currently no socioeconomic data specific to this fishery. Future reports will include new information as resources allow.*

## 2.3.4 Ongoing Research and Information Collection

Each year, the PIFSC reports on the status of economic data collections for select regional commercial fisheries. This supports a national economic data monitoring effort known as the Commercial Fishing Economic Assessment Index (CFEAI). Details on the CFEAI and access to data from other regions is available at: <https://www.st.nmfs.noaa.gov/data-and-tools/CFEAI-RFEAI/>

The table below represents the most recent data available for CFEAI metrics for select regional commercial fisheries for 2018. Entries for Marianas insular fisheries are bolded in red. These values represent the most recent year of data for key economic data monitoring parameters (fishing revenues, operating costs, and fixed costs). The assessment column indicates the most recent publication year for specific economic assessments (returns above operating cost, profit), where available.

PIFSC completed a cost-earnings survey of small boat fisheries in Guam and the CNMI during 2018-2019, to serve as an update to the previous 2011 cost-earnings survey (Hospital and Beavers, 2012; 2014). This 2018 survey collected data on fishing revenues, operating costs, and fixed costs, as well as numerous elements related to fishing behavior, market participation, and fishery demographics.

**Table 67. Pacific Islands Region 2018 Commercial Fishing Economic Assessment Index**

|                            | 2018 Projected CFEAI                         |   |   |  |  |
|----------------------------|--|---|---|--|--|
|                            | 2018 Reporting Year (e.g. 1/2018-12/2018)    |   |   |  |  |
|                            | Data   |   |   | Assessment   |  |
| Pacific Islands Fisheries  | Anticipated Fishing Revenue Most Recent Year | Anticipated Operating Cost Most Recent Year | Anticipated Fixed Cost Most Recent Year | Anticipated Returns Above Operating Costs (Quasi Rent) Assessment Most Recent Year | Anticipated Profit Assessment Most Recent Year |
| HI Longline                | 2018   | 2018  | 2013                                    | 2018   | 2016   |
| ASam Longline              | 2018   | 2018  | 2017                                    | 2018   | 2016   |
| HI Offshore Handline       | 2018   | 2014  | 2014                                    | 2018   | 2018   |
| HI Small Boat (pelagic)    | 2018   | 2014  | 2014                                    | 2017   | 2017   |
| HI Small Boat (bottomfish) | 2018   | 2014  | 2014                                    | 2017   | 2017   |
| HI Small Boat (reef)       | 2018   | 2014  | 2014                                    | 2017   | 2017   |
| <b>Guam Small boat</b>     | <b>2018</b>                                  | <b>2018</b>                                 | <b>2018</b>                             | <b>2018</b>  |  |
| <b>CNMI Small boat</b>     | <b>2018</b>                                  | <b>2018</b>                                 | <b>2018</b>                             | <b>2018</b>  |  |
| ASam Small boat            | 2018   | 2018  | 2018                                    | 2018   |  |

**Table 68. Pacific Islands Region 2019 Commercial Fishing Economic Assessment Index**

|                            | 2019 Projected CFEAI                         |   |   |  |  |
|----------------------------|--|---|---|--|--|
|                            | 2019 Reporting Year (e.g. 1/2019-12/2019)    |   |   |  |  |
|                            | Data   |   |   | Assessment   |  |
| Pacific Islands Fisheries  | Anticipated Fishing Revenue Most Recent Year | Anticipated Operating Cost Most Recent Year | Anticipated Fixed Cost Most Recent Year | Anticipated Returns Above Operating Costs (Quasi Rent) Assessment Most Recent Year | Anticipated Profit Assessment Most Recent Year |
| HI Longline                | 2019   | 2019  | 2019                                    | 2019   | 2016   |
| ASam Longline              | 2019   | 2019  | 2017                                    | 2019   | 2019   |
| HI Offshore Handline       | 2019   | 2014  | 2014                                    | 2018   | 2018   |
| HI Small Boat (pelagic)    | 2019   | 2014  | 2014                                    | 2017   | 2017   |
| HI Small Boat (bottomfish) | 2019   | 2014  | 2014                                    | 2017   | 2017   |
| HI Small Boat (reef)       | 2019   | 2014  | 2014                                    | 2017   | 2017   |
| <b>Guam Small boat</b>     | <b>2019</b>                                  | <b>2019</b>                                 | <b>2018</b>                             | <b>2019</b>  | <b>2020</b>                                    |
| <b>CNMI Small boat</b>     | <b>2019</b>                                  | <b>2019</b>                                 | <b>2018</b>                             | <b>2019</b>  | <b>2020</b>                                    |
| ASam Small boat            | 2019   | 2019  | 2018                                    | 2019   |  |

PIFSC also generates projections for upcoming fiscal years, and the table above provides the projected CFEAI report for 2019 (*all projected activities and analyses are subject to funding*). Based on early projections PIFSC intends to maintain ongoing economic data collections in the CNMI and Guam for small boat fisheries (Chan and Pan, 2019) during 2019. PIFSC intends for results of the 2018 cost-earnings survey, including a profit assessment, published by 2020.

Community social indicators have been generated for the CNMI and Guam (Kleiber *et al.*, 2018) in accordance with a national project to describe and evaluate community well-being measured through social, economic, and psychological welfare (<https://www.st.nmfs.noaa.gov/humandimensions/social-indicators/index>). However, these indicators rely on Census data, and cannot be updated until 2020 Census data becomes available.

PIFSC scientists will conduct research in the Marianas during 2019-2020 with the goal to engage the Marianas fishing community to better understand the nature of shark interactions and explore mitigation techniques aligned with community needs and values.

### **2.3.5 Relevant PIFSC Economics and Human Dimensions Publications: 2018**

Ayers, A.L., 2018. The commonwealth of the Northern Mariana Islands fishing community profile: 2017 update. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-66, 57 p. doi:10.7289/V5/TM-PIFSC-66.

Finkbeiner, E.M., Micheli, F., Bennett, N.J., Ayers, A.L., Le Cornu, E., and A.N. Doer, 2018. Exploring trade-offs in climate change response in the context of Pacific Island fisheries. *Marine Policy*. 88:359-364. doi:10.1016/j.marpol.2017.09.032.

Kleiber, D., Kotowicz, D., and J. Hospital, 2018. Applying national community social vulnerability indicators to fishing communities in the Pacific Island region. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-65, 63 p. doi:10.7289/V5/TM-PIFSC-65.

### **2.3.6 References**

Allen, S.D. and J.R. Amesbury, 2012. Commonwealth of the Northern Mariana Islands as a fishing community. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFSPIFSC-36, 89 p. [https://www.pifsc.noaa.gov/library/pubs/tech/NOAA\\_Tech\\_Memo\\_PIFSC\\_36.pdf](https://www.pifsc.noaa.gov/library/pubs/tech/NOAA_Tech_Memo_PIFSC_36.pdf).

Allen, S. and P. Bartram, 2008. Guam as a fishing community. Pacific Islands Fisheries Science Center Administrative Report H-08-01, 61 p. [https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_08-01.pdf](https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_08-01.pdf).

Ayers, A.L., 2018. The commonwealth of the Northern Mariana Islands fishing community profile: 2017 update. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-66, 57 p. doi:10.7289/V5/TM-PIFSC-66.

Chan, H.L. and M. Pan, 2019. Tracking economic performance indicators for small boat fisheries in America Samoa, Guam, and the Commonwealth of the Northern Mariana Islands. U.S.

- Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-79, 76 p. <https://doi.org/10.25923/8etp-x479>.
- Grace-McCaskey, C. 2014. Examining the potential of using secondary data to better understand human-reef relationships across the Pacific. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96818-5007. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-14-01, 69 p. [https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_14-01.pdf](https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_14-01.pdf).
- Hospital, J. and C. Beavers, 2012. Economic and social characteristics of Guam's small boat fisheries. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96822-2396. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-12-06, 60 p. + Appendices. [https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_12-06.pdf](https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_12-06.pdf).
- Hospital, J. and C. Beavers, 2014. Economic and Social Characteristics of Small Boat Fishing in the Commonwealth of the Northern Mariana Islands. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96818-5007. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-14-02, 58 p.+ Appendices. [https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_14-02.pdf](https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_14-02.pdf).
- Kotowicz, D.M. and S.D. Allen, 2015. Results of a survey of CNMI and Guam residents on the Marianas Trench Marine National Monument. Pacific Islands Fisheries Science Center, PIFSC Data Report, DR-13-009, 55 p. <https://www.pifsc.noaa.gov/library/pubs/DR-13-009.pdf>.
- Kotowicz, D.M. and L. Richmond, 2013. Traditional Fishing Patterns in the Marianas Trench Marine National Monument. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96822-2396. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-13-05, 54 p. [https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC\\_Admin\\_Rep\\_13-05.pdf](https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_13-05.pdf).
- National Coral Reef Monitoring Program (NCRMP) Socioeconomic Monitoring for Guam – infographic, available at: <https://www.coris.noaa.gov/monitoring/resources/GuamCoral.pdf>.
- PIFSC Socioeconomics Program, 2016. CNMI, American Samoa, and Guam Small Boat Fishery Trip Expenditure (2009 to present). Pacific Islands Fisheries Science Center, <https://inport.nmfs.noaa.gov/inport/item/20627>.
- Polovina, J. and Dreflak, K. (Chairs), Baker, J., Bloom, S., Brooke, S., Chan, V., Ellgen, S., Golden, D., Hospital, J., Van Houtan, K., Kolinski, S., Lumsden, B., Maison, K., Mansker, M., Oliver, T., Spalding, S., and P. Woodworth-Jefcoats, 2016. Pacific Islands Regional Action Plan: NOAA Fisheries climate science strategy. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-59, 33 p.
- Richmond, L., and D.M. Kotowicz, 2015. Equity and access in marine protected areas: The history and future of 'traditional indigenous fishing' in the Marianas Trench Marine National Monument. *Applied Geography*, 59, pp. 117-124.

Rubinstein, D., 2001. A Sociocultural Study of Pelagic Fishing Activities in Guam. Final progress report available from University of Hawaii Joint Institute for Marine and Atmospheric Research, Pelagic Fisheries Research Program. Also available at: <http://www.soest.hawaii.edu/PFRP/pdf/rubinstein01.pdf>.

Western Pacific Regional Fishery Management Council, 2016. Draft Fishery Ecosystem Plan for the Mariana Archipelago. Honolulu, HI. 114 p. + Appendices.

## 2.4 PROTECTED SPECIES

This section of the report summarizes information on protected species interactions in fisheries managed under the Mariana FEP. Protected species covered in this report include sea turtles, seabirds, marine mammals, sharks, and corals. Most of these species are protected under the Endangered Species Act (ESA), the Marine Mammal Protection Act (MMPA), and/or the Migratory Bird Treaty Act (MBTA). A list of protected species found in or near Mariana Archipelago waters and a list of critical habitat designations in the Pacific Ocean are included in Appendix B.

### 2.4.1 Indicators for Monitoring Protected Species Interaction

This report monitors the status of protected species interactions in the Marianas FEP fisheries using proxy indicators such as fishing effort, and changes in gear types as these fisheries do not have observer coverage. Creel surveys and logbook programs are not expected to provide reliable data about protected species interactions. Discussion of protected species interactions is focused on fishing operations in federal waters and associated transit through territorial waters.

### 2.4.2 FEP Conservation Measures

Bottomfish, precious coral, coral reef and crustacean fisheries managed under this FEP have no specific regulations in place to mitigate protected species interactions. Destructive gear such as bottom trawls, bottom gillnets, explosives and poisons are prohibited under this FEP, and these prohibitions benefit protected species by preventing potential interactions with non-selective fishing gear.

#### 2.4.2.1 ESA Consultations

ESA consultations were conducted by NMFS and the U.S. Fish and Wildlife Service (USFWS; for species under their jurisdiction) to ensure ongoing fisheries operations managed under the Marianas FEP are not jeopardizing the continued existence of any listed species or adversely modifying critical habitat. The results of these consultations conducted under section 7 of the ESA are briefly described below and summarized in Table 69.

NMFS concluded in an informal consultation dated April 29, 2015 that all fisheries managed under the Mariana Archipelago FEP are not likely to adversely affect the Indo-West Pacific DPS of scalloped hammerhead shark or ESA-listed reef-building corals.

In January 2018, oceanic whitetip sharks and giant manta rays were listed under the ESA (83 FR 4153 and 83 FR 2916, respectively). NMFS will reinitiate consultation for those two species for the applicable fisheries if NMFS determines that effects are likely. There is no record of giant manta ray incidental catches in Mariana fisheries, and NMFS is reviewing catch data on oceanic white tip shark incidental catch in these fisheries.

**Table 69. Summary of ESA consultations for Mariana Archipelago FEP Fisheries**

| <b>Fishery</b> | <b>Consultation date</b> | <b>Consultation type<sup>a</sup></b> | <b>Outcome<sup>b</sup></b> | <b>Species</b>        |
|----------------|--------------------------|--------------------------------------|----------------------------|-----------------------|
| Bottomfish     | 3/8/2008                 | BiOp                                 | NLAA                       | Loggerhead sea turtle |



| <b>Fishery</b>                     | <b>Consultation date</b> | <b>Consultation type<sup>a</sup></b> | <b>Outcome<sup>b</sup></b> | <b>Species</b>  |
|------------------------------------|--------------------------|--------------------------------------|----------------------------|---|
| (CNMI & Guam)                      | 6/3/2008                 | LOC                                  | NLAA                       | Green sea turtle, olive ridley sea turtle, hawksbill sea turtle, leatherback sea turtle, blue whale, fin whale, humpback whale, sei whale sperm whale   |
| Coral reef ecosystem (CNMI & Guam) | 3/7/2002                 | LOC                                  | NLAA                       | Loggerhead sea turtle, leatherback sea turtle, olive ridley sea turtle, green sea turtle, hawksbill sea turtle, humpback whale, blue whale, fin whale, sei whale, sperm whale   |
|                                    | 5/22/2002                | LOC (USFWS)                          | NLAA                       | Green, hawksbill, leatherback, loggerhead and olive ridley turtles, Newell's shearwater, short-tailed albatross, Laysan duck, Laysan finch, Nihoa finch, Nihoa millerbird, Micronesian megapode, 6 terrestrial plants |
|                                    | 6/3/2008                 | LOC                                  | NLAA                       | Green sea turtle, olive ridley sea turtle, hawksbill sea turtle, leatherback sea turtle, blue whale, fin whale, humpback whale, sei whale, sperm whale  |
| Crustaceans (CNMI & Guam)          | 9/28/2007                | LOC                                  | NLAA                       | Green sea turtle, loggerhead sea turtle, olive ridley sea turtle, hawksbill sea turtle, leatherback sea turtle, blue whale, humpback whale, sei whale, sperm whale  |
| Precious corals (CNMI & Guam)      | 10/4/1978                | BiOp                                 | Does not constitute threat | Sperm whale, leatherback sea turtle   |
| Precious corals (Guam)             | 12/20/2000               | LOC                                  | NLAA                       | Humpback whale, green sea turtle, hawksbill sea turtle  |
| All fisheries                      | 4/29/2015                | BE & LOC                             | NLAA                       | Reef-building corals, scalloped hammerhead shark (Indo-west Pacific DPS)  |

<sup>a</sup> BiOp = Biological Opinion; LOC = Letter of Concurrence; BE = Biological Evaluation

<sup>b</sup> LAA = likely to adversely affect; NLAA = not likely to adversely affect.

#### **2.4.2.1.1 Bottomfish Fishery**

In a Biological Opinion issued on March 8, 2002, NMFS concluded that the ongoing operation of the Western Pacific Region's bottomfish and seamount fisheries was not likely to jeopardize the continued existence of any threatened or endangered species under NMFS's jurisdiction or

destroy or adversely modify any critical habitat. In an informal consultation on June 3, 2008, NMFS concluded that Mariana Archipelago bottomfish fisheries are not likely to adversely affect four sea turtle species (leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales).

#### **2.4.2.1.2 Crustacean Fishery**

In an informal consultation completed on September 28, 2007, NMFS concluded that Mariana Archipelago crustacean fisheries are not likely to adversely affect five sea turtle species (loggerhead, leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales).

#### **2.4.2.1.3 Coral Reef Fishery**

In an informal consultation completed by NMFS on March 7, 2002, NMFS concluded that fishing activities conducted under the Coral Reef Ecosystems FMP are not likely to adversely affect endangered or threatened species or critical habitat under NMFS's jurisdiction. On May 22, 2002, the USFWS concurred with the determination of NMFS that the activities conducted under the Coral Reef Ecosystems FMP are not likely to adversely affect listed species under USFWS's exclusive jurisdiction (i.e., seabirds) and listed species shared with NMFS (i.e., sea turtles).

In an informal consultation completed in June 3, 2008, NMFS concluded that the Mariana Archipelago coral reef fisheries are not likely to adversely affect adversely affects four sea turtle species (leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales).

#### **2.4.2.1.4 Precious Coral Fishery**

In a Biological Opinion issued on October 4, 1978, NMFS concluded that the ongoing operation of the Western Pacific Region's precious coral fisheries was not likely to jeopardize the continued existence of any threatened or endangered species under NMFS's jurisdiction or destroy or adversely modify critical habitat. In an informal consultation completed on December 20, 2000, NMFS concluded that Mariana Archipelago precious coral fisheries are not likely to adversely affect humpback whales, green turtles, or hawksbill turtles.

#### **2.4.2.2 Non-ESA Marine Mammals**

The MMPA requires NMFS to annually publish a List of Fisheries (LOF) that classifies commercial fisheries in one of three categories based on the level of mortality and serious injury of marine mammals associated with that fishery. According to the 2019 LOF (84 FR 22051, May 16, 2019), the Guam and CNMI bottomfish fisheries operating under the Marianas FEP are classified as Category III fisheries (i.e., a remote likelihood of or no known incidental mortality and serious injury of marine mammals).

## 2.4.3 Status of Protected Species Interactions in the Marianas FEP Fisheries

### 2.4.3.1 Bottomfish and Coral Reef Fisheries

There are no observer data available for the Guam and CNMI bottomfish or coral reef fisheries. However based on current ESA consultations, these fisheries are not expected to interact with any ESA-listed species in federal waters around Guam or CNMI. NMFS has also concluded that the Mariana Archipelago bottomfish and coral reef commercial fisheries will not affect marine mammals in any manner not considered or authorized under the Marine Mammal Protection Act.

Based on fishing effort and other characteristics described in Chapter 1 of this report, no notable changes have been observed in the fishery. There is no other information to indicate that impacts to protected species from this fishery have changed in recent years.

### 2.4.3.2 Crustacean and Precious Coral Fisheries

There are currently no crustacean or precious coral fisheries operating in federal waters around Guam or CNMI. However based on current ESA consultations, crustacean fisheries are not expected to interact with any ESA-listed species in federal waters around Guam or CNMI. NMFS has also concluded that the Mariana Archipelago crustacean and precious coral commercial fisheries will not affect marine mammals in any manner not considered or authorized under the Marine Mammal Protection Act.

## 2.4.4 Identification of Emerging Issues

Several ESA-listed species are being evaluated for critical habitat designation (Table 70). If critical habitats are designated, they will be included in this SAFE report and impacts from FEP-managed fisheries will be evaluated under applicable mandates.

**Table 70. Candidate ESA species, and ESA-listed species being evaluated for critical habitat designation**

| Species                |                                | Listing Process                    |  |  | Post-Listing Activity   |               |
|------------------------|--------------------------------|------------------------------------|--|--|---|---------------|
| Common Name            | Scientific Name                | 90-Day Finding                     | 12-Month Finding / Proposed Rule               | Final Rule                                 | Critical Habitat  | Recovery Plan |
| Oceanic whitetip shark | <i>Carcharhinus longimanus</i> | Positive (81 FR 1376, 1/12/2016)   | Positive, threatened (81 FR 96304, 12/29/2016) | Listed as Threatened (83 FR 4153, 1/30/18) | Not determinable because of insufficient data (83 FR 4153, 1/30/18) | TBA           |
| Pacific bluefin tuna   | <i>Thunnus orientalis</i>      | Positive (81 FR 70074, 10/11/2016) | Not warranted (82 FR 37060, 8/8/17)            | N/A  | N/A   | N/A           |

| Species           |  | Listing Process                                 |   |   | Post-Listing Activity   |   |
|-------------------|--|---|---|---|---|---|
| Common Name       | Scientific Name  | 90-Day Finding                                  | 12-Month Finding / Proposed Rule  | Final Rule  | Critical Habitat  | Recovery Plan   |
| Giant manta ray   | <i>Manta birostris</i>   | Positive (81 FR 8874, 2/23/2016)                | Positive, threatened (82 FRN 3694, 1/12/2017)                                 | Listed as Threatened (83 FR 2916, 1/22/18)                          | Not determinable because of insufficient data (83 FR 2916, 1/22/18) | TBA   |
| Reef manta ray    | <i>Manta alfredi</i>   | Positive (81 FR 8874, 2/23/2016)                | Not warranted (82 FRN 3694, 1/12/2017)  | N/A   | N/A   | N/A   |
| Corals            | N/A  | Positive for 82 species (75 FR 6616, 2/10/2010) | Positive for 66 species (77 FR 73219, 12/7/2012)                              | 20 species listed as threatened (79 FR 53851, 9/10/2014)            | In development, proposal expected TBA                               | In development, expected TBA, interim recovery outline in place |
| Cauliflower coral | <i>Pocillopora meandrina</i>   | Positive (83 FR 47592, 9/20/2018)               | TBA (status review ongoing)   | TBA   | N/A   | N/A   |
| Giant Clams       | <i>Hippopus hippopus</i> , <i>H. porcellanus</i> , <i>Tridacna costata</i> , <i>T. derasa</i> , <i>T. gigas</i> , <i>T. Squamosa</i> , and <i>T. tevoroa</i> | Positive (82 FR 28946, 06/26/2017)              | TBA (status review ongoing)   | TBA   | N/A   | N/A   |
| Green sea turtle  | <i>Chelonia mydas</i>  | Positive (77 FR 45571, 8/1/2012)                | Identification of 11 DPSs, endangered and threatened (80 FR 15271, 3/23/2015) | 11 DPSs listed as endangered and threatened (81 FR 20057, 4/6/2016) | In development, proposal expected TBA <sup>a</sup>                  | TBA   |

| Species                |                               | Listing Process   |   |  | Post-Listing Activity  |               |
|------------------------|-------------------------------|---|---|--|--|---------------|
| Common Name            | Scientific Name               | 90-Day Finding  | 12-Month Finding / Proposed Rule  | Final Rule   | Critical Habitat   | Recovery Plan |
| Leatherback sea turtle | <i>Dermochelys coriacea</i>   | Positive 90-day finding on a petition to identify the Northwest Atlantic leatherback turtle as a DPS (82 FR 57565, 12/06/2017)  | TBA   | TBA  | N/A  | N/A           |
| Humpback whale         | <i>Megaptera novaeangliae</i> | Positive 90-day finding on petition to classify the North Pacific population as DPS and delist the DPS (78 FR 53391, 8/29/2013) | Revision of species-wide listing and listing of four DPSs as threatened or endangered (80 FR 22304) | Revision of species wide listing; Western North Pacific DPS listed as endangered (81 FR 62259, 9/8/2016) | Critical habitat of Western North Pacific DPS in development, proposal expected 2019 | TBA           |

<sup>a</sup> NMFS and USFWS have been tasked with higher priorities regarding sea turtle listings under the ESA, and do not anticipate proposing green turtle critical habitat designations in the immediate future.

#### 2.4.5 Identification of Research, Data, and Assessment Needs

The following research, data, and assessment needs for insular fisheries were identified by the Council's Protected Species Advisory Committee and Plan Team:

- Improve the precision of commercial and non-commercial fisheries data to improve understanding of potential protected species impacts.
- Define and evaluate innovative approaches to derive robust estimates of protected species interactions in insular fisheries.

## 2.5 CLIMATE AND OCEANIC INDICATORS

### 2.5.1 Introduction

Over the past few years, the Council has incorporated climate change into the overall management of the fisheries over which it has jurisdiction. This 2018 Annual SAFE Report includes a now standard chapter on indicators of climate and oceanic conditions in the Western Pacific region. These indicators reflect global climate variability and change as well as trends in local oceanographic conditions.

The reasons for the Council's decision to provide and maintain an evolving discussion of climate conditions as an integral and continuous consideration in their deliberations, decisions, and reports are numerous:

- Emerging scientific and community understanding of the impacts of changing climate conditions on fishery resources, the ecosystems that sustain those resources, and the communities that depend upon them;
- Recent Federal Directives including the 2010 implementation of a National Ocean Policy that identified Resiliency and Adaptation to Climate Change and Ocean Acidification as one of nine National priorities as well as the development of a Climate Science Strategy by NMFS in 2015 and the subsequent development of the Pacific Islands Regional Action Plan for climate science; and
- The Council's own engagement with NOAA as well as jurisdictional fishery management agencies in American Samoa, CNMI, Guam, and Hawaii as well as fishing industry representatives and local communities in those jurisdictions.

In 2013, the Council began restructuring its Marine Protected Area/Coastal and Marine Spatial Planning Committee to include a focus on climate change, and the committee was renamed as the Marine Planning and Climate Change (MPCC) Committee. In 2015, based on recommendations from the committee, the Council adopted its Marine Planning and Climate Change Policy and Action Plan, which provided guidance to the Council on implementing climate change measures, including climate change research and data needs. The revised Pelagic FEP (February 2016) included a discussion on climate change data and research as well as a new objective (Objective 9) that states the Council should consider the implications of climate change in decision-making, with the following sub-objectives:

- To identify and prioritize research that examines the effects of climate change on Council-managed fisheries and fishing communities.
- To ensure climate change considerations are incorporated into the analysis of management alternatives.
- To monitor climate change related variables via the Council's Annual Reports.
- To engage in climate change outreach with U.S. Pacific Islands communities.

Beginning with the 2015 report, the Council and its partners began providing continuing descriptions of changes in a series of climate and oceanic indicators.

This annual report focuses previous years' efforts by refining existing indicators and improving communication of their relevance and status. Future reports will include additional indicators as the information becomes available and their relevance to the development, evaluation, and revision of the FEPs becomes clearer. Working with national and jurisdictional partners, the Council will make all datasets used in the preparation of this and future reports available and easily accessible.

### **2.5.2 Response to Previous Plan Team and Council Recommendations**

At its 170<sup>th</sup> meeting from June 20-22, 2017, the Council directed staff to support the development of community training and outreach materials and activities on climate change. In addition, the Council directed staff to coordinate a “train-the-trainers” workshop that includes NOAA scientists who presented at the 6th Marine Planning and Climate Change Committee (MPCCC) meeting and the MPCCC committee members in preparation for community workshops on climate and fisheries. The Council and NOAA partnered to deliver the workshops in the fall of 2017 to the MPCCC members in Hawaii (with the Hawaii Regional Ecosystem Advisory Committee), as well as American Samoa, Guam, and the CNMI (with their respective Advisory Panel groups). Feedback from workshop participants has been incorporated into this year's climate and oceanic indicator section. To prepare for community outreach, Guam-based MPCCC members conducted a climate change survey and shared the results with the MPCCC at its 7<sup>th</sup> meeting on April 10<sup>th</sup> and 11<sup>th</sup>, 2018.

Prior to holding its 8<sup>th</sup> meeting, the MPCCC was disbanded in early 2019, re-allocating its responsibilities among its members already on other committees or teams, such as the Fishery Ecosystem Plan Teams.


The Council also directed staff to explore funding avenues to support the development of additional oceanic and climate indicators, such as wind and extratropical storms. These indicators were added to this module by corresponding Plan Team members in 2018.

### **2.5.3 Conceptual Model**

In developing this chapter, the Council relied on a number of recent reports conducted in the context of the U.S. National Climate Assessment including, most notably, the 2012 Pacific Islands Regional Climate Assessment (PIRCA) and the Ocean and Coasts chapter of the 2014 report on a Pilot Indicator System prepared by the National Climate Assessment and Development Advisory Committee (NCADAC).

The Advisory Committee Report presented a possible conceptual framework designed to illustrate how climate factors can connect to and interact with other ecosystem components to impact ocean and coastal ecosystems and human communities. The Council adapted this model with considerations relevant to the fishery resources of the Western Pacific region:

**Indicators of Change to Archipelagic Coastal and Marine Systems\***  
*(Items in red to be monitored for 2015 Annual Reports of the Archipelagic Fishery Ecosystem Plans for the Western Pacific Region)*



**Figure 23. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability**

As described in the 2014 NCADAC report, the conceptual model presents a “simplified representation of climate and non-climate stressors in coastal and marine ecosystems.” For the purposes of this Annual Report, the modified Conceptual Model allows the Council and its partners to identify indicators of interest to be monitored on a continuing basis in coming years. The indicators shown in red were considered for inclusion in the Annual SAFE Reports, though the final list of indicators varied somewhat. Other indicators will be added over time as data become available and an understanding of the causal chain from stressors to impacts emerges.

The Council also hopes that this Conceptual Model can provide a guide for future monitoring and research. This guide will ideally enable the Council and its partners to move forward from observations and correlations to understanding the specific nature of interactions, and to develop capabilities to predict future changes of importance in the developing, evaluating, and adapting of FEPs in the Western Pacific region.



## 2.5.4 Selected Indicators


The primary goal for selecting the Indicators used in this (and future reports) is to provide fisheries-related communities, resource managers, and businesses with climate-related situational awareness. In this context, Indicators were selected to:

- Be fisheries relevant and informative;
- Build intuition about current conditions in light of changing climate;
- Provide historical context; and
- Recognize patterns and trends.


In this context, this section includes the following climate and oceanic indicators:

- Atmospheric concentration of carbon dioxide (CO<sub>2</sub>)
- Oceanic pH at Station ALOHA;
- Oceanic Niño Index (ONI);
- Pacific Decadal Oscillation (PDO);
- Tropical cyclones;
- Sea surface temperature (SST);
- Coral Thermal Stress Exposure
- Chlorophyll-A
- Rainfall
- Sea Level (Sea Surface Height)


Figure 24 and Figure 25 provide a description of these indicators and illustrate how they are connected to each other in terms of natural climate variability and anthropogenic climate change.



**Figure 24. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability**



**Figure 25 Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of anthropogenic climate change**



**Figure 26. Regional spatial grids representing the scale of the climate change indicators being monitored**

#### 2.5.4.1 Atmospheric Concentration of Carbon Dioxide at Mauna Loa

Rationale: Atmospheric carbon dioxide is a measure of what human activity has already done to affect the climate system through greenhouse gas emissions. It provides quantitative information in a simplified, standardized format that decision makers can easily understand. This indicator

demonstrates that the concentration (and, in turn, warming influence) of greenhouse gases in the atmosphere has increased substantially over the last several decades.


Status: Atmospheric CO<sub>2</sub> is increasing exponentially. This means that atmospheric CO<sub>2</sub> is increasing at a faster rate each year. In 2018, the annual mean concentration of CO<sub>2</sub> was 409 ppm. In 1959, the first year of the time series, it was 316 ppm. The annual mean passed 350 ppm in 1988 and 400 ppm in 2015 (NOAA 2019b).

Description: Monthly mean atmospheric carbon dioxide (CO<sub>2</sub>) at Mauna Loa Observatory, Hawaii in parts per million (ppm) from March 1958 to present. The observed increase in monthly average carbon dioxide concentration is primarily due to CO<sub>2</sub> emissions from fossil fuel burning. Carbon dioxide remains in the atmosphere for a very long time, and emissions from any location mix throughout the atmosphere in about one year. The annual oscillations at Mauna Loa, Hawaii are due to the seasonal imbalance between the photosynthesis and respiration of plants on land. During the summer growing season photosynthesis exceeds respiration and CO<sub>2</sub> is removed from the atmosphere, whereas outside the growing season respiration exceeds photosynthesis and CO<sub>2</sub> is returned to the atmosphere. The seasonal cycle is strongest in the northern hemisphere because of this hemisphere's larger land mass.

Timeframe: Annual, monthly.

Region/Location: Mauna Loa, Hawaii but representative of global atmospheric carbon dioxide concentration.

Measurement Platform: *In-situ* station.



**Figure 27. Monthly mean (red) and seasonally-corrected (black) atmospheric carbon dioxide at Mauna Loa Observatory, Hawaii**

#### 2.5.4.2 Oceanic pH

Rationale: Oceanic pH is a measure of how greenhouse gas emissions have already impacted the ocean (NOAA PMEL 2019). This indicator demonstrates that oceanic pH has decreased significantly over the past several decades (i.e., the ocean has become more acidic). Increasing ocean acidification limits the ability of marine organisms to build shells and other calcareous structures. Recent research has shown that pelagic organisms such as pteropods and other prey for commercially-valuable fish species are already being negatively impacted by increasing acidification (Feely et al., 2016). The full impact of ocean acidification on the pelagic food web is an area of active research (Fabry et al., 2008).

Status: The ocean is roughly 9.4% more acidic than it was nearly 30 years ago at the start of this time series. Over this time, pH has declined by 0.0389 at a constant rate. In 2017, the most recent year for which data are available, the average pH was 8.07. Additionally, small variations seen over the course of the year are now outside the range seen in the first year of the time series. The highest pH value reported for the most recent year (8.0831) is lower than the lowest pH value reported in the first year of the time series (8.0845).


Description: Trends in surface (5 m) pH at Station ALOHA, north of Oahu (22.75°N, 158°W), collected by the Hawai'i Ocean Time Series (HOT) from October 1988 to 2017 (2018 data are not yet available). Oceanic pH is a measure of ocean acidity, which increases as the ocean absorbs carbon dioxide from the atmosphere. Lower pH values represent greater acidity. Oceanic

pH is calculated from total alkalinity (TA) and dissolved inorganic carbon (DIC). Total alkalinity represents the ocean's capacity to resist acidification as it absorbs CO<sub>2</sub> and the amount of CO<sub>2</sub> absorbed is captured through measurements of DIC. The multi-decadal time series at Station ALOHA represents the best available documentation of the significant downward trend in oceanic pH since the time series began in 1988. Oceanic pH varies over both time and space, though the conditions at Station ALOHA are considered broadly representative of those across the Western and Central Pacific's pelagic fishing grounds.

Timeframe: Monthly.

Region/Location: Station ALOHA: 22.75°N, 158°W.

Measurement Platform: *In-situ* station.



**Figure 28. Oceanic pH (black) and its trend (red) at Station ALOHA from 1989 – 2016**

#### 2.5.4.3 Oceanic Niño Index

Rationale: The El Niño – Southern Oscillation (ENSO) cycle is known to have impacts on Pacific fisheries including tuna fisheries. The ONI focuses on ocean temperature, which has the most direct effect on these fisheries.


Status: In 2018, the ONI transitioned from a weak La Niña to neutral conditions.

Description: The three-month running mean of satellite remotely-sensed sea surface temperature (SST) anomalies in the Niño 3.4 region (5°S – 5°N, 120° – 170°W). The Oceanic Niño Index (ONI) is a measure of the El Niño – Southern Oscillation (ENSO) phase. Warm and cool phases, termed El Niño and La Niña respectively, are based in part on an ONI threshold of  $\pm 0.5$  °C being met for a minimum of five consecutive overlapping seasons. Additional atmospheric indices are needed to confirm an El Niño or La Niña event, as the ENSO is a coupled ocean-atmosphere phenomenon. The atmospheric half of ENSO is measured using the Southern Oscillation Index.

Timeframe: Every three months.

Region/Location: Niño 3.4 region, 5°S – 5°N, 120° – 170°W.

Measurement Platform: *In-situ* station, satellite, model.



**Figure 29. Oceanic Niño Index from 1950-2018 (top) and 2000–2018 (bottom) with El Niño periods in red and La Niña periods in blue**

#### 2.5.4.4 Pacific Decadal Oscillation

Rationale: The Pacific Decadal Oscillation (PDO) was initially named by fisheries scientist Steven Hare in 1996 while researching connections between Alaska salmon production cycles and Pacific climate. Like ENSO, the PDO reflects changes between periods of persistently warm or persistently cool ocean temperatures, but over a period of 20 to 30 years (versus six to 18 months for ENSO events). The climatic finger prints of the PDO are most visible in the Northeastern Pacific, but secondary signatures exist in the tropics.


Status: The PDO was positive, or warm, for much of 2018. In March and June, the index dipped just below zero but returned to a positive value the following months. PDO index values were not yet available for the last three months of 2018 at the time of publication.

Description: The PDO is often described as a long-lived El Niño-like pattern of Pacific climate variability. As seen with the better-known ENSO, extremes in the PDO pattern are marked by widespread variations in the Pacific Basin and the North American climate. In parallel with the ENSO phenomenon, the extreme cases of the PDO have been classified as either warm or cool, as defined by ocean temperature anomalies in the northeast and tropical Pacific Ocean. When SST is below average in the interior North Pacific and warm along the North American coast, and when sea level pressures are below average in the North Pacific, the PDO has a positive value. When the climate patterns are reversed, with warm SST anomalies in the interior and cool SST anomalies along the North American coast, or above average sea level pressures over the North Pacific, the PDO has a negative value. (<https://www.ncdc.noaa.gov/teleconnections/pdo/>).

Timeframe: Annual, monthly.

Region/Location: Pacific Basin north of 20°N.

Measurement Platform: *In-situ* station, satellite, model.



**Figure 30. Pacific Decadal Oscillation from 1950–2018 (top) and 2000–2018 (bottom) with positive warm periods in red and negative cool periods in blue**

#### 2.5.4.5 Tropical Cyclones

Rationale: The effects of tropical cyclones are numerous and well known. At sea, storms disrupt and endanger shipping traffic as well as fishing effort and safety. The Hawai`i longline fishery, for example, has had serious problems with vessels dodging storms at sea, delayed departures, and inability to make it safely back to Honolulu because of bad weather. When cyclones encounter land, their intense rains and high winds can cause severe property damage, loss of life, soil erosion, and flooding. Associated storm surge, the large volume of ocean water pushed toward shore by cyclones' strong winds, can cause severe flooding and destruction.

Status:

*Eastern North Pacific.* Overall, the 2018 eastern Pacific hurricane season featured well above average activity. There were 22 named storms, of which 12 became hurricanes and 9 became major hurricanes - category 3 or higher on the Saffir-Simpson Hurricane Wind Scale. This compares to the long-term averages of 15 named storms, 8 hurricanes, and 4 major hurricanes. There were also 3 tropical depressions that did not reach tropical storm strength. In terms of Accumulated Cyclone Energy (ACE), which measures the strength and duration of tropical storms and hurricanes, activity in the basin in 2018 was the 3<sup>rd</sup> highest on record, behind 1990 and 1992. Summary inserted from <https://www.nhc.noaa.gov/text/MIATWSEP.shtml>.

*Central North Pacific.* Tropical cyclone activity in 2018 was high. The ACE index was the second highest since 1980, second only to 2015, and well above the 1981 – 2010 average of just under 20 ( $\times 10^4$  knots<sup>2</sup>). Of note was Hurricane Lane, which reached Category 5 strength and passed within 110 miles of Honolulu. Lane was only the second Category 5 hurricane to pass within 250 miles of Hawaii, with the last being Hurricane John in 1994. Some of the impacts associated with Hurricane Lane include widespread reports of more than 40 inches of rain the islands of Hawaii and Kauai. There was one preliminary report of more than 52 inches of rain. At least one fatality was blamed on Hurricane Lane. Summary inserted from <https://www.ncdc.noaa.gov/sotc/tropical-cyclones/201808>.

*Western North Pacific.* Tropical cyclone activity was roughly average. The ACE Index was slightly above average in the Western North Pacific. Of note was Super Typhoon Yutu which made landfall on the islands of Tinian and Saipan as a Category 5 equivalent typhoon with estimated winds of 180 mph and a central minimum pressure of 905 mb. This marked the second strongest tropical cyclone to impact any U.S. territory on record. The storm devastated most of Tinian and Saipan with nearly every structure on the two islands being damaged or destroyed, including the Saipan International Airport. There were two fatalities reported in the Northern Marianas. Summary inserted from <https://www.ncdc.noaa.gov/sotc/tropical-cyclones/201810>.

*South Pacific.* Tropical cyclone activity and the ACE Index were below average in 2018.

Description: This indicator uses historical data from the NOAA National Climate Data Center (NCDC) International Best Track Archive for Climate Stewardship to track the number of tropical cyclones in the western, central, eastern, and southern Pacific basins. This indicator also monitors the Accumulated Cyclone Energy (ACE) Index and the Power Dissipation Index which are two ways of monitoring the frequency, strength, and duration of tropical cyclones based on wind speed measurements.



The annual frequency of storms passing through each basin is tracked and a stacked time series plot shows the representative breakdown of Saffir-Simpson hurricane categories.

Every cyclone has an ACE Index value, which is a number based on the maximum wind speed measured at six-hourly intervals over the entire time that the cyclone is classified as at least a tropical storm (wind speed of at least 34 knots; 39 mph). Therefore, a storm's ACE Index value accounts for both strength and duration. This plot shows the historical ACE values for each hurricane/typhoon season and has a horizontal line representing the average annual ACE value.

Timeframe: Annual.

Region/Location:


Eastern North Pacific: east of 140° W, north of the equator.

Central North Pacific: 180° - 140° W, north of the equator.

Western North Pacific: west of 180°, north of the equator.

South Pacific: south of the equator.

Measurement Platform: Satellite.



**Figure 31. 2018 Pacific basin tropical cyclone tracks**




Figure 32. 2018 tropical storm totals by region




Figure 33. 2018 Accumulated Cyclone Energy (ACE) Index by region

#### 2.5.4.6 Sea Surface Temperature and Anomaly

Rationale: Sea surface temperature is one of the most directly observable existing measures for tracking increasing ocean temperatures. SST varies in response to natural climate cycles such as the El Niño – Southern Oscillation (ENSO) and is projected to rise as a result of anthropogenic climate change. Both short-term variability and long-term trends in SST impact the marine ecosystem. Understanding the mechanisms through which organisms are impacted and the time scales of these impacts is an area of active research.

Status: Annual mean SST was 28.26°C in 2018. Over the period of record, annual SST has increased at a rate of 0.022°C/year. Monthly SST values in 2018 ranged from 27.02 – 29.17°C, within the climatological range of 25.48 – 30.43 °C. The annual anomaly was 0.188 °C hotter than average, with intensification in the northern islands.

Note that from the top to bottom in **Error! Reference source not found.**, panels show climatological SST (1982-2017), 2018 SST anomaly, time series of monthly mean SST, and time series of monthly SST anomaly. The white box in the upper panels indicates the area over which SST is averaged for the time series plots.




Description: Satellite remotely-sensed monthly sea surface temperature (SST) is averaged across the Marianas Grid (13° – 21°N, 144° – 146°E). A time series of monthly mean SST averaged over the American Samoa Grid Region is presented. Additionally, spatial climatology and anomalies are shown. Data from NOAA Pathfinder v5.3 (NOAA 2019c).

Timeframe: Monthly.

Region/Location: Marianas Grid (13° – 21°N, 144° – 146°E).

Measurement Platform: Satellite.

Sourced from: NOAA OceanWatch (2018).



**Figure 34. Sea surface temperature climatology and anomalies from 1982-2018**

#### **2.5.4.7 Coral Thermal Stress Exposure: Degree Heating Weeks**

Rationale: Degree heating weeks are one of the most widely used metrics for assessing exposure to coral bleaching-relevant thermal stress.

Status: After a series of stress events in 2013, 2014, 2016, and 2017, the Marianas have shown no significant coral heat stress since.

Description: Here we present a metric of exposure to thermal stress that is relevant to coral bleaching. Degree Heating Weeks (DHW) measure time and temperature above a reference ‘summer maximum’, presented as rolling sum weekly thermal anomalies over a 12-week period. Higher DHW measures imply a greater likelihood of mass coral bleaching or mortality from thermal stress.


The NOAA Coral Reef Watch program uses satellite data to provide current reef environmental conditions to quickly identify areas at risk for [coral bleaching](#). Bleaching is the process by which corals lose the symbiotic algae that give them their distinctive colors. If a coral is severely bleached, disease and death become likely.

The NOAA Coral Reef Watch (CRW) daily 5-km satellite coral bleaching Degree Heating Week (DHW) product presented here shows accumulated heat stress, which can lead to coral bleaching and death. The scale goes from 0 to 20 °C-weeks. The DHW product accumulates the instantaneous bleaching heat stress (measured by Coral Bleaching HotSpots) during the most-recent 12-week period. It is directly related to the timing and intensity of coral bleaching. Significant coral bleaching usually occurs when DHW values reach 4 °C-weeks. By the time DHW values reach 8 °C-weeks, widespread bleaching is likely and significant mortality can be expected (NOAA Coral Reef Watch 2019).

Timeframe: 2013-2018, Daily data.

Region/Location: Global.

Measurement Platform: [NOAA/NESDIS operational daily global 5km geostationary-polar-orbiting \(Geo-Polar\) Blended Night-only SST Analysis](#)



**Figure 35. Coral Thermal Stress Exposure measured at CNMI Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)**

#### 2.5.4.8 Chlorophyll-A and Anomaly

Rationale: Chlorophyll-A is one of the most directly observable measures we have for tracking increasing ocean productivity.

Status: Annual mean Chl-A was 0.054 mg/m<sup>3</sup> in 2018. Over the period of record, annual Chl-A has shown weak but significant linear decrease at a rate of 0.00014 mg/m<sup>3</sup>. Monthly Chl-A values in 2018 ranged from 0.051-0.059 mg/m<sup>3</sup>, within the climatological range of 0.041 – 0.088 mg/m<sup>3</sup>. The annual anomaly was essentially in line with climatological values -- 0.003 mg/m<sup>3</sup> higher than average.




Description: Chlorophyll-A Concentration from 1998-2018, derived from the MODIS Ocean Color sensor aboard the NASA Aqua Satellite. A monthly climatology was generated across the entire period (1982-2018) to provide both a 2018 spatial anomaly, and an anomaly time series.

The following text was inserted from the OceanWatch Central Pacific Node (NOAA 2019a). The MODIS (Moderate Resolution Imaging Spectro-radiometer) sensor was deployed onboard the NASA Aqua satellite. It is a multi-disciplinary sensor providing data for the ocean, land, aerosol, and cloud research and is used for detecting chlorophyll-a concentrations in the world's oceans, among other applications. Aqua MODIS views the entire Earth's surface every two days, acquiring data in 36 spectral bands. The data available here is the latest reprocessing from June 2015, which NASA undertook to correct for some sensor drift issues.

Timeframe: 2003-2018, Daily data available, Monthly means shown.

Region/Location: Global.

Measurement Platform: *MODIS sensor on NASA Aqua Satellite*



**Figure 36. Chlorophyll-A (Chl-A) and Chl-A Anomaly from 1982-2018**



#### 2.5.4.9 Rainfall (CMAP Precipitation)

Rationale: Rainfall may have substantive effects on the nearshore environment and is a potentially important co-variate with the landings of particular stocks.

Description: The CPC Merged Analysis of Precipitation (CMAP) is a technique which produces pentad and monthly analyses of global precipitation in which observations from rain gauges are merged with precipitation estimates from several satellite-based algorithms, such as infrared and microwave (NOAA 2002). The analyses are on a 2.5 x 2.5 degree latitude/longitude grid and extend back to 1979. CMAP Precipitation data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <https://www.esrl.noaa.gov/psd/>. The data are comparable (but should not be confused with) similarly combined analyses by the [Global Precipitation Climatology Project](#) described in Huffman et al. (1997).


It is important to note that the input data sources to make these analyses are not constant throughout the period of record. For example, SSM/I (passive microwave - scattering and emission) data became available in July 1987; prior to that the only microwave-derived estimates available are from the MSU algorithm (Spencer 1993) which is emission-based thus precipitation estimates are available only over oceanic areas. Furthermore, high temporal resolution IR data from geostationary satellites (every 3-hr) became available during 1986; prior to that, estimates from the OPI technique (Xie and Arkin 1997) are used based on OLR from orbiting satellites.

The merging technique is thoroughly described in Xie and Arkin (1997). Briefly, the methodology is a two-step process. First, the random error is reduced by linearly combining the satellite estimates using the maximum likelihood method, in which case the linear combination coefficients are inversely proportional to the square of the local random error of the individual data sources. Over global land areas the random error is defined for each time period and grid location by comparing the data source with the rain gauge analysis over the surrounding area. Over oceans, the random error is defined by comparing the data sources with the rain gauge observations over the Pacific atolls. Bias is reduced when the data sources are blended in the second step using the blending technique of Reynolds (1988).

Timeframe: Monthly.

Region/Location: Global.

Measurement Platform: *In-situ* station gauges and satellite data.



**Figure 37. CMAP precipitation across the Marianas Grid with 2018 values in red**

### 2.5.3.9 Sea Level (Sea Surface Height and Anomaly)

Rationale: Rising sea levels can result in a number of coastal impacts, including inundation of infrastructure, increased damage resulting from storm-driven waves and flooding, and saltwater intrusion into freshwater supplies.

Description: Monthly mean sea level time series, including extremes


Timeframe: Monthly

Region/Location: Observations from selected sites within the Samoan Archipelago


Measurement Platform: Satellite and *in situ* tide gauges

#### 2.5.3.9.1 Basin-Wide Perspective

This image of the mean sea level anomaly for March 2019 compared to 1993-2013 climatology from satellite altimetry provides a glimpse into how the current weak El Niño continues to affect sea level across the Pacific Basin. The image captures the fact that sea level continues to be lower in the Western Pacific and higher in the Central and Eastern Pacific (a standard pattern during El Niño events - this basin-wide perspective provides a context for the location-specific sea level/sea surface height images that follow).




**Figure 38a. Sea surface height and anomaly**



**Figure 38b.** Quarterly time series of mean sea level anomalies during 2018 show no pattern of El Niño throughout the year according to satellite altimetry measurements of sea level height (unlike 2015).

[http://sealevel.jpl.nasa.gov/science/elninopdo/latestdata/archive/index.cfm?y=2017\)](http://sealevel.jpl.nasa.gov/science/elninopdo/latestdata/archive/index.cfm?y=2017)




### 2.5.3.9.2 Local Sea Level

These time-series from *in situ* tide gauges provide a perspective on sea level trends within each Archipelago (Tide Station Time Series from NOAA/COOPS).

The following figures and descriptive paragraphs were inserted from NOAA (2018). Figure 31 shows the monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents. The long-term linear trend is also shown, including its 95% confidence interval. The plotted values are relative to the most recent [Mean Sea Level datum established by CO-OPS](#). The calculated trends for all stations are available as a [table in millimeters/year and in feet/century](#). If present, solid vertical lines indicate times of any major earthquakes in the vicinity of the station and dashed vertical lines bracket any periods of questionable data or datum shift.

The relative sea level rise trend is 3.68 millimeters/year with a 95% confidence interval of +/- 3.98 mm/yr based on monthly mean sea level data from 1993 to 2018 which is equivalent to a change of 1.21 feet in 100 years. Trend for 1948-1993 is -0.85 +/- 1.76 mm/yr.



**Figure 39. Monthly mean sea level without regular seasonal variability due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents**

### 2.5.5 References

- Aviso, 2019. El Niño Bulliten – Latest plots. Ocean indicator products, Center for Topographic Studies of the Ocean and Hydrosphere, Centre National D'études Spatiales. Available at <https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/el-nino-bulletin.html>. Updated April 2019.
- Fabry, V.J., Seibel, B.A., Feely, R.A., and J.C. Orr, 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65, pp. 414-432.
- Feely, R.A., Alin, S.R., Carter, B., Bednarsek, N., Hales, B., Chan, F., Hill, T.M., Gaylord, B., Sanford, E., Byrne, R.H., Sabine, C.L., Greeley, D., and L. Juranek, 2016. Chemical and biological impacts of ocean acidification along the west coast of North America. *Estuarine, Coastal and Shelf Science*, 183, pp. 260-270. doi: 10.1016/j.ecss.2016.08.043.
- Huffman, G.J., Adler, R.F., Arkin, P., Chang, A., Ferraro, R., Gruber, A., Janowiak, J., McNab, A., Rudolf, B. and U. Schneider, 1997. The global precipitation climatology project (GPCP) combined precipitation dataset. *Bulletin of the American Meteorological Society*, 78(1), pp.5-20.
- Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.K., Hnilo, J.J., Fiorino, M. and G.L. Potter, 2002. NCEP–DOE AMIP-II Reanalysis (R-2), *B. Am. Meteorol. Soc.*, 83, pp. 1631–1643,
- Keeling, C.D., Bacastow, R.B., Bainbridge, A.E., Ekdahl, C.A., Guenther, P.R., and L.S. Waterman, 1976. Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii. *Tellus*, 28, pp. 538-551.
- Knapp, K.R., M.C. Kruk, D.H. Levinson, H.J. Diamond, and C.J. Neumann, 2010: The International Best Track Archive for Climate Stewardship (IBTrACS): Unifying tropical cyclone best track data. *Bulletin of the American Meteorological Society*, 91, pp. 363-376. doi:10.1175/2009BAMS2755.1.
- Li, X., Pichel, W.G., Clemente-Colón, P., and J.F. Sapper, 2001a. Deriving the operational nonlinear multi-channel sea surface temperature algorithm coefficients for NOAA-15 AVHRR/3, *Int. J. Remote Sens.*, 22(4), pp. 699 - 704.
- Li, X., Pichel, W.G., Clemente-Colón, P., Krasnopolsky, V., and J.F Sapper, 2001b. Validation of coastal sea and lake surface temperature measurements derived from NOAA/AVHRR Data, *Int. J. Remote Sens.*, 22(7), pp. 1285-1303.
- Liu, G., Heron, S.F., Eakin, C.M., Muller-Karger, F.E., Vega-Rodriguez, M., Guild, L.S., De La Cour, J.L., Geiger, E.F., Skirving, W.J., Burgess, T.F. and A.E. Strong, 2014. Reef-scale thermal stress monitoring of coral ecosystems: new 5-km global products from NOAA Coral Reef Watch. *Remote Sensing*, 6(11), pp.11579-11606.
- Mantua, N., 2018. The Pacific Decadal Oscillation. Available at <http://research.jisao.washington.edu/pdo/>. Accessed March 2018.

- Mélin, F., V. Vantrepotte, A. Chuprin, M. Grant, T. Jackson, and S. Sathyendranath, 2017. "Assessing the fitness-for-purpose of satellite multi-mission ocean color climate data records: A protocol applied to OC-CCI chlorophyll-a data." *Remote sensing of environment*, 203, pp. 139-151.
- NOAA, 2002. CPC Merged Analysis of Precipitation. National Weather Service, National Centers for Environmental Prediction, Climate Prediction Center. Available at [https://www.cpc.ncep.noaa.gov/products/global\\_precip/html/wpage.cmap.html](https://www.cpc.ncep.noaa.gov/products/global_precip/html/wpage.cmap.html). Updated 25 September 2002.
- NOAA, 2018. Sea level Trends. NOAA Tides & Currents. Available at [https://tidesandcurrents.noaa.gov/sltrends/sltrends\\_station.shtml?stnid=1770000](https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?stnid=1770000). Updated 8 August 2018.
- NOAA, 2019a. Chlorophyll a Concentration, Aqua MODIS. OceanWatch – Central Pacific Node. NASA/GSFC/OBPG. Available at <https://oceanwatch.pifsc.noaa.gov/contact.html>. Updated 21 February 2019.
- NOAA, 2019b. Full Mauna Loa CO<sub>2</sub> record. Trends in Atmospheric Carbon Dioxide. NOAA Earth System Research Laboratory, Global Monitoring Division. Available from <https://www.esrl.noaa.gov/gmd/ccgg/trends/full.html>. Updated 28 May 2019.
- NOAA, 2019c. Sea Surface Temperature Datasets. AVHRR Pathfinder v.5, AVRR GAC v.5, and GOES-POES v.5. OceanWatch – Central Pacific Node. NASA/GSFC/OBPG. Available at <https://oceanwatch.pifsc.noaa.gov/contact.html>. Updated 21 February 2019.
- NOAA Coral Reef Watch, 2019, updated daily. Version 3.1 Daily Global 5-km Satellite Coral Bleaching Degree Heating Week Product, 2013-2018. College Park, Maryland, USA: NOAA Coral Reef Watch. Available at <https://coralreefwatch.noaa.gov/satellite/hdf/index.php>.
- NOAA NCEI, 2019. State of the Climate: Hurricanes and Tropical Storms for Annual 2018. Available from <http://www.ncdc.noaa.gov/sotc/tropical-cyclones/201713>. Accessed 30 March 2018.
- NOAA PMEL, 2019. A primer on pH. NOAA Pacific Marine Environmental Laboratory, Carbon Program. Available at <https://www.pmel.noaa.gov/co2/story/A+primer+on+pH>.
- Reynolds, R.W., 1988. A real-time global sea surface temperature analysis. *Journal of Climate*, 1(1), pp. 75-87.
- Savchenko, A., Ouzounov, D., Ahmad, S., Acker, J., Leptoukh, G., Koziana, J., and D. Nickless, 2004. Terra and Aqua MODIS products available from NASA GES DAAC. *Advances in Space Research*, 34(4), pp. 710-714.
- Spencer, R.W., 1993. Global oceanic precipitation from the MSU during 1979—91 and comparisons to other climatologies. *Journal of Climate*, 6(7), pp.1301-1326.

State of Environmental Conditions in Hawaii and the U.S. Affiliated Pacific Islands under a Changing Climate, 2017. Coordinating Authors: J.J. Marra and M.C. Kruk. Contributing Authors: M. Abecassis; H. Diamond; A. Genz; S.F. Heron; M. Lander; G. Liu; J. T. Potemra; W.V. Sweet; P. Thompson; M.W. Widlansky; and P. Woodworth-Jefcoats. September, 2017. NOAA NCEI.

Stowe, L.L., Davis, P.A., and E.P. McClain, 1999. Scientific basis and initial evaluation of the CLAVR-1 global clear/cloud classification algorithm for the advanced very high resolution radiometer. *J. Atmos. Oceanic Technol.*, 16, pp. 656-681.

Thoning, K.W., Tans, P.P., and W.D. Komhyr, 1989. Atmospheric carbon dioxide at Mauna Loa Observatory 2. Analysis of the NOAA GMCC data, 1974-1985. *Journal of Geophysical Research*, 94, pp. 8549-8565.

Walton C.C., Pichel, W.G., Sapper, J.F., and D.A. May, 1998. The development and operational application of nonlinear algorithms for the measurement of sea surface temperatures with the NOAA polar-orbiting environmental satellites, *J. Geophys. Res.*, 103(C12), pp. 27999-28012.

Xie, P. and P.A. Arkin, 1997. Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bulletin of the American Meteorological Society*, 78(11), pp. 2539-2558.



## **2.6 ESSENTIAL FISH HABITAT**

### **2.6.1 Introduction**

The Magnuson-Stevens Fishery Conservation and Management Act (MSA) includes provisions concerning the identification and conservation of essential fish habitat (EFH), and under the EFH final rule, habitat areas of particular concern (HAPC) (50 Code of Federal Regulations [CFR] 600.815). The MSA defines EFH as “those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity.” HAPC are those areas of EFH identified pursuant to 50 CFR 600.815(a)(8), and meeting one or more of the following considerations: (1) ecological function provided by the habitat is important; (2) habitat is sensitive to human-induced environmental degradation; (3) development activities are, or will be, stressing the habitat type; or (4) the habitat type is rare.

The National Marine Fisheries Service (NMFS) and regional fishery management councils must describe and identify EFH in fishery management plans (FMPs), minimize to the extent practicable the adverse effects of fishing on EFH, and identify other actions to encourage the conservation and enhancement of EFH. Federal agencies that authorize, fund, or undertake actions that may adversely affect EFH must consult with NMFS, and NMFS must provide conservation recommendations to federal and state agencies regarding actions that would adversely affect EFH. Regional fishery management councils also have the authority to comment on federal or state agency actions that would adversely affect the habitat, including EFH, of managed species.

The EFH Final Rule strongly recommends regional fisheries management councils and NMFS to conduct a review and revision of the EFH components of fisheries management plans every five years (600.815(a)(10)). The council’s FEPs state that new EFH information should be reviewed, as necessary, during preparation of the annual reports by the Plan Teams. Additionally, the EFH Final Rule states, “Councils should report on their review of EFH information as part of the annual Stock Assessment and Fishery Evaluation (SAFE) report prepared pursuant to §600.315(e).” The habitat portion of the annual report is designed to meet the FEP requirements and EFH Final Rule guidelines regarding EFH reviews.

National Standard 2 guidelines recommend that the SAFE report summarize the best scientific information available concerning the past, present, and possible future condition of EFH described by the FEPs.

#### **2.6.1.1 EFH Information**

The Western Pacific Regional Fishery Management Council (Council) FMPs include identification and description of EFH, lists of prey species and locations for each managed species, and optionally, HAPC. Impact-oriented components of FMPs include federal fishing activities that may adversely affect EFH; non-federal fishing activities that may adversely affect EFH; non-fishing activities that may adversely affect EFH; conservation and enhancement recommendations; and a cumulative impacts analysis on EFH. The last two components include the research and information needs section that feeds into the Council’s Five Year Research Priorities and the EFH update procedure that is described in the FEP.

The Council has described EFH for five management unit species (MUS) under its management authority: pelagic (PMUS), bottomfish (BMUS), crustaceans (CMUS), coral reef ecosystem (CREMUS), and precious corals (PCMUS).

EFH reviews of the biological components, including the description and identification of EFH, lists of prey species and locations, and HAPC, consist of three to four parts:

- Updated species descriptions, which can be found appended to the SAFE report. These can be used to directly update the FEP;
- Updated EFH levels of information tables, which can be found in Section 2.6.4;
- Updated research and information needs, which can be found in Section 2.6.5. These can be used to directly update the FEP; and
- An analysis that distinguishes EFH from all potential habitats used by the species, which is the basis for an options paper for the Council. This part is developed if enough information exists to refine EFH.

### **2.6.1.2 Habitat Objectives of FEP**

The habitat objective of the FEP is to refine EFH and minimize impacts to EFH, with the following sub-objectives:

- Review EFH and HAPC designations every five years based on the best available scientific information and update such designations based on the best available scientific information, when available, and
- Identify and prioritize research to: assess adverse impacts to EFH and HAPC from fishing (including aquaculture) and non-fishing activities, including, but not limited to, activities that introduce land-based pollution into the marine environment.

This annual report reviews the precious coral EFH components and non-fishing impacts components, resetting the five-year timeline for review. The Council's support of non-fishing activities research is monitored through the program plan and five year research priorities.

### **2.6.1.3 Response to Previous Council Recommendations**

At its 172<sup>nd</sup> meeting in March 2018, the Council recommended that staff develop an omnibus amendment updating the non-fishing impact to EFH sections of the FEPs, incorporating the non-fishing impacts EFH review report by Minton (2017) by reference. An options paper has been developed. The CNMI Joint Advisory Group provided comments on the non-fishing impacts review at a meeting held November 15, 2017, in Garapan. The Guam Joint Advisory Group also reviewed the report at their meeting held on November 17, 2017, in Tumon.

At its 173<sup>rd</sup> meeting in June 2018, the Council directed staff to develop options to redefine EFH and any HAPC for precious corals in Hawaii for Council consideration for an FEP amendment.

At its 174<sup>th</sup> meeting in October 2018, the Council directed staff to prepare an amendment to the Hawaii FEP to revise the Precious Corals EFH and selected the following preliminarily preferred options for the staff to further analyze:

- Action 1: Option 4 - Revise existing beds and designate new beds as EFH
- Action 2: Option 2 - Update Geographic Extent and Habitat Characteristics
- Action 3: Option 1 - Update the FEP narratives

An FEP amendment is being developed to present to the Council in mid-2019.

## **2.6.2 Habitat Use by MUS and Trends in Habitat Condition**

The Mariana Archipelago is a chain of islands in the western Pacific roughly oriented north-south. It is anchored at the southern end by the relatively large island of Guam at 13.5° north latitude. The Commonwealth of the Northern Mariana Islands (CNMI) stretch off to the north. The entire chain is approximately 425 miles long. The archipelago was named by Spanish explorers in the 16<sup>th</sup> Century in honor of Spanish Queen Mariana of Austria.

The total land area of Guam is approximately 212 square miles and its EEZ is just over 84,000 square miles. The CNMI consists of 14 main islands. From north to south these are: Farallon de Pajaros, Maug, Asuncion, Agrihan, Pagan, Alamagan, Guguan, Sarigan, Anatahan, Farallon de Medinilla, Saipan, Tinian, Aguijan, and Rota. Only Saipan, Rota, and Tinian are permanently inhabited, with 90% of the population residing on the island of Saipan. The total land area of the CNMI is 176.5 square miles and its EEZ is almost 300,000 square miles.

Guam and the southern islands of the CNMI are limestone, with level terraces and fringing coral reefs. The CNMI's northern islands are volcanic and sparsely inhabited, with active volcanoes on several islands, including Anatahan, Pagan, and Agrihan (the highest, at 3,166 feet). The archipelago has a tropical maritime climate moderated by seasonal northeast trade winds. While there is little seasonal temperature variation, there is a dry season (December to June) and a rainy season (July to November). The rainy season coincides with hurricane season, and the Mariana Archipelago is periodically impacted by powerful typhoons.

The Mariana Trench is located to the east of the chain and includes the deepest point in the world's oceans. The vertical measurement from the seafloor to Mount Tapotchau is 37,752 ft.

Essential fish habitat in the Marianas for the four MUS comprises all substrate from the shoreline to the 700 m isobath. The entire water column is described as EFH from the shoreline to the 700 m isobath, and the water column to a depth of 400 m is described as EFH from the 700 m isobath to the limit or boundary of the EEZ. While the coral reef ecosystems surrounding the islands in the Marianas have been subject to monitoring program through PIFSC CRED biennially since 2003, surveys are focused on the nearshore environments surrounding the islands (PIFSC 2010).

The mission of the PIFSC CREP is to “provide high-quality, scientific information about the status of coral reef ecosystems of the U.S. Pacific islands to the public, resource managers, and policymakers on local, regional, national, and international levels” (PIFSC 2011). CREP's Reef Assessment and Monitoring Program (RAMP) conducts comprehensive ecosystem monitoring surveys at about 50 island, atoll, and shallow bank sites in the Western Pacific region on a one-to three-year schedule (Brainard et al. 2008). CREP coral reef monitoring reports provide the most comprehensive description of nearshore habitat quality in the region. The benthic habitat mapping program provides information on the quantity of habitat.




Figure 40. Substrate EFH Limit of 700 m isobath around the Mariana Archipelago (from Ryan et al. 2009)


### 2.6.2.1 Habitat Mapping

Interpreted IKONOS benthic habitat maps in the 0 – 30 m depth range have been completed for all islands in the CNMI (Miller et al., 2011). Mapping products for the Marianas are available from the Pacific Islands Benthic Habitat Mapping Center.

**Table 71. Summary of habitat mapping in CNMI**

| Depth Range | Timeline/Mapping Product                    | Progress  | Source   |
|-------------|---|---|--|
| 0-30 m      | IKONOS Benthic Habitat Maps                 | All Islands   | Miller et al. (2011)   |
|             | 2000-2010 Bathymetry                        | 70%   | DesRochers (2016)  |
|             | 2011-2015 Multibeam Bathymetry              | -   | DesRochers (2016)  |
|             | 2011-2015, Satellite Worldview 2 Bathymetry | 15%   | DesRochers (2016)  |
| 30-150 m    | 2000-2010 Bathymetry                        | 85%   | DesRochers (2016)  |
|             | 2011-2015 Multibeam Bathymetry              | -   | DesRochers (2016)  |
| 15-2000 m   | Multibeam Bathymetry                        | Complete around all islands except Guam, Rota, and Agrigan                        | <a href="#">Pacific Islands Benthic Habitat Mapping Center</a> |
|             | Derived Products                            | Backscatter available for all 60 m multibeam Geomorphology products – see website | <a href="#">Pacific Islands Benthic Habitat Mapping Center</a> |

The land and seafloor area surrounding the islands of the Marianas as well as primary data coverage are reproduced from Miller et al. (2011) in Figure 41.



**Figure 41. CNMI Land and Seafloor Area and Primary Data Coverage**

### 2.6.2.2 Benthic Habitat

Juvenile and adult life stages of coral reef MUS and crustaceans including spiny and slipper lobsters and Kona crab extends from the shoreline to the 100 m isobath (64 FR 19067, April 19, 1999). All benthic habitat is considered EFH for crustaceans species (64 FR 19067, April 19, 1999), while the type of bottom habitat varies by family for coral reef species (69 FR 8336, February 24, 2004). Juvenile and adult bottomfish EFH extends from the shoreline to the 400 m isobath (64 FR 19067, April 19, 1999), and juvenile and adult deepwater shrimp habitat extends from the 300 m isobath to the 700 m isobath (73 FR 70603, November 21, 2008).

#### 2.6.2.2.1 RAMP Indicators

Benthic percent cover of coral, macroalgae, and crustose coralline algae from CRED are found in the following tables. CRED uses the benthic towed-diver survey method to monitor changes in benthic composition. In this method, “a pair of scuba divers (one collecting fish data, the other collecting benthic data) is towed about 1 m above the reef roughly 60 m behind a small boat at a constant speed of about 1.5 kt. Each diver maneuvers a towboard platform, which is connected to the boat by a bridle and towline and outfitted with a communications telegraph and various survey equipment, including a downward-facing digital SLR camera (Canon EOS 50D, Canon Inc., Tokyo). The benthic towed diver records general habitat complexity and type (e.g., spur and groove, pavement), percent cover by functional-group (hard corals, stressed corals, soft corals, macroalgae, crustose coralline algae, sand, and rubble), and for macroinvertebrates (crown-of-thorns sea stars, sea cucumbers, free and boring urchins, and giant clams).

Towed-diver surveys are typically 50 minutes long and cover about two to three kilometers of habitat. Each survey is divided into five-minute segments, with data recorded separately per segment to allow for later location of observations within the ~200-300 meter length of each segment. Throughout each survey, latitude and longitude of the survey track are recorded on the

small boat using a GPS; after the survey, diver tracks are generated with the GPS data and a layback algorithm that accounts for position of the diver relative to the boat. (PIFSC, 2016).

**Table 72. Mean percent cover of live coral from RAMP sites collected from towed-diver surveys in the Mariana Archipelago**

| Year                | 2003  | 2005  | 2007  | 2009  | 2011  | 2014  |
|---------------------|-------|-------|-------|-------|-------|-------|
| Agrihan             | 16.03 | 15.45 | 13.68 | 16.03 | 19.83 |       |
| Aguijan             | 17.88 | 17.25 | 11.68 | 15.61 | 21.88 | 33.46 |
| Alamagan            | 18.23 | 17.39 | 22.21 | 23.34 | 30.28 | 27.58 |
| Anatahan            | 7.93  |       |       |       |       |       |
| Arakane             | 24.06 | 11.83 |       |       |       |       |
| Asuncion            | 18.15 | 15.58 | 15.66 | 18.57 | 28    | 40.56 |
| Farallon de Pajaros | 10.13 | 4.82  | 4.94  | 11.28 | 11.69 | 16.45 |
| Guam                | 19.58 | 23.3  | 11.72 | 13.71 | 19.06 | 17.58 |
| Guguan              | 23    | 10.18 | 26.58 | 24.97 | 30.23 | 37.23 |
| Maug                | 26.86 | 21.43 | 26.25 | 28.09 | 38    | 46.17 |
| Pagan               | 18.51 | 9.84  | 12.04 | 13.09 | 16.23 | 27.87 |
| Pathfinder          | 24.17 | 24.75 |       |       |       |       |
| Rota                | 8.98  | 6.04  | 4.36  | 4.45  | 9.94  | 17.39 |
| Saipan              | 20.85 | 10.63 | 10.18 | 10.18 | 13.73 | 24.99 |
| Santa Rosa          | 7.31  | 7.8   |       |       |       |       |
| Sarigan             | 18.02 | 12.88 | 14.21 | 23.37 | 18.01 | 31.98 |
| Stingray            | 54.86 |       |       |       |       |       |
| Supply              | 38.75 |       |       |       |       |       |
| Tatsumi             | 7.92  |       |       |       |       |       |
| Tinian              | 12.46 | 8.99  | 8.08  | 9.33  | 12.02 | 17.37 |

**Table 73. Mean percent cover of macroalgae from RAMP sites collected from towed-diver surveys in the Mariana Archipelago**

| Year                | 2003  | 2005  | 2007  | 2009  | 2011  | 2014  |
|---------------------|-------|-------|-------|-------|-------|-------|
| Agrihan             | 48.25 | 22.65 | 8.55  | 3.2   | 4.63  |       |
| Aguijan             | 44.56 | 38.81 | 28.31 | 20.8  | 21.52 | 25.1  |
| Alamagan            | 41.21 | 26.03 | 15.65 | 15.47 | 12.81 | 8.33  |
| Anatahan            | 14.31 |       |       |       |       |       |
| Arakane             | 52.26 | 45.75 |       |       |       |       |
| Asuncion            | 51.1  | 5.37  | 19.11 | 7.54  | 7.47  | 3.86  |
| Farallon de Pajaros | 60.2  | 4.32  | 3.38  | 0.05  | 0.91  | 0.18  |
| Guam                | 46.19 | 52.67 | 43.22 | 26.82 | 29.61 | 41.64 |
| Guguan              | 45    | 10.18 | 19.5  | 17    | 12.59 | 8.66  |
| Maug                | 45.91 | 27.2  | 8.17  | 3.26  | 4.37  | 12.01 |
| Pagan               | 45.96 | 18.4  | 16.74 | 9.84  | 7.36  | 19.3  |

|            |       |       |       |       |       |       |
|------------|-------|-------|-------|-------|-------|-------|
| Pathfinder | 37.29 | 29    |       |       |       |       |
| Rota       | 54.34 | 56.05 | 38.76 | 30.95 | 35.16 | 29.33 |
| Saipan     | 48.57 | 30.75 | 31.87 | 20.39 | 15.26 | 25.18 |
| Santa Rosa | 42.5  | 70.54 |       |       |       |       |
| Sarigan    | 42.23 | 23.95 | 16.47 | 12.51 | 9.41  | 11.55 |
| Stingray   | 33.89 |       |       |       |       |       |
| Supply     | 19.17 |       |       |       |       |       |
| Tatsumi    | 67.22 |       |       |       |       |       |
| Tinian     | 46.94 | 56.38 | 39.95 | 30.4  | 25.92 | 34.91 |

**Table 74. Mean percent cover of crustose coralline algae from RAMP sites collected from towed-diver surveys in the Mariana Archipelago**

| Year                | 2003  | 2005  | 2007  | 2009  | 2011  | 2014  |
|---------------------|-------|-------|-------|-------|-------|-------|
| Agrihan             | 8.64  | 5.7   | 9.94  | 5.57  | 3.91  |       |
| Aguijan             | 14.69 | 10.59 | 12.67 | 7.32  | 11.47 | 18.33 |
| Alamagan            | 7.63  | 4.85  | 10.29 | 5.33  | 4.29  | 6.25  |
| Anatahan            | 7.72  |       |       |       |       |       |
| Arakane             | 5.28  | 3.58  |       |       |       |       |
| Asuncion            | 7.96  | 8.99  | 9.53  | 3.67  | 4.62  | 2.19  |
| Farallon de Pajaros | 3.44  | 8.03  | 5.39  | 2.94  | 2.29  | 0.05  |
| Guam                | 12.75 | 4.04  | 8.54  | 6.13  | 9.39  | 6.9   |
| Guguan              | 17.13 | 15    | 12.95 | 14.59 | 7.35  | 9.91  |
| Maug                | 10.22 | 7.53  | 12.32 | 7.73  | 5.38  | 8.23  |
| Pagan               | 6.61  | 12.41 | 14.16 | 8.42  | 6.33  | 2.48  |
| Pathfinder          | 5.56  | 10    |       |       |       |       |
| Rota                | 18.39 | 4.56  | 12.42 | 5.22  | 6.67  | 5.49  |
| Saipan              | 10.04 | 8.74  | 15.03 | 8.27  | 6.31  | 5.61  |
| Santa Rosa          | 7.13  | 0.55  |       |       |       |       |
| Sarigan             | 10.64 | 3.24  | 7.58  | 3.84  | 2.59  | 4.57  |
| Stingray            | 1.54  |       |       |       |       |       |
| Supply              | 35    |       |       |       |       |       |
| Tatsumi             | 6.11  |       |       |       |       |       |
| Tinian              | 6.25  | 5.18  | 16.16 | 4.07  | 7.59  | 5.96  |

### 2.6.2.3 Oceanography and Water Quality

The water column is also designated as EFH for selected MUS life stages at various depths. For larval stages of all species except deepwater shrimp, the water column is EFH from the shoreline to the EEZ. Coral reef species egg and larval EFH is to a depth of 100 m; crustaceans, 150m; and bottomfish, 400 m. Please see the Ecosystem and Climate Change section for information related to oceanography and water quality.



### 2.6.3 Report on Review of EFH Information

One EFH review was drafted this year; the review of the biological components of crustaceans EFH can be found in Appendix C.

### 2.6.4 EFH Levels

NMFS guidelines codified at 50 C.F.R. § 600.815 recommend Councils organize data used to describe and identify EFH into the following four levels:

Level 1: Distribution data are available for some or all portions of the geographic range of the species.

Level 2: Habitat-related densities of the species are available.

Level 3: Growth, reproduction, or survival rates within habitats are available.

Level 4: Production rates by habitat are available.

The Council adopted a fifth level, denoted Level 0, for situations in which there is no information available about the geographic extent of a particular managed species' life stage. The existing level of data for individual MUS in each fishery are presented in tables per fishery. In subsequent SAFE reports, each fishery section will include the description of EFH method, method used to assess the value of the habitat to the species, description of data sources used if there was analysis; and description of method for analysis.

#### 2.6.4.1 Precious Corals

Essential Fish Habitat for precious corals was originally designated in Amendment 4 to the Precious Corals Fishery Management Plan (64 FR 19067, April 19, 1999), using the level of data found in the table.

**Table 75. Level of EFH information available for the Western Pacific PCMUS**

| Species  | Pelagic Phase<br>(Larval Stage) | Benthic Phase | Source(s)                               |
|--|---------------------------------|---------------|---|
| <b>Pink Coral (<i>Corallium</i>)</b>                               |                                 |               |   |
| <i>Pleurocorallium secundum</i> (prev. <i>Corallium secundum</i> ) | 0                               | 1             | Figueroa and Baco (2014); HURL Database |
| <i>C. regale</i>   | 0                               | 1             | HURL Database                           |
| <i>Hemicorallium laauense</i> (prev. <i>C. laauense</i> )          | 0                               | 1             | HURL Database                           |
| <b>Gold Coral</b>  |                                 |               |   |
| <i>Kulamanamana haumea</i>   | 0                               | 1             | Sinniger et al. (2013); HURL Database   |
| <i>Callogorgia gilberti</i>  | 0                               | 1             | HURL Database                           |
| <i>Narella</i> spp.  | 0                               | 1             | HURL Database                           |
| <b>Bamboo Coral</b>  |                                 |               |   |
| <i>Lepidisis olapa</i>   | 0                               | 1             | HURL Database                           |

| Species  | Pelagic Phase (Larval Stage) | Benthic Phase | Source(s)                     |
|--|------------------------------|---------------|-------------------------------|
| <i>Acanella</i> spp.   | 0                            | 1             | HURL Database                 |
| <b>Black Coral</b>   |                              |               |                               |
| <i>Antipathes griggsi</i> (prev. <i>Antipathes dichotoma</i> ) | 0                            | 2             | Opresko (2009); HURL Database |
| <i>A. grandis</i>  | 0                            | 1             | HURL Database                 |
| <i>Mysiopathes ulex</i> (prev. <i>A. ulex</i> )                | 0                            | 1             | Opresko (2009); HURL Database |

#### 2.6.4.2 Bottomfish and Seamount Groundfish

Essential Fish Habitat for bottomfish and seamount groundfish was originally designated in Amendment 6 to the Bottomfish and Seamount Groundfish FMP (64 FR 19067, April 19, 1999).

**Table 76. Level of EFH information available for Western Pacific BMUS and seamount groundfish MUS complexes**

| Life History Stage                                   | Eggs | Larvae | Juvenile | Adult |
|--|------|--------|----------|-------|
| Bottomfish: (scientific/English common)              |      |        |          |       |
| <i>Aphareus rutilans</i> (red snapper/silvermouth)   | 0    | 0      | 0        | 2     |
| <i>Aprion virescens</i> (gray snapper/jobfish)       | 0    | 0      | 1        | 2     |
| <i>Caranx ignobilis</i> (giant trevally/jack)        | 0    | 0      | 1        | 2     |
| <i>C. lugubris</i> (black trevally/jack)             | 0    | 0      | 0        | 2     |
| <i>Epinephelus faciatus</i> (blacktip grouper)       | 0    | 0      | 0        | 1     |
| <i>E. quernus</i> (sea bass)                         | 0    | 0      | 1        | 2     |
| <i>Etelis carbunculus</i> (red snapper)              | 0    | 0      | 1        | 2     |
| <i>E. coruscans</i> (red snapper)                    | 0    | 0      | 1        | 2     |
| <i>Lethrinus amboinensis</i> (ambon emperor)         | 0    | 0      | 0        | 1     |
| <i>L. rubrioperculatus</i> (redgill emperor)         | 0    | 0      | 0        | 1     |
| <i>Lutjanus kasmira</i> (blueline snapper)           | 0    | 0      | 1        | 1     |
| <i>Pristipomoides auricilla</i> (yellowtail snapper) | 0    | 0      | 0        | 2     |
| <i>P. filamentosus</i> (pink snapper)                | 0    | 0      | 1        | 2     |
| <i>P. flavipinnis</i> (yelloweye snapper)            | 0    | 0      | 0        | 2     |
| <i>P. seiboldi</i> (pink snapper)                    | 0    | 0      | 1        | 2     |
| <i>P. zonatus</i> (snapper)                          | 0    | 0      | 0        | 2     |
| <i>Pseudocaranx dentex</i> (thicklip trevally)       | 0    | 0      | 1        | 2     |
| <i>Seriola dumerili</i> (amberjack)                  | 0    | 0      | 0        | 2     |
| <i>Variola louti</i> (lunartail grouper)             | 0    | 0      | 0        | 2     |
|  |      |        |          |       |
| Seamount Groundfish:                                 |      |        |          |       |
| <i>Beryx splendens</i> (alfonsin)                    | 0    | 1      | 2        | 2     |
| <i>Hyperoglyphe japonica</i> (ratfish/butterfish)    | 0    | 0      | 0        | 1     |
| <i>Pseudopentaceros richardsoni</i> (armorhead)      | 0    | 1      | 1        | 3     |

### 2.6.4.3 Crustaceans

Essential Fish Habitat for crustaceans MUS was originally designated in Amendment 10 to the Crustaceans FMP (64 FR 19067, April 19, 1999). EFH definitions were also approved for deepwater shrimp through an amendment to the Crustaceans FMP in 2008 (73 FR 70603, November 21, 2008).

**Table 77. Level of EFH information available for the Western Pacific crustacean MUS**

| Life History Stage  | Eggs | Larvae | Juvenile | Adult |
|---|------|--------|----------|-------|
| Crustaceans: (English common\scientific)                  |      |        |          |       |
| Spiny lobster ( <i>Panulirus marginatus</i> )             | 2    | 1      | 1-2      | 2-3   |
| Spiny lobster ( <i>Panulirus pencillatus</i> )            | 1    | 1      | 1        | 2     |
|   |      |        |          |       |
| Common slipper lobster ( <i>Scyllarides squammosus</i> )  | 2    | 1      | 1        | 2-3   |
| Ridgeback slipper lobster ( <i>Scyllarides haanii</i> )   | 2    | 0      | 1        | 2-3   |
| Chinese slipper lobster ( <i>Parribacus antarcticus</i> ) | 2    | 0      | 1        | 2-3   |
|   |      |        |          |       |
| Kona crab ( <i>Ranina ranina</i> )                        | 1    | 0      | 1        | 1-2   |

### 2.6.4.4 Coral Reef

Essential Fish Habitat for coral reef ecosystem species was originally designated in the Coral Reef Ecosystem FMP (69 FR 8336, February 24, 2004). An EFH review of CREMUS will not be undertaken until the Council completes its process of re-designating certain CREMUS into the ecosystem component classification. Ecosystem component species do not require EFH designations, as they are not a managed species.

### 2.6.5 Research and Information Needs

Based, in part, on the information provided in the tables above the Council identified the following scientific data which are needed to more effectively address the EFH provisions:

#### 2.6.5.1 All FMP Fisheries

- Distribution of early life history stages (eggs and larvae) of management unit species by habitat.
- Juvenile habitat (including physical, chemical, and biological features that determine suitable juvenile habitat).
- Food habits (feeding depth, major prey species etc.).
- Habitat-related densities for all MUS life history stages.
- Growth, reproduction, and survival rates for MUS within habitats.

#### 2.6.5.2 Bottomfish Fishery

- Inventory of marine habitats in the EEZ of the Western Pacific region.
- Data to obtain a better SPR estimate for American Samoa's bottomfish complex.
- Baseline (virgin stock) parameters (CPUE, percent immature) for the Guam/NMI deep-water and shallow-water bottomfish complexes.

- High resolution maps of bottom topography/currents/water masses/primary productivity.
- Habitat utilization patterns for different life history stages and species.

### 2.6.5.3 Crustaceans Fishery

- Identification of post-larval settlement habitat of all CMUS.
- Identification of “source/sink” relationships in the NWHI and other regions (i.e. relationships between spawning sites settlement using circulation models, genetic techniques, etc.).
- Establish baseline parameters (CPUE) for the Guam/Northern Marianas crustacean populations.
- Research to determine habitat-related densities for all CMUS life history stages in American Samoa, Guam, Hawaii, and CNMI.
- High resolution mapping of bottom topography, bathymetry, currents, substrate types, algal beds, and habitat relief.

### 2.6.5.4 Precious Corals Fishery

- Distribution, abundance, and status of precious corals in the CNMI and Guam.

## 2.6.6 References

64 FR 19067. Fisheries Off West Coast States and in the Western Pacific; Pelagic Fisheries, Amendment 8; Crustacean Fisheries, Amendment 10; Bottomfish and Seamount Groundfish Fisheries, Amendment 6; Precious Corals Fisheries, Amendment 4, Rule. *Federal Register* 64 (19 April 1999): 19067-19069. Downloaded from <https://www.govinfo.gov/content/pkg/FR-1999-04-19/pdf/99-9728.pdf>.

69 FR 8336. Fisheries off West Coast States and in the Western Pacific; Coral Reef Ecosystems Fishery Management Plan for the Western Pacific, Final Rule. *Federal Register* 69 (24 February 2004): 8336-8349. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.

73 FR 70603. Fisheries in the Western Pacific; Crustacean Fisheries; Deepwater Shrimp, Final Rule. *Federal Register* 73 (21 November 2008): 70603-70605. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2008-11-21/pdf/E8-27773.pdf>.

Brainard, R., Asher, J., Gove, J., Helyer, J., Kenyon, J., Mancini, F., Miller, J., Myhre, S., Nadon, M., Rooney, J., Schroeder, R., Smith, E., Vargas-Angel, B., Vogt, S., Vroom, P., Balwani, S., Craig, P., DesRochers, A., Ferguson, S., Hoeke, R., Lammers, M., Lundblad, E., Maragos, J., Moffitt, R., Timmers, M., and O. Vetter, 2008. Coral reef ecosystem monitoring report for American Samoa: 2002-2006. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 472 pp.

DesRochers, A., 2016. “Benthic Habitat Mapping.” NOAA Fisheries Center, Honolulu, HI. Presentation. Presented 6 April 2016.

- Figueroa, D.F. and A.R. Baco, 2014. Complete mitochondrial genomes elucidate phylogenetic relationships of the deep-sea octocoral families Coralliidae and Paragorgiidae. *Deep Sea Research Part II: Topical Studies in Oceanography*, 99, pp.83-91.
- Hawaii Mapping Research Group, 2014. Main Hawaiian Islands Multibeam Bathymetry and Backscatter Synthesis. School of Ocean and Earth Science and Technology, University of Hawaii at Mānoa. <http://www.soest.hawaii.edu/HMRG/multibeam/index.php>. Updated 20 November 2014. Accessed April 4, 2016.
- McCoy, K., Heenan, A., Asher, J., Ayotte, P., Gorospe, K., Gray, A., Lino, K., Zamzowm J., and I. Williams, 2017. Pacific Reef Assessment and Monitoring Program Data Report: Ecological monitoring 2016 - reef fishes and benthic habitats of the main Hawaiian Islands, Northwestern Hawaiian Islands, Pacific Remote Island Areas, and American Samoa. NOAA Pacific Islands Fisheries Science Center. PIFSC Data Report DR-17-001. 66 pp.
- Miller, J., Battista, T., Pritchett, A., Rohmann, S., and J. Rooney, 2011. Coral Reef Conservation Program Mapping Achievements and Unmet Needs. NOAA Coral Reef Conservation Program, Silver Spring, MD, 68 pp.
- Minton, D. 2017. Non-fishing effects that may adversely affect essential fish habitat in the Pacific Islands region, Final Report. NOAA National Marine Fisheries Service, Contract AB-133F-15-CQ-0014. 207 pp.
- Opresko, D.M., 2009. A New Name for the Hawaiian Antipatharian Coral Formerly Known as *Antipathes dichotoma* (Cnidaria: Anthozoa: Antipatharia) 1. *Pacific Science*, 63(2), pp.277-292.
- PIFSC, 2010. Coral reef ecosystems of the Mariana Archipelago: a 2003–2007 overview. NOAA Pacific Islands Fisheries Science Center, PIFSC Special Publication, SP-10-002, 38 pp.
- PIFSC, 2011. Coral reef ecosystems of American Samoa: a 2002–2010 overview. NOAA Fisheries Pacific Islands Fisheries Science Center, PIFSC Special Publication, SP-11-02, 48 pp.
- PIFSC, 2016. Pacific Islands Fisheries Science Center Ecosystem Sciences Coral Reef Ecosystem Survey Methods. Benthic Monitoring. [http://www.pifsc.noaa.gov/cred/survey\\_methods.php](http://www.pifsc.noaa.gov/cred/survey_methods.php). Updated 1 April 2016. Accessed 5 April 2016.
- Ryan, W.B.F., S.M. Carbotte, S.M., Coplan, J.O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., and R. Zemsky, 2009. Global Multi-Resolution Topography synthesis, *Geochem. Geophys. Geosyst.*, 10, Q03014. doi: 10.1029/2008GC002332
- Sinniger, F., Ocana, O.V., and A.R. Baco, 2013. Diversity of zoanths (Anthozoa: Hexacorallia) on Hawaiian seamounts: description of the Hawaiian gold coral and additional zoanths. *PLoS one*, 8(1), p.e52607.

## **2.7 MARINE PLANNING**

### **2.7.1 Introduction**

Marine planning is a science-based management tool being utilized regionally, nationally and globally to identify and address issues of multiple human uses, ecosystem health and cumulative impacts in the coastal and ocean environment. The Council's efforts to formalize incorporation of marine planning in its actions began in response to Executive Order (EO) 13547, *Stewardship of the Ocean, Our Coasts, and the Great Lakes*. EO 13158, *Marine Protected Areas*, proposes that agencies strengthen the management, protection, and conservation of existing MPAs, develop a national system of MPAs representing diverse ecosystems, and avoid causing harm to MPAs through federal activities. MPAs, or marine managed areas (MMAs) are one tool used in fisheries management and marine planning.

At its 165<sup>th</sup> meeting in March 2016, in Honolulu, Hawai'i, the Council approved the following objective for the FEPs: Consider the Implications of Spatial Management Arrangements in Council Decision-making. The following sub-objectives apply:

- Identify and prioritize research that examines the positive and negative consequences of areas that restrict or prohibit fishing to fisheries, fishery ecosystems, and fishermen, such as the Bottomfish Fishing Restricted Areas (BRFAs), military installations, NWHI restrictions, and Marine Life Conservation Districts (MLCDs).
- Establish effective spatially-based fishing zones.
- Consider modifying or removing spatial-based fishing restrictions that are no longer necessary or effective in meeting their management objectives.
- As needed, periodically evaluate the management effectiveness of existing spatial-based fishing zones in Federal waters.

In order to monitor implementation of this objective, this annual report includes the Council's spatially-based fishing restrictions or MMAs, the goals associated with those, and the most recent evaluation. Council research needs are identified and prioritized through the Five Year Research Priorities and other processes, and are not tracked in this report.

In order to meet the EFH and National Environmental Policy Act (NEPA) mandates, this annual report tracks activities that occur in the ocean that are of interest to the Council, and incidents or facilities that may contribute to cumulative impact. NMFS is responsible for NEPA compliance, and the Council must assess the environmental effects of ocean activities for the FEP's EFH cumulative impacts section.


### **2.7.2 Response to Previous Council Recommendations**

There are no Council recommendations indicating review deadlines for Marianas MMAs.

### **2.7.3 Marine Managed Areas Established under FEPs**

Council-established MMAs were compiled in Table 78 from 50 CFR § 665, Western Pacific Fisheries, the Federal Register, and Council amendment documents. All regulated fishing areas

and large scale access restrictions, including the Mariana Trench Marine National Monument, are shown in Figure 42.



\* The Coral Reef Ecosystem Regulatory Area excluded the portion of EEZ waters 0-3 miles around the CNMI. The Bottomfish Management Subarea was divided in the CNMI Inshore Area, which was that portion of the EEZ shoreward of 3 nautical miles of the shoreline of CNMI, and the CNMI Offshore Area, which was that portion of the EEZ seaward of 3 nautical miles from the CNMI shoreline.

Figure 42. Regulated fishing areas of the Mariana Archipelago

Table 78. MMAs established under FEPs from [50 CFR § 665](#)

| Name                              | FEP                 | Island | 50 CFR /FR /Amendment Reference   | Marine Area (km <sup>2</sup> ) | Fishing Restriction  | Goals   | Most Recent Evaluation | Review Deadline |
|-----------------------------------|---------------------|--------|---|--------------------------------|--|---|------------------------|-----------------|
| Pelagic Restrictions              |                     |        |   |                                |  |   |                        |                 |
| Guam Longline Prohibited Area     | Pelagic             | Guam   | 665.806(a)(3)<br><a href="#">57 FR 7661 Pelagic FMP Am. 5</a>                     | 50,192.88                      | Longline fishing prohibited  | Prevent gear conflicts between longline vessels and troll/handline vessels.   | 1992                   | -               |
| CNMI Longline Prohibited Area     | Pelagic             | -      | 665.806(a)(4)<br><a href="#">76 FR 37287</a>                                      | 88,112.68                      | Longline fishing prohibited  | Reduce potential for nearshore localized fish depletion from longline fishing, and to limit catch competition and gear conflicts between the CNMI-based longline and trolling fleets. | 2011                   | -               |
| Bottomfish Restrictions           |                     |        |   |                                |  |   |                        |                 |
| Guam Large Vessel Prohibited Area | Mariana Archipelago | Guam   | 665.403(a)<br><a href="#">71 FR 64474</a><br>Bottomfish FMP Am. 9                 | 29,384.06                      | Vessels ≥ 50 feet prohibited   | To maintain viable participation and bottomfish catch rates by small vessels in the fishery.  | 2006                   | -               |
| Other Restrictions                |                     |        |   |                                |  |   |                        |                 |
| Guam No Anchor Zone               | Mariana Archipelago | Guam   | 665.399<br><a href="#">69 FR 8336</a><br><a href="#">Coral Reef Ecosystem FEP</a> | 138,992.51                     | Anchoring by all fishing vessels ≥ 50 ft. prohibited on the offshore southern banks located in the U.S. EEZ off Guam | Minimize adverse human impacts on coral reef resources.   | 2004                   | -               |



## 2.7.4 Fishing Activities and Facilities

There are no proposed or existing offshore aquaculture projects in Federal waters of neither Guam nor CNMI.

## 2.7.5 Non-Fishing Activities and Facilities

The following section includes activities or facilities associated with known uses and predicted future uses. The Plan Team will add to this section as new facilities are proposed and/or built. Due to the sheer volume of ocean activities and the annual frequency of this report, only major activities on multi-year planning cycles are tracked in this report. Activities which are no longer reasonably foreseeable or have been replaced with another planning activity are removed from the report, though may occur in previous reports.

### 2.7.5.1 Alternative Energy Facilities

There are no proposed or existing alternative energy facilities in Federal waters of neither Guam nor CNMI.

### 2.7.5.2 Military Training and Testing Activities and Impacts

The Department of Defense major planning activities in the region are summarized in Table 79. Activities that are no longer reasonably foreseeable or have been replaced with another planning activity were removed from the report, though may occur in previous reports.

## 2.7.6 Pacific Islands Regional Planning Body Report

In June 2018, President Trump signed the EO 13840 *Regarding the Ocean Policy to Advance Economic, Security, and Environmental Interests of the United States*, which revoked EO 13547. The new EO eliminated the mandate for the federal government to participate in ocean planning at a regional level and eliminated the regional planning bodies. As such, the Pacific Islands Regional Planning Body (RPB) no longer exists and ocean planning will now occur at a local level led by Hawaii and the territories.

However, EO 13840 established a policy focused on public access to marine data and information, and requires federal agencies to 1) coordinate activities regarding ocean-related matters and 2) facilitate the coordination and collaboration of ocean-related matters with governments and ocean stakeholders. To that end, the [American Samoa Coastal and Marine Spatial Planning Data Portal](https://marinecadastre.gov/viewers/) was created by Marine Cadastre (<https://marinecadastre.gov/viewers/>). The intent is for it to be expanded to include the Marianas, PRIA, and Hawaii and be titled the Pacific Islands Regional Marine Planner.

Spatial planning has occurred in CNMI in Saipan Lagoon. CNMI Division of Coastal Resources Management developed the Saipan Lagoon Use Management Plan, which was updated in 2017 and can be found at <https://dcrm.gov.mp/current-projects/saipan-lagoon-use-management-planning/>. There is also an associated mapping tool that can be found at <http://dcrm.maps.arcgis.com/apps/webappviewer/index.html?id=7def562d70014be58112bc62b1bf9902>.

**Table 79. Department of Defense major planning activities**

| Action  | Description  | Phase   | Impacts   |
|---|--|---|---|
| <a href="#">Guam and CNMI Military Relocation SEIS</a>              | Relocate Marines to Guam and build a cantonment/family housing unit on Finegayan/AAFB, a live-fire individual training range complex at the Ritidian Unit of the Guam National Wildlife Refuge.  | ROD published August 29, 2015.<br><br>Suit filed for segmentation and range of reasonable alternatives under NEPA, requesting that DON vacate the ROD. DOJ asked US District Court for the NMI to dismiss the plaintiff's complaint with prejudice to prevent refiling ( <a href="http://www.saipantribune.com/index.php/doj-federal-court-lacks-jurisdiction/">http://www.saipantribune.com/index.php/doj-federal-court-lacks-jurisdiction/</a> ). | Surface danger zone established at Ritidian – access restricted during training. Access will be negotiated between the Navy and USFWS.<br><br>Northern District Wastewater Treatment Plant is non-compliant with NPDES permit; until plant is upgraded, increased wastewater discharge associated with buildup will significantly impact nearshore water quality. DOD to fund plant upgrades – see Economic Adjustment Committee Implementation Plan. |
| <a href="#">Mariana Islands Training and Testing – Supplemental</a> | The supplement to the 2015 Final EIS/OEIS is being prepared to support ongoing and future activities conducted at sea and on Farallon de Medinilla (FDM) beyond 2020. New information, including an updated acoustic effects model, updated marine mammal density data, and evolving and emergent BSIA, will be used to update the MITT. | The 2019 MITT Final Supplemental EIS/OEIS is expected in spring 2020.<br><br>Public Comment and Open House Public Meetings to take place in March and April of 2019.  | Likely access and habitat impacts similar to previous analysis.   |

|  |   |   |   |
|--|---|---|---|
| <a href="#">CNMI Joint Military Training</a>                         | Establish unit and combined level training ranges on Tinian and Pagan.  | Supplemental Draft EIS expected in late 2018 or early 2019.<br><br>Suit filed for segmentation and range of reasonable alternatives under NEPA. DOJ asked US District Court for the NMI to dismiss the plaintiff's complaint with prejudice to prevent refileing.   | Significant access and habitat impacts.   |
| <a href="#">Divert Activities and Exercises, Air Force, Marianas</a> | Improve airports in CNMI for expanding mission requirements in Western Pacific.   | The USAF has published a NOI to prepare a SEIS for the proposed Tinian Divert Infrastructure Improvements. The NOI began the public scoping process for the SEIS, which ended on May 31, 2018. Substantive comments received during the public scoping period will be taken into consideration during preparation of the Draft SEIS.  | Adverse impacts to EFH minimal; access near Port of Tinian fuel transfer facility affected. |
| Garapan Anchorage  | Military Pre-Positioned Ships anchor and transit.   | Expired Memorandum of Understanding with the CNMI government. As of October 2018, MOU had not been signed.  | Access, invasive species, unmitigated damage to reefs.                                      |
| Farallon de Medinilla  | Restricted airspace covering the island to 12 nmi radius to conduct military training scenarios using air-to-ground ordnance delivery, naval gunfire, lasers and special operations training. | Final rule published March 13, 2017, effective June 22, 2017, designating a new area, R-2701A, that surrounds existing R-2701, encompassing airspace between a 3 nmi radius and 12 nmi radius of FDM (82 FR 13389).<br><br>Proposed surface danger zone to 12 nmi.<br><br>Damage to submerged lands and fisheries to be included within consultation establishing continued US interest in the island and compensation to the CNMI (Report to the President on 902 Consultations, 2017) | Access – to fishing grounds and transit to fishing grounds - and damage to submerged lands. |

## 2.7.7 References

- 50 CFR 665. Fisheries in the Western Pacific. Title 50 *Code of Federal Regulations*, Pt. 665. Electronic Code of Federal Regulations data current as of March 16, 2016. Accessed at [http://www.ecfr.gov/cgi-bin/retrieveECFR?gp=&SID=b28abb7da3229173411daf43959fcbd1&n=50y13.0.1.1.2&r=PART&ty=HTML#\\_top](http://www.ecfr.gov/cgi-bin/retrieveECFR?gp=&SID=b28abb7da3229173411daf43959fcbd1&n=50y13.0.1.1.2&r=PART&ty=HTML#_top).
- 57 FR 7661. Pelagic Fisheries of the Western Pacific Region, Final Rule. *Federal Register* 57 (4 March 1992): 7661-7665. Downloaded from <http://www.wpcouncil.org/pelagic/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 69 FR 8336. Fisheries Off West Coast States and in the Western Pacific; Coral Reef Ecosystems Fishery Management Plan for the Western Pacific, Final Rule. *Federal Register* 69 (24 February 2004): 8336-8349. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 71 FR 64474. Fisheries in the Western Pacific; Western Pacific Bottomfish and Seamount Groundfish Fisheries; Guam Bottomfish Management Measures, Final Rule. *Federal Register* 71 (2 November 2006): 64474-64477. Downloaded from <http://www.wpcouncil.org/bottomfish/Documents/FMP/Bottomfish%20A9%20Final%20Rule%202006.pdf>.
- 76 FR 37287. Western Pacific Pelagic Fisheries; Prohibiting Longline Fishing Within 30 nm of the Northern Mariana Islands, Final Rule. *Federal Register* 76 (27 June 2011): 37287-37289. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2011-06-27/pdf/2011-16039.pdf>.
- 78 FR 32996. Western Pacific Fisheries; Fishing in the Marianas Trench, Pacific Remote Islands, and Rose Atoll Marine National Monuments, Final Rule. *Federal Register* 78 (3 June 2013): 32996-33007. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 80 FR 46252. Department of Defense; Department of the Navy. Record of Decision for the Mariana Islands Training and Testing Final Environmental Impact Statement/Overseas Environmental Impact Statement, Notice. *Federal Register* 80 (4 August 2015): 46252. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2015-08-04/pdf/2015-19050.pdf>.
- 80 FR 55838. Department of Defense; Department of the Navy. Record of Decision for the Final Supplemental Environmental Impact Statement for Guam and Commonwealth of the Northern Mariana Islands Military Relocation, Notice. *Federal Register* 80 (17 September 2015): 55838-55839. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2015-09-17/pdf/2015-23244.pdf>.
- 81 FR 92791. Department of Defense; United States Air Force. Record of Decision for Divert Activities and Exercises, Commonwealth of the Northern Mariana Islands, Notice.

- Federal Register* 81 (20 December 2016): 92791. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2016-12-20/pdf/2016-30488.pdf>.
- 82 FR 13389. Modification of Restricted Area R-7201; Farallon De Medinilla Island, Mariana Islands, Final Rule. *Federal Register* 82 (13 March 2017): 13389-13390. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2017-03-13/pdf/FR-2017-03-13.pdf>.
- CNMI Joint Military Training EIS/OEIS. "DOD to Issue Revise Draft EIS on CJMT". Department of Defense. Published 18 February, 2016. Accessed February 28, 2017. <http://www.cnmijointmilitarytrainingeis.com/announcements/25>.
- De La Torre, F. 2016. "DOJ: Federal court lacks jurisdiction." Saipan Tribune. Published 25 November 2016. <http://www.saipantribune.com/index.php/doj-federal-court-lacks-jurisdiction/>.
- Hofschneider, A. 2017. "The U.S. Military Won't Bomb Pagan Just Yet." Honolulu Civil Beat. Published 9 March 2017. [http://www.civilbeat.org/2017/03/the-u-s-military-wont-bomb-pagan-or-tinian-just-yet/?mc\\_cid=1a464a317d&mc\\_eid=abaf3b9d93](http://www.civilbeat.org/2017/03/the-u-s-military-wont-bomb-pagan-or-tinian-just-yet/?mc_cid=1a464a317d&mc_eid=abaf3b9d93).
- U.S. Department of Defense. 2015. Draft Environmental Impact Statement: Joint Military Training, Commonwealth of the Northern Mariana Islands. Accessed 10 March 2017. <https://cnmimarines.s3.amazonaws.com/static/DraftEIS/Cover%20thru%20Readers%20Guide.pdf>.
- U.S. Department of Defense, Department of the Air Force. 2016. Final environmental Impact Statement: Divert Activities and Exercises, Commonwealth of the Northern Mariana Islands. Accessed 17 March 2016. <http://www.pacafdivertmarianaseis.com/archive>.
- U.S. Department of the Interior. Report to the President on 902 Consultations. 2017. Special Representatives of the United States and the Commonwealth of the Northern Mariana Islands. January 2017. Accessed 12 March 2018. <https://www.doi.gov/sites/doi.gov/files/uploads/902-consultations-report-january-2017.pdf>.
- WPRFMC. 2015. Report of the CNMI Advisory Panel to the Western Pacific Regional Fishery Management Council, June 2015.
- WPRFMC. 2018. Fishery Management Plan and Fishery Ecosystem Plan Amendments. Available from <http://www.wpcouncil.org/fishery-plans-policies-reports/current-fishery-ecosystem-plans-sorted-by-island-areas/>.

### **3 DATA INTEGRATION**

#### **3.1 INTRODUCTION**

##### **3.1.1 Potential Indicators for Nearshore Fisheries**

The purpose of this section (“Chapter 3”) of the Stock Assessment and Fishery Evaluation (SAFE) annual report is to identify and evaluate potential fishery ecosystem relationships between fishery parameters and ecosystem variables to assess how changes in the ecosystem affect fisheries in the Mariana Archipelago and across the Western Pacific region. “Fishery ecosystem relationships” are those associations between various fishery-dependent data measures (e.g., catch, effort, or catch-per-unit-effort), and other environmental attributes (e.g., precipitation, sea surface temperature, primary productivity) that may contribute to observed trends or act as potential indicators of the status of prominent stocks in the fishery. These analyses represent a first step in a sequence of exploratory analyses that will be utilized to inform new assessments of what factors may be useful going forward.

To support the development of Chapter 3 of the annual SAFE report, staff from the Council, NMFS PIFSC and PIRO, and Triton Aquatics (consultants), held a SAFE Report Data Integration Workshop (hereafter, “the Workshop”) convened on November 30, 2016 to identify potential fishery ecosystem relationships relevant to local policy in the WPR and determine appropriate methods to analyze them. The archipelagic fisheries group developed nearly 30 potential fishery ecosystem relationships to examine across bottomfish, coral reef, and crustacean fisheries based on data reliability, suitability of methodology, repeatability on an annual basis, and how well analyses could potentially inform management decisions.

Brief introductory analyses, presented in this section and initially introduced in the 2017 report, were intended to be “proof of concept” such that similar evaluations could be carried out on remaining fishery data for the Mariana Archipelago in the future. However, the Archipelagic Fishery Ecosystem Plan Team determined that the quantitative analyses presented here were not sufficient to act as a model for future evaluations. Using the direction from the Plan Team, the data integration module was updated for the Hawaii Archipelagic Annual SAFE Report for the 2018 report, but each of the remaining archipelagic reports still contains data integration assessments from 2017. The Annual SAFE Report for the Mariana Archipelago will be updated in the following year similar to the Annual SAFE Report for the Hawaii Archipelago pending Plan Team support.

Going forward, relationships deemed potentially relevant will be emphasized and recommended for further analysis. In subsequent years, this chapter will be updated with these analyses through the SAFE report process as the strength of certain fishery ecosystem relationships relevant to advancing ecosystem-based fishery management are determined.

To begin, this chapter described feedback from the Plan Team, SSC, and Council members on the initial drafts of the data integration module. Next, the chapter includes brief descriptions of past work on fishery ecosystem relationship assessment in coral reefs of the U.S. Western Pacific, followed by initial evaluations of relationships previously recommended for evaluation by participants of the Workshop using current data streams from the Mariana Archipelago. The

evaluations completed were exploratory in nature, being the first step of analyses to know which comparisons may be more useful to focus on going forward.

Going forward with the analyses and presentation of results for the data integration chapter of the Marianas Archipelago Annual SAFE Report, the Plan Team suggested several improvements to implement in the coming year: standardizing and correcting values in CPUE time series, incorporating longer stretches of phase lag, completing comparisons on the species-level and by dominant gear types, incorporating local knowledge on shifts in fishing dynamics over the course of the time series, and utilizing the exact environmental data sets presented in the ecosystem consideration chapter of the annual report. Many of these recommendations were applied to datasets from Hawaii in 2018, and will similarly be done for Marianas data integration analyses in the upcoming report cycles. Implementation of these suggestions will allow for the preparation of a more finalized version of the data integration chapter in the coming report cycles.

### 3.1.2 2018 Recommendations for Chapter Development

At the most recent FEP Plan Team Meeting held on April 30<sup>th</sup> – May 1<sup>st</sup>, 2018, participants were presented preliminary data integration results shown here, and provided detailed recommendations to support the ongoing development of the data integration section of the Archipelagic Annual SAFE Report. These suggestions, both general and specific, will be implemented in the coming year to ensure that more refined analyses comprise the data integration section. FEP Plan Team participants recommended that:

- CPUE data should be standardized and calculated in a more robust fashion, measuring the average catch per unit effort rate over the course of a year to analyze variance.
- Analyses of fishery performance data against environmental variables should focus on dominant gear types rather than the entirety of the fishery or other gear aggregates (e.g. purse seine harvest of *Selar crumenophthalmus* in the MHI).
- There should be additional phase lag implemented in the analyses
- Local knowledge of fishery dynamics, especially pertaining to shifting gear preferences, should be utilized. Changes in dynamics that may have impacted observed fishery trends over the course of available time series, both discreetly and long-term for taxa-specific and general changes should be emphasized.
- Spatial specificity and precision should be increased for analyses of environmental variables in relation to areas commonly fished.

The analyses presented in the data integration chapter of the 2018 Hawaii Annual SAFE Report are a reflection of a thoughtful re-approaching to these data integration evaluations based on this feedback. Additional data can be added to either time series as they are made available. Incorporating such recommendations into the 2018 version of the Mariana Archipelago Annual SAFE Report will mark the beginning of a standardized process to implement current data integration analyses on an annual basis. Doing so will promote more proactive management action with respect to ecosystem-based fishery management objectives.

### 3.1.3 Past Work

Richards et al. (2012) performed a study on a range environmental factors that could potentially affect the distribution of large-bodied coral reef fish in Mariana Archipelago. Large-bodied reef fish were determined to typically be at the greatest risk of overfishing, and their distribution in the region was shown to be negatively associated with human population density. Additionally, depth, sea surface temperature (SST), and distance to deep water were identified as important environmental factors to large-bodied coral reef fish, whereas topographic complexity, benthic habitat structure, and benthic cover had little association with reef fish distribution in the Mariana Archipelago.

Kitiona et al. (2016) completed a study of the impacts climate and/or ecosystem change on coral reefs fish stocks of American Samoa using climate and oceanic indicators (see Section 2.5.4). The evaluation of environmental variables showed that certain climate parameters (e.g., SST anomaly, sea level height, precipitation, and tropical storm days) are likely linked to fishery performance. It was also noted that larger natural disturbances in recent decades, such as cyclones and tsunamis, negatively impacted reef fish assemblages and lowered reef fishery CPUE in American Samoa (Ochavillo et al., 2012).

On a larger spatial scale, an analysis of various drivers on coral reef fish populations across 37 U.S.-affiliated islands in the Central and Western Pacific was performed by Williams et al. (2015), and evaluated relationships between fish biomass in these reefs with human and environmental factors. Again, reef fish assemblages were negatively associated with increasing human population density (even at relatively low levels) across the WRP, but were positively associated with elevated levels of ocean productivity across islands. The authors warned, however, that the ability of reefs surrounding uninhabited islands to maintain fish populations varies, and that high biomass observed in remote areas (e.g. the NWHI) may not necessarily be reflective of baselines or recovery response levels for all reef systems.

A common method of EBFM used in coral reef ecosystems is the implementation of biological reference points, statistical indicators of potential overfishing used to help determine how a fishery is performing relative to these points at a given time (McClanahan et al., 2007). Hawhee (2007) adapted this idea, generating biological reference points in the form of CPUE-based proxies to be used as indicators for reef fish stocks in the WPR. However, the devised method was determined to be inappropriate for application in management of reef stocks in the U.S. Western Pacific due to the lack of a historical CPUE to use as a baseline for the reference points and their limit thresholds (Remington and Field, 2016).


## 3.2 PRECIPITATION

### 3.2.1 Guam

Participants of the Workshop determined that the potential fishery ecosystem relationships between precipitation levels and atulai and opelu (bigeye scad and mackerel scad, *Selar crumenophthalmus* and *Decapterus macarellus*, respectively) were among the highest priority of those involving coral reef fisheries in the Mariana Archipelago. It has been suggested that the recruitment of small tropical pelagic fish is related to annual rainfall and subsequent runoff



enrichment (Longhurst and Pauly, 1987). The direct freshwater and nutrient input to reefs associated with increased precipitation can alter the physiochemical composition of the water, and it has been shown that reef assemblages are positively associated with this sort of increased ocean productivity (Williams et al. 2015). Data for precipitation in the Mariana Archipelago was gathered from local databases maintained by the National Weather Service (NWS-G). The time series of total annual precipitation from showed a non-significant, slightly variable trend over the last 30 years ( $R^2 = 0.05$ ,  $CV = 19.5$ ; Figure 43).



**Figure 43. Total annual precipitation (in.) in Guam from 1980-2016**


### 3.2.1.1 Evaluating relationship with atulai

Total annual estimated atulai catch in the Guam recreational coral reef fishery according to shore- and boat-based creel surveys showed no general trend over the last thirty years, with relatively large variability likely due to several years of catch orders of magnitude greater than previous or subsequent years (e.g., 2009;  $R^2 = 0.01$ ;  $CV = 119.5$ ; Figure 44). Combined effort statistics between shore- and boat-based creel survey statistics could not be generated because the proxies used to measure effort in each survey are different (i.e. number of gear hours versus number of boat trips). Similarly, because effort could not be standardized across the data sets, CPUE could not be generated on the individual family level at which these evaluations are taking place.


Examining effort, Guam shore-based creel survey data show that there are considerable differences in the number of samples recorded across gear types. The most frequently sampled gear in the shore-based survey was hook and line by an order of magnitude, and had catch estimated to be several times greater than that in the expanded dataset (Figure 45a-b). Effort data also revealed that, despite catch statistics, the gill net had been sampled the least frequently among the top gears (Figure 45a-b). Boat-based effort data show that bottom fishing was sampled approximately twice as much than the other three top gears, but the difference in the expanded estimates between were at least an order of magnitude greater (Figure 45c-d).

Generally, each of the time series for prominent gear types in Guam showed a slight shift but seemingly no net change over the course of available data despite interannual variability.


Total estimated atulai catch and rainfall in Guam showed no statistical association with one another such that would allow for assessment of the fishery ecosystem relationship between the two ( $R^2=0.02$ ; Figure 46). However, there seemed to be a slight observable negative relationship between the two ( $r = -0.15$ ), indicating that catch may have experienced a minor decrease in years with more rainfall. Additionally, there was no association between annual rainfall amounts and total estimated atulai catch in Guam when only considering shore-based data, boat-based data, or prominent gear types.



**Figure 44. Time series of total annual estimated (i.e. expanded) landings of atulai in kilograms from Guam shore-and boat-based creel survey records from 1982-2016**



**Figure 45. Time series of total sampled (left) and expanded (right) effort for top gear types in shore-based (top) and boat-based (bottom) creel surveys in Guam from 1982-2016**




**Figure 46. Linear regression between total atulai catch (kg) in the Guam shore-based and boat-based creel survey records and total annual rainfall (in.) from 1982-2016**

### 3.2.1.2 Evaluating relationship with *D. macarellus*

*Decapterus macarellus* (i.e. mackerel scad) records from creel surveys in Guam were scant and had high variability, with estimated catch for many years being close to zero while others had close to 8,000 kg ( $R^2 = 0.01$ ;  $CV = 278.4$ ; Figure 47). Several years where mackerel scad catch

data were available, they indicated a total amount landed of just a few kilograms (e.g. 1999, 2001, 2013, etc.; Figure 47). Because there were 17 of 35 total years with available mackerel scad catch data across gear types for the entire territory since 1982, many with extremely low catch estimates, the time series were not able to be used for comparison to rainfall records in the same region over the last thirty years.




**Figure 47. Time series of total annual expanded landings of *Decapterus macarellus* (kg) in Guam shore-and boat-based creel survey records from 1982-2016**

In summary, no fishery ecosystem relationship could be established between atulai or mackerel scad catch with precipitation in Guam from 1982 till present without the incorporation of phase lag, and no standardized index/threshold characteristic of the association between the parameters could be identified representative of an immediate population response. The general lack of recreational harvest data for mackerel scad in Guam hindered the ability to determine whether a relationship exists with rainfall in that portion of the fishery. Analyses including atulai data had similar comparisons with rainfall data completed in the MHI as well, though no notable relationship between atulai catch and annual precipitation was identified there.

### 3.3 SEA SURFACE TEMPERATURE


Sea surface temperature (SST) is a commonly used diagnostic tool in monitoring climate change and its affects both regionally and globally, as it is representative of changes in ocean temperatures over time that can affect coastal fisheries (see Section 2.5.4). The potential influence of temperature-derived variables in fishery ecosystem relationships for U.S. Western Pacific coral reef stocks was deemed to be among the highest priority by the participants of the Workshop. Data for SST was gathered from the NOAA's AVHRR Pathfinder v5.0 through the OceanWatch program in the Central Pacific (NOAA/NESDIS/OceanWatch).

A time series of SST for the CNMI from 1985-2016 is shown in Figure 48. SST here had slightly less variability over time than Guam (CV = 0.55), again indicating relative stability. Unlike Guam, the CNMI did not seem to be observably increasing or decreasing over the time series of available data. The hottest temperature in the last three decades was approximately 29°C, where preceding SST had largely been stable over time. The average SST over the course of evaluated data was 28.8°C, slightly warmer than observed in Guam. The lowest recorded SST over the course of the time series was just about 27.5°C in the year 1996 (Figure 48).



**Figure 48. Time series of SST (°C) in the CNMI from 1985-2016 (CV = 0.55)**

A time series of SST for Guam from 1985-2016 is shown in Figure 49. Temperature had low variability over time (CV = 1.38), suggesting relative stability. There was also a seeming increase in temperature over the last three decades, with some of the hottest temperatures recorded observed in the last five years. The average SST over the course of evaluated data was 28.6°C. The highest recorded SST over the course of the time series was just over 29°C in the year 1999, whereas the lowest was earlier in the 1990s (27.7°C; Figure 49).



**Figure 49. Time series of SST (°C) from 1985-2016 in Guam (CV = 1.38)**

### 3.3.1 CNMI

#### 3.3.1.1 Evaluating relationship for entire reef fishery

A plot showing the relationship between SST and catch time series from the recreational coral reef fishery in the CNMI from 2000-2016 is depicted in Figure 50. Landings were variable over the course of the time series (CV = 19.4), but less so than observed in catch time series in Guam. Total annual catch in the fishery has been observably decreasing over the last decade and a half despite an abrupt increase in 2013 resulting in the recorded maximum catch over this period (~338,000 kg). Recent recorded catch levels (i.e. for 2016) were the lowest for the fishery through the available time series of data (~165,000 kg; Figure 50).

In performing comparisons between fishery parameters and environmental variables such as SST, data were grouped in taxa categories based on family due to scarcity of data on the species level in many cases. Table 80 displays the different dominant family groups considered as well as their common names.

Linear regressions and correlation analyses performed on the time series of recreational coral reef fishery catch (kg) and annual mean SST from the CNMI are reported in Table 81. The comparisons between the two parameters showed a negatively significant relationship between 2000 and 2016 ( $R^2 = 0.30$ ,  $p = 0.02$ ; Table 81; Figure 51). The relationship between the total annual catch and average annual SST for the whole fishery were associated such that for every degree Celsius of temperature increase, catch would decrease by approximately 105,000 kg (Figure 51).





Figure 50. Time series of total annual catch (kg; blue) for the CNMI recreational coral reef fishery plotted alongside average annual SST (°C; black) from 2000-2016

Table 80. Families in creel surveys from the U.S. Western Pacific analyzed in this report

| Four-letter code | Family        | Common Name              |
|------------------|---------------|--------------------------|
| LUTJ             | Lutjanidae    | snappers                 |
| LETH             | Lethrinidae   | emperors                 |
| CARA             | Carangidae    | jacks/mackerel/trevally  |
| ACAN             | Acanthuridae  | unicornfish/tang         |
| SERR             | Serranidae    | Sea bass/grouper         |
| SIGA             | Siganidae     | rabbitfish               |
| SCAR             | Scaridae      | parrotfish               |
| MULL             | Mullidae      | goatfish                 |
| MUGI             | Mugilidae     | mullet                   |
| LABR             | Labridae      | wrasse                   |
| HOLO             | Holocentridae | squirrelfish/soldierfish |
| BALI             | Balistidae    | triggerfish              |

Table 81. Correlation coefficients (*r*) between recreational coral reef fishery catch (kg) and SST (°C) in the CNMI for 12 top taxa harvested from 2000-2016

| Taxa Code             | Total Catch  | LUTJ | LETH  | CARA  | ACAN  | SERR  | SIGA | SCAR  | MULL  | MUGI | LABR | HOLO | BALI  |
|-----------------------|--------------|------|-------|-------|-------|-------|------|-------|-------|------|------|------|-------|
| <b>n = 17</b>         |              |      |       |       |       |       |      |       |       |      |      |      |       |
| <i>p</i>              | <b>0.02</b>  | 0.49 | 0.54  | 0.26  | 0.70  | 0.91  | 0.99 | 0.88  | 0.06  | -    | 0.59 | 0.91 | 0.82  |
| <i>r</i>              | <b>-0.55</b> | 0.18 | -0.16 | -0.29 | -0.10 | -0.03 | 0.00 | -0.04 | -0.47 | -    | 0.14 | 0.03 | -0.06 |
| <i>R</i> <sup>2</sup> | <b>0.30</b>  | 0.03 | 0.02  | 0.09  | 0.01  | 0.00  | 0.00 | 0.00  | 0.22  | -    | 0.02 | 0.00 | 0.00  |




**Figure 51. Linear regression showing the correlation between total annual catch (kg) in creel survey records and average annual SST (°C) in the CNMI from 2000-2016**

### 3.3.1.2 Evaluating relationship for dominant taxa

Correlation and regression analyses were performed on prominent taxa in the CNMI recreational coral reef fishery, and it was found that no individual taxa had significant relationships with SST data (Table 81). The strongest associations between fishery catch and SST were observed from the Mullids ( $R^2 = 0.22$ ,  $p = 0.06$ ; Figure 52a), Carangids ( $R^2 = 0.09$ ,  $p = 0.26$ ; Figure 52b), and Lutjanids ( $R^2 = 0.03$ ,  $p = 0.49$ ; Figure 52c). While the relationship between catch and temperature for families Mullidae and Carangidae were negative, the Lutjanidae family had a positive relationship (Table 81).






**Figure 52. Linear regressions showing the three top correlations between total annual catch (kg) from creel survey records and average annual SST (°C) in the CNMI from for (a) Mullids, (b) Carangids, and (c) Lutjanids from 2000–2016**

### 3.3.2 Guam

#### 3.3.2.1 Evaluating relationship for entire reef fishery

An individual plot depicting the comparisons of time series of SST and catch from the recreational coral reef fishery in Guam from 1985-2016 is shown in Figure 53. Landings were variable over the course of the time series (CV = 28.1) though relatively stable, especially before the year 2000. There was a relatively abrupt observed decrease in total annual catch from 1998 to 2005, where recorded landings went from over half a million kg to approximately 180,000 kg in less than a decade. Catch has slightly rebounded since that minimum, with landings reaching over 400,000 kg in six of the last seven years (Figure 53).


Multiple linear regressions and correlation analyses were performed on time series of recreational coral reef fishery catch and annual mean SST from Guam (Table 82). Evaluations measuring the association between SST and total catch for the entirety of the recreational coral reef fishery in Guam showed a negatively significant relationship between 1985 and 2016 ( $R^2 = 0.20$ ,  $p = 0.02$ ; Table 82; Figure 54). The relationship between the total annual catch and average annual SST were associated such that for every degree Celsius of temperature increase, catch would decrease by approximately 120,000 kg (Figure 54).



**Figure 53. Time series of total annual catch (kg; blue) in the Guam shore-and boat-based creel survey records plotted with average annual SST (°C; black) from 1985-2016**


**Table 82. Correlation coefficients ( $r$ ) between recreational coral reef fishery catch (in kg) and SST ( $^{\circ}\text{C}$ ) in Guam for 12 top taxa harvested from 1985-2016**

| Taxa Code | Total Catch | LUTJ  | LETH  | CARA | ACAN  | SERR  | SIGA  | SCAR  | MULL  | MUGI  | LABR  | HOLO  | BALI  |
|-----------|-------------|-------|-------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| $n = 28$  |             |       |       |      |       |       |       |       |       |       |       |       |       |
| $p$       | 0.02        | 0.01  | 0.00  | 0.01 | 0.39  | 0.01  | 0.00  | 0.00  | 0.01  | 0.01  | 0.00  | 0.00  | 0.00  |
| $r$       | -0.45       | -0.80 | -0.48 | 0.17 | -0.50 | -0.54 | -0.71 | -0.51 | -0.56 | -0.66 | -0.60 | -0.63 | -0.43 |
| $R^2$     | 0.20        | 0.64  | 0.23  | 0.03 | 0.25  | 0.30  | 0.50  | 0.26  | 0.31  | 0.43  | 0.35  | 0.39  | 0.18  |

**Figure 54. Linear regression between total annual catch (kg) for shore- and boat-based creel survey records and average annual SST ( $^{\circ}\text{C}$ ) in Guam from 1985-2016**

### 3.3.2.2 Evaluating relationship for dominant taxa

Comparisons were made for the time series of catch for prevalent taxa in Guam's recreational reef fishery as well, and it was found that all except for the Acanthuridae family showed negative statistically significant correlations with SST (Table 82). The strongest relationship observed was of that between SST and annual Lutjanidae catch, where the regression suggested that for every degree Celsius of temperature increase, catch would decrease by approximately 7,500 kg ( $R^2 = 0.64$ ,  $p = 0.00$ ; Table 82; Figure 55a). The next two strongest associations observed were for families Siganidae ( $R^2 = 0.50$ ,  $p = 0.00$ ; Figure 55b) and Mugilidae ( $R^2 = 0.43$ ,  $p = 0.01$ ; Figure 55c). The regressions performed with temperature for taxa, suggesting negative relationships with temperature, also showed that for every degree of temperature increase in degrees Celsius, Siganidae and Mugilidae recreational catch in Guam would decrease by approximately 10,000 kg and 7,500 kg, respectively.



**Figure 55. Linear regressions showing three top correlations between total annual catch (kg) for shore-and boat-based creel survey records and average annual SST (°C) in Guam for (a) Lutjanids, (b) Siganids, and (c) Mugilids from 1985–2016**

In summary, Guam and the CNMI had fishery ecosystem relationships that could be identified for the entirety of the recreational coral reef fishery. The relationship between the total annual catch and average annual SST in Guam were associated such that for every degree Celsius of temperature increase, catch would decrease by approximately 120,000 kg. The relationship between the total annual catch and average annual SST in the CNMI were associated such that for every degree Celsius of temperature increase, catch would decrease by approximately 105,000 kg.

In Guam, the linear regressions performed showed that all evaluated taxa except for the Acanthurids had a statistically significant negative relationship with average annual temperature. The three strongest associations with SST were with the Lutjanids, Siganids, and Mugilids, such that the total annual catch for each would decrease by approximately 7,500-10,000 kg for every increase in SST by one degree Celsius. In the CNMI, conversely, there were no individual family groups whose catch data had statistically significant associations with temperature, though the strongest associations observed were the Mullids (relatively close to the threshold of significance,  $p = 0.06$ ), Carangids, and Lutjanids. The relationships for families Mullidae and Carangidae were negative, though the Lutjanidae family displayed a positive relationship with SST.

### 3.4 PRIMARY PRODUCTIVITY

#### 3.4.1 CNMI


Concentrations of the pigment chlorophyll-*a* are commonly used as an index of phytoplankton biomass that represents primary production, a commonly utilized tool in identifying eutrophication also noted to be among the highest priority fishery ecosystem relationships in the WPR by participants of the Workshop (Islam and Tanaka, 2004). In Pacific regions where interannual precipitation and associated coastal runoff are relatively high, the physiochemistry of nearshore reefs is especially impacted from accompanying nutrient input resulting in increased primary production (Ansell et al., 1996).

Long-term changes in regional primary productivity have the potential to change reef fish population abundance due to the susceptibility of these assemblages in shallow areas of coastal reefs to variations in water chemistry, especially when combined with the variability of other environmental parameters like sea surface temperature (Kitiona et al. 2016). For example, it has been suggested that warming ocean temperatures coupled with decreasing environmental productivity led to waning reef fish assemblages in the Southern California Bight, likely due to a reduction in upwelling that isolated nutrients at depth (Roemmich and McGowan, 1995). With recent progress in satellite and fluorometric measurements of oceanic surface waters, time series of global and regional primary production estimated using concentrations of chlorophyll-*a* have become increasingly available, and can be used for evaluating the impact of environmental productivity on reef fish population abundance and the marine food web in general (Behrenfed et al., 2006; Messié and Radenac, 2006). Data for the study at hand were gathered from the ESA Ocean Colour Climate Change Initiative dataset version 3.1.


Considering the Ocean Colour Climate Change Initiative dataset (v3.1) for CNMI, the time series of fluorometric chlorophyll-*a* concentrations ( $\text{mg}/\text{m}^3$ ) for the years 1998-2016 in the

region is shown in Figure 56. The chlorophyll concentrations had less variability than Guam (CV = 6.28), but was relatively higher in overall average concentration. Unlike Guam, however, pigment levels appeared to have been decreasing over the course of the time series despite the non-significant nature of the associated regression. Over the 15 years of evaluated data, the average chlorophyll-*a* concentration was 0.049 mg/m<sup>3</sup>, though the lowest recorded level was seen in 2014 at 0.042 mg/m<sup>3</sup> Figure 56.

A time series of fluorometric chlorophyll-*a* concentrations (mg/m<sup>3</sup>) for the years 1998-2016 in Guam is shown in Figure 57. Pigment concentration in the upper 200 meters had moderate variability over the course of the time series (CV=7.03). Also, there seemed to be a slight increase in pigment concentrations over the course of collected data despite the lack of a significant trend over the same time. The average chlorophyll-*a* concentration over this time was 0.048 mg/m<sup>3</sup>, with the highest recorded levels being observed in 2005 at 0.055 mg/m<sup>3</sup> and the lowest occurring earlier in 2002 (0.042 mg/m<sup>3</sup>; Figure 57).



**Figure 56. Time series of fluorometric chlorophyll-*a* concentrations (mg/m<sup>3</sup>) around the CNMI from 1998-2016 (CV=6.28)**



**Figure 57. Time series of fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>) around Guam from 1998-2016 (CV=7.03)**

### 3.4.1.1 Evaluating relationship for entire reef fishery

A plot showing the relationship between these same chlorophyll levels and catch time series from the recreational coral reef fishery in the CNMI from 2000-2016 is depicted in Figure 58. Catch, again, was even more variable than the environmental data evaluated (CV=19.4), and was at about the same levels as Guam. Total annual catch in the fishery has been decreasing over the last decade and a half despite a spike in catch during 2013 that gave the maximum observed annual catch over this time series (~338,000 kg). The levels of current catch (i.e., for 2014-2016) are the lowest for the entirety of the recreational fishery over the past decade and a half (~165,000 kg; Figure 58).

In pattern with the analyses completed for Guam, linear regressions and correlation analyses were conducted for the time series of the CNMI recreational coral reef fishery catch (with phase lag) with fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>) gathered for the 15 years between 2000-2014. The chlorophyll-a concentrations and total annual catch for the all harvested taxa had a positive relationship between 2000 and 2014, though the relationship was far from being considered statistically significant ( $r = 0.32$ ,  $p = 0.25$ ; Table 83; Figure 59). Though not significant, the regression was extrapolated to determine that, following this pattern, every increase of 0.01 mg/m<sup>3</sup> in chlorophyll-a concentration would cause increase by nearly 62,000 kg two years later for all the CNMI recreational reef fishery ( $R^2=0.11$ ,  $p = 0.25$ ; Figure 59).





Figure 58. Comparison of the CNMI recreational reef fish catch (kg; black) from creel survey records with two years of time lag (t+2 years) and fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>; blue) from 2000-2014 ( $r = 0.32$ )

Table 83. Correlation coefficients ( $r$ ) from comparisons of time series of the CNMI recreational coral reef fishery annual catch (kg) and fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>) from 2000-2014

| Taxa Code             | Total Catch | LUTJ  | LETH  | CARA | ACAN | SERR | SIGA  | SCAR  | MULL | MUGI  | LABR  | HOLO | BALI |
|-----------------------|-------------|-------|-------|------|------|------|-------|-------|------|-------|-------|------|------|
| <b>n = 15</b>         |             |       |       |      |      |      |       |       |      |       |       |      |      |
| <i>p</i>              | 0.25        | 0.47  | 0.14  | 0.67 | 0.37 | 0.09 | 0.72  | 0.80  | 0.99 | 0.83  | 0.83  | 0.10 | 0.72 |
| <i>r</i>              | 0.32        | -0.20 | -0.04 | 0.12 | 0.25 | 0.45 | -0.10 | -0.07 | 0.00 | -0.06 | -0.06 | 0.44 | 0.10 |
| <i>R</i> <sup>2</sup> | 0.11        | 0.04  | 0.00  | 0.02 | 0.06 | 0.20 | 0.01  | 0.01  | 0.00 | 0.00  | 0.00  | 0.20 | 0.01 |






**Figure 59. Linear regression between total annual catch (kg) phase lag (t+2 years) and fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>) in CNMI from 2000-2014**

### 3.4.1.2 Evaluating relationship for dominant taxa

Out of the many linear regressions completed for catch time series of dominant taxa in the CNMI's recreational coral reef fishery, none of them were determined to be significantly related to the recorded chlorophyll-*a* concentrations from the same area (Table 83). Of the 12 analyzed groups, the three with the strongest (non-significant) relationship with local chlorophyll concentrations were the Serranids, the Acanthurids, and the Holocentrids ( $R^2 = 0.20, 0.20, 0.06$ , respectively; Figure 60a-c). It is interesting to note that, unlike Guam, the overall relationship between pigment concentration and catch for the entirety of the reef fishery in the region was positive, though non-significant ( $r = 0.32, p = 0.25$ ), and the strongest determined associations among the analyzed taxa were all positive as well (Table 83).




**Figure 60. Linear regressions showing the three top correlations between total annual catch (kg) for the CNMI from creel survey records with phase lag (t+2 years) and fluorometric chlorophyll-*a* concentrations (mg/m<sup>3</sup>) for (a) Holocentrids, (b) Serranids, and (c) Acanthurids from 2000–2014**

### 3.4.2 Guam

#### 3.4.2.1 Evaluating relationship for entire reef fishery

A plot depicting the comparison of the fluorometric chlorophyll-*a* concentrations and recreational coral reef fishery catch time series from 1998 - 2014 in Guam is shown in Figure 61. Catch levels were relatively variable over the course of the time series when considering the variation in pigment levels (CV=26.2; Figure 61). A gradual drop in total annual catch was observed starting from 1998 before stabilizing in the late 2000s, where recorded catch decreased to approximately a quarter million, and rose back up to over half a million kilograms in more recent years; it is of note that the minimum catch and maximum chlorophyll concentration depicted in this plot both occurred in the year 2005 (Figure 61).


Linear regressions and correlation analyses were conducted for the time series of the Guam recreational coral reef fishery catch (with phase lag) with fluorometric chlorophyll-*a* concentrations ( $\text{mg}/\text{m}^3$ ) gathered from the Ocean Colour Climate Change Initiative dataset (v3.1) for the 17 years between 1998 and 2014. It was found that the chlorophyll concentrations and total annual catch for the all harvested taxa had a negative relationship between 1989 and 2015, though it was slightly over the threshold of significance ( $r = -0.45$ ,  $p = 0.02$ ; Table 84; Figure 62). The association was statistically significant, and it was determined that for every increase of  $0.01 \text{ mg}/\text{m}^3$  in chlorophyll-*a* concentration, catch would approximately decrease by 180,000 kg after two years all of the Guam recreational fishery ( $R^2 = 0.20$ ,  $p = 0.02$ ; Table 84; Figure 62).



**Figure 61. Comparison of Guam recreational reef fish catch for shore-and boat-based creel survey records (kg; black) with two years of time lag (t+2 years) and fluorometric chlorophyll-*a* concentrations ( $\text{mg}/\text{m}^3$ ; blue) from 1998-2014**

**Table 84. Correlation coefficients ( $r$ ) from comparisons of time series of for shore-and boat-based creel survey records in Guam (kg) and fluorometric chlorophyll- $a$  concentrations ( $\text{mg}/\text{m}^3$ ) for 12 top taxa harvested from 1998 - 2014. Significant correlations are indicated in bold ( $\alpha=0.05$ )**

| Taxa Code               | Total Catch | LUTJ  | LETH  | CARA  | ACAN  | SERR  | SIGA  | SCAR  | MULL  | MUGI         | LABR  | HOLO  | BALI         |
|-------------------------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|--------------|-------|-------|--------------|
| <b>n = 17</b>           |             |       |       |       |       |       |       |       |       |              |       |       |              |
| <b><math>p</math></b>   | 0.07        | 0.62  | 0.16  | 0.73  | 0.44  | 0.51  | 0.17  | 0.42  | 0.08  | <b>0.04</b>  | 0.47  | 0.21  | <b>0.03</b>  |
| <b><math>r</math></b>   | -0.45       | -0.13 | -0.36 | -0.09 | -0.20 | -0.17 | -0.35 | -0.21 | -0.43 | <b>-0.50</b> | -0.19 | -0.32 | <b>-0.53</b> |
| <b><math>R^2</math></b> | 0.20        | 0.02  | 0.13  | 0.01  | 0.04  | 0.03  | 0.12  | 0.04  | 0.19  | <b>0.25</b>  | 0.03  | 0.11  | <b>0.28</b>  |



**Figure 62. Linear regression between total annual catch (kg) for Guam shore-and boat-based creel survey records with phase lag (t+2 years) and fluorometric chlorophyll- $a$  concentrations ( $\text{mg}/\text{m}^3$ ) from 1998-2014**

### 3.4.2.2 Evaluating relationship for dominant taxa


The several linear regression and correlation analyses performed for time series of catch on the taxa level of Guam's recreational reef fishery showed that for dominant taxa in the fishery, and only two of the 12 analyzed groups had statistically significant relationships with local chlorophyll concentrations: the Balistids and the Mugilids (Table 84). The relationship between catch of species in the Balistidae group and chlorophyll concentration was shown to have negatively significant relationship such that for every increase of  $0.01 \text{ mg}/\text{m}^3$  in chlorophyll- $a$  concentration, catch would drop by more than 1,700 kg two years later when harvesting members of the Balistidae family ( $R^2=0.28$ ,  $p = 0.03$ ; Table 84; Figure 63a). The relationship between catch of members of the Mugilidae group and chlorophyll concentration was also shown to be negatively significant, but to a lesser degree. With a rise of  $0.01 \text{ mg}/\text{m}^3$  in chlorophyll- $a$

levels, recreational catch of the Mugilids would decrease by approximately over 4,600 kg after two years for the group ( $R^2=0.25$ ,  $p = 0.04$ ; Table 84; Figure 63b;). The next strongest relationship as determined by the regressions was not significant, but was similarly negative (Mullidae;  $R^2=0.19$ ,  $p=0.08$ ; Table 84; Figure 63c); all four of these potential fishery ecosystem relationships, however, were positive.

In the CNMI, there were no statistically significant relationships discovered between chlorophyll concentrations and any of the 12 prevalent taxa evaluated in this study, nor to the total fishery annual catch in its entirety. The lack of identifiable associations could have been attributed to the relatively short time series of data available for comparison at 15 years. While there were several families observed that had relationships on the cusp of being deemed significant according to resulting coefficients of determination, such as Serranidae and Holocentridae, they were positively associated.

In summary for Guam, it was determined that there existed a negatively significant relationship between reef recreational catch and fluorometric chlorophyll-*a* concentrations ( $\text{mg}/\text{m}^3$ ) from the Ocean Colour Climate Change Initiative dataset (v3.1) for the entirety of the fishery. For every increase of  $0.01 \text{ mg}/\text{m}^3$  in chlorophyll-*a* concentration, catch would approximately decrease by 180,000 kg across all harvested taxa two years later. Potential statistically significant fishery ecosystem relationships were also observed for the Balistidae and Mugilidae groups, where the catch of each group would decrease by approximately 1,700 and 4,600 kg, respectively, given two years of phase lag with a similar increase in fluorometric chlorophyll.

Uncertainty levels were relatively high in evaluations including chlorophyll-*a* concentrations due to the nature of incorporating phase lag and not smoothing the catch data. The largest issue in performing comparison analyses between catch from reef fisheries in the Mariana Archipelago and fluorometric chlorophyll-*a* concentrations was the relatively short time series (i.e. small sample size). Robust, homogenous time series highlighting interdecadal patterns in these regions were difficult to obtain due to time series merging several sources of chlorophyll concentration to elongate the range of continuous data. For example, the ESA's OCC CCI dataset only permitted the use of less than two decades of data when evaluating the territories with the incorporation of phase lag. The length of the applied lag has a large impact in the patterns observed, so the relatively short extent of the available time series may obfuscate some of the identified relationships.



**Figure 63. Linear regressions showing the three top correlations between total annual catch (kg) for Guam for shore- and boat-based creel survey records with phase lag (t+2 years) and fluorometric chlorophyll-a concentrations (mg/m<sup>3</sup>) for (a) Balistidae, (b) Mugilidae, and (c) Mullidae from 1998–2014.**

### 3.5 MULTIVARIATE ASSESSMENTS OF OTHER ECOSYSTEM VARIABLES

#### 3.5.1 Non-metric Multidimensional Scaling

There were several other prioritized fishery ecosystem relationships for coral reefs in the Mariana Archipelago involving environmental parameters that were not to be addressed in this initial evaluation including: the Oceanic Niño Index (ONI), the Pacific Decadal Oscillation (PDO), sea level height, pH, dissolved oxygen, and salinity. Further descriptions of these climate and oceanic indicators are available in Section **Error! Reference source not found.** Sea surface height data were aggregated from the Ocean Service, Tides, and Currents, and Sea Level database operated (NOAA/NOS/CO-OPS). Basin-wide data ONI were taken from NOAA's Nation Centers for Environmental Information- Equatorial Pacific Sea Surface Temperature Database (Climate Prediction Center Internet Team 2015). Similarly, PDO data were obtained from NOAA's Earth System Research Laboratory Physical Sciences Division originally derived from OI.v1 and OI.v2 SST parameters (NOAA PDO). Salinity data for American Samoa were gathered from Simple Ocean Data Assimilation (SODA) version 3.3.1 (Carton and Giese 2008). Rainfall estimates were obtained through the National Weather Service in American Samoa (NWS-G).


Non-metric multidimensional scaling (NMS), a form of multivariate analysis that orders sample units along synthetic axes to reveal patterns of composition and relative abundance (Peck, 2016), is most commonly utilized when looking to identify patterns in heterogeneous species response data (Peck, 2016). For this study, NMS was used to help identify associations between coral reef fishery parameters and environmental factors using the program PCORD 7. To ensure the same length of time series for all catch and environmental variables considered, data was analyzed from 1989-2015 to allow for the inclusion of more parameters (e.g. pH) for which longer-term time series were unavailable. The generated axes represent the best fit of patterns of redundancy in the catch data used as input, and the resulting ordination scores are a rank-order depiction of associations in the original dataset.

NMS produces robust results even in the presence of outliers by avoiding parametric and distributional assumptions (Peck, 2016). The only assumption to be met in NMS is that the relationship between the original rank ordered distances between sample units and the reduced distances in the final solution should be monotonic; that is, the slope of the association between the two is flat or positive, as determined by the stress statistic. In the most general terms, interpretable and reliable ordination axes have stress less than 10 up to 25 for datasets with large sample size, but large stress scores (i.e. greater than 30) may suggest that the final ordination results have little association with the original data matrix. Additionally, NMS ordination scores vary depending on the number of dimensions/axes designated to be solved (Peck, 2016). Dimensionality (i.e. number of axes for the final solution) for each test was identified through PCORD result recommendations based on final stress being lower than that for 95% of randomized runs (i.e.  $p \leq 0.05$ ). Tau is a statistic that represents the rank correlations of the ordination scores to the original data matrices, and was used to identify explanatory variables with associations to the ordination axes. For the MHI test, data from 13 species/taxa groups from 1989 - 2015 (27 years) were included along with 10 variables of environmental data collected during the same time period.

### 3.5.1.1 CNMI


The resulting ordination scores from the NMS analysis performed on boat-based expanded creel survey catch records and the previously mentioned environmental parameters recommended a one dimensional solution, which accounting for 87.2% of the cumulated variance observed in the CNMI boat-based creel survey data. The NMS final stress was moderate for the real runs (13.9), but low relative to stress from the randomization runs (31.0; Figure 64). The final ordination scores for the families considered were scaled on a gradient relative to the individual ordination axis, the overlying environmental joint biplot is situated to the left of the final ordination points (Figure 64).

The only environmental parameter included in this analysis that displayed a significant relationship with the lone axis was PDO, though that association was negative. ( $\tau = -0.47$ ), Although this NMS run was not able to identify any other environmental parameters significantly correlated to the ordination axis, additionally relatively strong associations exist between sea level height ( $\tau = 0.33$ ) and pH ( $-0.31$ ; Figure 65). Replicate NMS runs had similar stress levels for the final generated result.



**Figure 64. NMS stress plot showing the stress test to determine dimensionality for the final solution for the CNMI multivariate analysis; a one-axis solution was recommended**






**Figure 65. One-dimensional scatterplot overlaid with a joint biplot depicting ordination scores resulting from an NMS analysis on creel survey expanded catch data and prominent environmental parameters in the CNMI from 2000-2014**


### 3.5.1.2 Guam

The Guam NMS identified two orthogonal axes for the final solution that accounted for 93.6% of the cumulative observed variance in shore- and boat-based creel survey data from Guam. The final stress for the Guam NMS barely less than 10, though it was notable lower than the average final stress from randomizations (14.2; Figure 66). A majority of the families were clustered in ordination space, with the notable exception of Carangidae (Figure 67).

The final ordination scores for the Guam NMS did not show any environmental parameters with a statistically significant correlation to the first axis ( $r^2 = 0.62$ ; Figure 67). SST ( $\tau = -0.50$ ) and SSTA ( $\tau = -0.50$ ) were both negatively associated with the Axis 2 ( $r^2 = 0.32$ ), and pH had a significantly positive relationship with the axis ( $\tau = 0.56$ ). Additionally, Axis 2 was shown to also be negatively associated with pH ( $\tau = -0.37$ ; Figure 67). Replicate NMS runs had similar stress levels for the final generated result.



**Figure 66. NMS scree plot showing the stress test to determine dimensionality for the final solution for the Guam multivariate analysis; two-axis solution was recommended**



**Figure 67. Two-dimensional scatterplot overlaid with a joint biplot depicting ordination scores resulting from an NMS analysis on creel survey expanded catch data and prominent environmental parameters in Guam from 1989-2014**

Ultimately, stress values for all analyses were relatively low, suggesting that the generated ordination scores were robust and useful for interpretation relative to the ordination axes. Nearly all included environmental parameters had a statistically significant relationship with at least one ordination axis in at least one of the final solutions, suggesting that these parameters likely intertwine in complicated processes to produce observed impacts on coral reef fisheries in the U.S. Western Pacific. Though a fishery ecosystem relationship may have not been explicitly identified in NMS runs of this preliminary evaluation, it does not preclude the possibility that an association may still exist.

### 3.6 REFERENCES

Ansell, A.D., Gibson, R.N., Barnes, M., and Press, U.C.L., 1996. Coastal fisheries in the Pacific Islands. *Oceanography and Marine Biology: an annual review*, 34(395), 531 p.

- Behrenfeld, M.J., T O'Malley, R., Siegel, D.A., McClain, C.R., Sarmiento, J.L., Feldman, G.C., Milligan, A.J., Falkowski, P.G., Letelier, R.M. and Boss, E.S., 2006. Climate-driven trends in contemporary ocean productivity. *Nature*, 444(7120), 752 p.
- Carton, J.A. and Giese B.S., 2008. A Reanalysis of Ocean Climate Using Simple Ocean Data Assimilation (SODA), *Mon. Weather Rev.*, 136, pp. 2999-3017.
- Climate Prediction Center Internet Team, 2015. Cold and warm episodes by season. NOAA Center for Weather and Climate Prediction, Maryland, United States. Accessed March 2017. Available from [http://www.cpc.ncep.noaa.gov/products/analysis\\_monitoring/ensostuff/ensoyears.shtml](http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml).
- Hawhee, J.M., 2007. Western Pacific Coral Reef Ecosystem Report. Report prepared for the Western Pacific Regional Fishery Management Council, Honolulu, HI 96813 USA, 185.
- Islam, M.S., and Tanaka, M., 2004. Impacts of pollution on coastal and marine ecosystems including coastal and marine fisheries and approach for management: a review and synthesis. *Marine pollution bulletin*, 48(7), pp. 624-649.
- Kitiona, F., Spalding, S., and Sabater, M., 2016. The impacts of climate change on coastal fisheries in American Samoa. University of Hawaii at Hilo, HI 96720 USA, 18 p.
- Longhurst, A.R. and Pauly, D., 1987. The Ecology of Tropical Oceans. *Academic Press Inc., London*, 407 p.
- McClanahan, T.R., Graham, N.A., MacNeil, M.A., Muthiga, N.A., Cinner, J.E., Bruggemann, J.H., and Wilson, S.K., 2011. Critical thresholds and tangible targets for ecosystem-based management of coral reef fisheries. *Proceedings of the National Academy of Sciences*, 108(41), pp.17230-17233.
- Messié, M. and Radenac, M.H., 2006. Seasonal variability of the surface chlorophyll in the western tropical Pacific from SeaWiFS data. *Deep Sea Research Part I: Oceanographic Research Papers*, 53(10), pp. 1581-1600.
- NOAA/NESDIS/OceanWatch – Central Pacific, n.d. OceanWatch – Sea-surface Temperature, AVHRR Pathfinder v5.0 + GAC – Monthly. NOAA, Maryland, USA. Accessed March 2017. Available from <<http://oceanwatch.pifsc.noaa.gov/thredds/dodsC/pfgac/monthly>>.
- NOAA/NOS/CO-OPS, n.d. Observed Water Levels. NOAA, Maryland, United States. Accessed March 2017. Available from <https://tidesandcurrents.noaa.gov/waterlevels.html>.
- NOAA PDO, n.d. Pacific Decadal Oscillation Index. NOAA, Maryland, United States. Accessed March 2017. Available from <http://research.jisao.washington.edu/pdo/PDO.latest>.
- NWS-G, n.d. National Weather Service Forecast Office, Tiyan, Guam. NOAA National Weather Service. Accessed March 2017.

- Ocean Colour Climate Change Initiative dataset, Version 3.1, European Space Agency. Accessed August 2017. Available from <<http://www.esa-oceancolour-cci.org/>>.
- Ochavillo, D., 2012. Coral Reef Fishery Assessment in American Samoa. Department of Marine and Wildlife Resources, Pago Pago, American Samoa 96799 USA, 29 p.
- Peck, J.E., 2016. Multivariate Analysis for Ecologists: Step-by- Step, Second edition. MjM Software Design, Gleneden Beach, OR. 192 p.
- Remington, T.R. and Field, D.B., 2016. Evaluating biological reference points and data-limited methods in Western Pacific coral reef fisheries. Report prepared for the Western Pacific Regional Fishery Management Council, Honolulu, HI 96813 USA, 134 p.
- Roemmich, D. and McGowan, J., 1995. Climatic warming and the decline of zooplankton in the California Current. *Science: New York then Washington*, pp. 1324-1324.
- Richards B.L., Williams I.D., Vetter O.J., and Williams G.J., 2012. Environmental factors affecting large-bodied coral reef fish assemblages in the Mariana Archipelago. *PLoS ONE* 7(2), e31374.
- Weng, K.C. and Sibert, J.R., 2000. Analysis of the Fisheries for two pelagic carangids in Hawaii. Joint Institute for Marine and Atmospheric Research, University of Hawaii at Manoa, Honolulu, HI 96822 USA, 78 p.
- Williams, I.D., Baum, J.K., Heenan, A., Hanson, K.M., Nadon, M.O., and Brainard, R.E., 2015. Human, Oceanographic and Habitat Drivers of Central and Western Pacific Coral Reef Fish Assemblages. *PLoS ONE* 10(5): e0129407.

**APPENDIX A: LIST OF MANAGEMENT UNIT SPECIES****CNMI****1. Bottomfish Multi-species Stock Complex (FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b>                     | <b>Scientific Name</b>             |
|---|---|------------------------------------|
| 214                                       | red snapper, silvermouth (lehi)         | <i>Aphareus rutilans</i>           |
| 213                                       | grey snapper, jobfish                   | <i>Aprion virescens</i>            |
| 112                                       | giant trevally, jack                    | <i>Caranx ignobilis</i>            |
| 111                                       | black trevally, jack                    | <i>Caranx lugubris</i>             |
| 231                                       | blacktip grouper                        | <i>Epinephelus fasciatus</i>       |
| 241                                       | lunartail grouper<br>(lyretail grouper) | <i>Variola lauti</i>               |
| 203                                       | red snapper (ehu)                       | <i>Etelis carbunculus</i>          |
| 210                                       | red snapper (onaga)                     | <i>Etelis coruscans</i>            |
| none                                      | ambon emperor                           | <i>Lethrinus amboinensis</i>       |
| 350                                       | redgill emperor                         | <i>Lethrinus rubrioperculatus</i>  |
| 253                                       | blueline snapper                        | <i>Lutjanus kasmira</i>            |
| none                                      | yellowtail snapper                      | <i>Pristipomoides auricilla</i>    |
| 212                                       | pink snapper (paka)                     | <i>Pristipomoides filamentosus</i> |
| 209                                       | yelloweye snapper                       | <i>Pristipomoides flavipinnis</i>  |
| 207                                       | pink snapper (kalekale)                 | <i>Pristipomoides seiboldi</i>     |
| 204                                       | flower snapper (gindai)                 | <i>Pristipomoides zonatus</i>      |
| 220                                       | amberjack                               | <i>Seriola dumerili</i>            |

**2. Crustacean deep-water shrimp complex (non-FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>   |
|---|---------------------|--------------------------|
| 508                                       | deepwater shrimp    | <i>Heterocarpus</i> spp. |

**3. Crustacean spiny lobster complex (non-FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>        |
|---|---------------------|-------------------------------|
| 504                                       | spiny lobster       | <i>Panulirus marginatus</i>   |
| 504                                       | spiny lobster       | <i>Panulirus penicillatus</i> |

**4. Crustacean slipper lobster complex (non-FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b> |
|---|---------------------|------------------------|
| 505                                       | Slipper lobster     | Scyllaridae            |

**5. Crustacean Kona crab complex (non-FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b> |
|---|---------------------|------------------------|
| 502                                       | Kona crab           | <i>Ranina ranina</i>   |

**6. Precious coral black coral complex (non-FSSI)**

| <b>DFW<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>      |
|---|---------------------|-----------------------------|
| none                                      | Black Coral         | <i>Anitpathes dichotoma</i> |
| none                                      | Black Coral         | <i>Antipathes grandis</i>   |
| none                                      | Black Coral         | <i>Antipathes ulex</i>      |

**7. Exploratory area precious coral (except black coral; non-FSSI)**

| <b>DFW Creel Species Code</b> | <b>Species Name</b> | <b>Scientific Name</b>      |
|-------------------------------|---------------------|-----------------------------|
| none                          | Pink coral          | <i>Corallium secundum</i>   |
| none                          | Pink coral          | <i>Corallium regale</i>     |
| none                          | Pink coral          | <i>Corallium laauense</i>   |
| none                          | Bamboo coral        | <i>Lepidisis olapa</i>      |
| none                          | Bamboo coral        | <i>Acanella</i> spp.        |
| none                          | Gold Coral          | <i>Gerardia</i> spp.        |
| none                          | Gold Coral          | <i>Callogorgia gilberti</i> |
| none                          | Gold Coral          | <i>Narella</i> spp.         |
| none                          | Gold Coral          | <i>Calyptrophora</i> spp.   |

**8. Coral reef ecosystem (non-FSSI)**

| <b>DFW Creel Species Code</b> | <b>Species Name</b>    | <b>Scientific Name</b>           | <b>Grouping</b> |
|-------------------------------|------------------------|----------------------------------|-----------------|
| 357                           | Bigeye Emperor         | <i>Monotaxis grandoculus</i>     | Lethrinidae     |
| 353                           | Blackspot Emperor      | <i>Lethrinus harak</i>           | Lethrinidae     |
| 310                           | Emperor (mafute/misc.) | <i>Lethrinus</i> sp.             | Lethrinidae     |
| 356                           | Flametail Emperor      | <i>Lethrinus fulvus</i>          | Lethrinidae     |
| 351                           | Longnose Emperor       | <i>Lethrinus olivaceus</i>       | Lethrinidae     |
| 352                           | Orangefin Emperor      | <i>Lethrinus erythracanthus</i>  | Lethrinidae     |
| 361                           | Ornate Emperor         | <i>Lethrinus ornatus</i>         | Lethrinidae     |
| 358                           | Stout Emperor          | <i>Gymnocranius</i> sp.          | Lethrinidae     |
| 355                           | Yellowlips Emperor     | <i>Lethrinus xanthochilis</i>    | Lethrinidae     |
| 359                           | Yellowspot emperor     | <i>Gnathodentex aurolineatus</i> | Lethrinidae     |
| 354                           | Yellowstripe Emperor   | <i>Lethrinus obsoletus</i>       | Lethrinidae     |
| 362                           | Yellowtail Emperor     | <i>Lethrinus atkinsoni</i>       | Lethrinidae     |
| 115                           | Bigeye Trevally        | <i>Caranx sexfasciatus</i>       | Carangidae      |
| 113                           | Bluefin Trevally       | <i>Caranx melampygus</i>         | Carangidae      |
| 114                           | Brassy Trevally        | <i>Caranx papuesis</i>           | Carangidae      |
| 105                           | EE: Juvenile Jacks     | <i>Canranx</i> sp.               | Carangidae      |
| 104                           | Jacks (misc.)          | <i>Caranx</i> sp.                | Carangidae      |
| 101                           | Leatherback            | <i>Scomberoides lysan</i>        | Carangidae      |

|     |                          |                                     |              |
|-----|--------------------------|-------------------------------------|--------------|
| 103 | Mackerel Scad            | <i>Decapterus macarellus</i>        | Carangidae   |
| 410 | Rainbow Runner           | <i>Elagatis bipinnulatus</i>        | Carangidae   |
| 117 | Small-spotted pompano    | <i>Trachinotus bailloni</i>         | Carangidae   |
| 116 | Snubnose pompano         | <i>Trachinotus blochii</i>          | Carangidae   |
| 110 | Yellow Spotted Trevally  | <i>Carangoides orthogrammus</i>     | Carangidae   |
| 380 | Bluebanded Surgeonfish   | <i>Acanthurus lineatus</i>          | Acanthuridae |
| 383 | Bluelined Surgeon        | <i>Acanthurus nigroris</i>          | Acanthuridae |
| 384 | Bluespine Unicornfish    | <i>Naso unicornis</i>               | Acanthuridae |
| 381 | Convict Tang             | <i>Acanthurus triostegus</i>        | Acanthuridae |
| 319 | Orangespine Unicornfish  | <i>Naso lituratus</i>               | Acanthuridae |
| 318 | Surgeonfish (misc.)      | <i>Acanthurus</i> sp.               | Acanthuridae |
| 320 | Unicornfish (misc.)      | <i>Naso</i> sp.                     | Acanthuridae |
| 382 | Yellowfin Surgeonfish    | <i>Acanthurus xanthopterus</i>      | Acanthuridae |
| 102 | Bigeye Scad              | <i>Selar crumenophthalmus</i>       | Atulai       |
| 239 | Coral Grouper            | <i>Epinephelus corallicola</i>      | Serranidae   |
| 237 | Flagtail Grouper         | <i>Cephalopholis urodeta</i>        | Serranidae   |
| 206 | Grouper (misc.)          | Serranidae                          | Serranidae   |
| 233 | Highfin Grouper          | <i>Epinephelus maculatus</i>        | Serranidae   |
| 234 | Honeycomb Grouper        | <i>Epinephelus merra</i>            | Serranidae   |
| 235 | Marbled Grouper          | <i>Epinephelus polyphkadion</i>     | Serranidae   |
| 236 | Peacock Grouper          | <i>Cephalopholis argus</i>          | Serranidae   |
| 244 | Pink Grouper             | <i>Saloptia powelli</i>             | Serranidae   |
| 238 | Saddleback Grouper       | <i>Plectropomus laevis</i>          | Serranidae   |
| 242 | Tomato Grouper           | <i>Cephanopholis sonnerati</i>      | Serranidae   |
| 240 | White Lyretail Grouper   | <i>Variola albimarginata</i>        | Serranidae   |
| 243 | Yellow Banded Grouper    | <i>Cephalopholis igarashiensis</i>  | Serranidae   |
| 316 | Snapper (misc. shallow)  | Lutjanidae                          | Lutjanidae   |
| 250 | Humpback Snapper         | <i>Lutjanus gibbus</i>              | Lutjanidae   |
| 251 | Onespot Snapper          | <i>Lutjanus monostigmus</i>         | Lutjanidae   |
| 254 | Red Snapper              | <i>Lutjanus bohar</i>               | Lutjanidae   |
| 208 | Smalltooth Jobfish       | <i>Aphareus furca</i>               | Lutjanidae   |
| 371 | Dash & Dot Goatfish      | <i>Parupeneus barberrinus</i>       | Mullidae     |
| 321 | Goatfish (juvenile-misc) | Mullidae                            | Mullidae     |
| 322 | Goatfish (misc.)         | Mullidae                            | Mullidae     |
| 323 | Sidespot Goatfish        | <i>Parupeneus pleurostigma</i>      | Mullidae     |
| 372 | Two-barred Goatfish      | <i>Parupeneus bifasciatus</i>       | Mullidae     |
| 370 | Yellowstripe Goatfish    | <i>Mulloidichthys flavolineatus</i> | Mullidae     |
| 314 | Parrotfish (misc.)       | <i>Scarus</i> sp.                   | Scaridae     |
| 315 | Seagrass Parrotfish      | <i>Leptoscarus vaigiensis</i>       | Scaridae     |



|     |                        |                                     |                   |
|-----|------------------------|-------------------------------------|-------------------|
| 506 | Octopus                | <i>Octopus i.</i>                   | Mollusk           |
| 510 | Squid                  | <i>Teuthida</i>                     | Mollusk           |
| 516 | Trochus                | <i>Trochus</i> sp.                  | Mollusk           |
| 522 | Clam/bivalve           | Bivalvia                            | Mollusk           |
| 106 | Mullet                 | Mugilidae                           | Mugilidae         |
| 304 | Rabbitfish (hitting)   | <i>Siganus</i> sp.                  | Siganidae         |
| 306 | Rabbitfish (h.feda)    | <i>Siganus punctatus</i>            | Siganidae         |
| 307 | Rabbitfish (menahac)   | <i>Siganus</i> sp.                  | Siganidae         |
| 308 | Rabbitfish (sesjun)    | <i>Siganus spinus</i>               | Siganidae         |
|     | Bolbometopon muricatum | <i>Bumphead parrotfish</i>          |                   |
| 391 | Cheilinus undulatus    | <i>Napoleon wrasse</i>              |                   |
|     | Reef sharks (misc)     | Carcharhinidae                      | Carcharhinidae    |
|     | Hammerhead shark       | Sphyrnidae                          | Carcharhinidae    |
| 338 | Angelfish              | Pomacanthidae                       | Other CRE-Finfish |
| 338 | Butterflyfish          | Chaetodontidae                      | Other CRE-Finfish |
| 324 | Bigeye/glasseye        | <i>Heteropriacanthus cruentatus</i> | Other CRE-Finfish |
| 396 | Blue Razorfish         | <i>Xyrichtys pavo</i>               | Other CRE-Finfish |
| 397 | Bronzespot Razorfish   | <i>Xyrichtys celebicus</i>          | Other CRE-Finfish |
| 260 | Cardinal Misc.         | Apogonidae                          | Other CRE-Finfish |
| 162 | Cornetfish             | <i>Fistularia commersonii</i>       | Other CRE-Finfish |
| 332 | Damsel fish            | Pomacentridae                       | Other CRE-Finfish |
| 341 | Filefish (misc)        | Monacanthidae                       | Other CRE-Finfish |
| 340 | Flounder (misc)        | <i>Bothus</i> sp.                   | Other CRE-Finfish |
| 328 | Fusilier (misc.)       | Caesionidae                         | Other CRE-Finfish |
| 325 | Goggle-eye             | <i>Priacanthus hamrur</i>           | Other CRE-Finfish |
| 195 | Lizardfish misc.       | Synodontidae                        | Other CRE-Finfish |
| 180 | Milkfish               | <i>Chanos chanos</i>                | Other CRE-Finfish |
| 329 | Mojarra                | <i>Gerres</i> sp.                   | Other CRE-Finfish |
| 140 | Moray eel              | Muraenidae                          | Other CRE-Finfish |
| 170 | Needlefish             | Belonidae                           | Other CRE-Finfish |
| 343 | Picasso Trigger        | <i>Rhinecanthus aculeatus</i>       | Other CRE-Finfish |
| 348 | Pufferfish             | Tetraodontidae                      | Other CRE-Finfish |
| 395 | Razorfish (misc)       | Tribe Novaculini                    | Other CRE-Finfish |
| 130 | Scorpionfishes         | Scorpaenidae                        | Other CRE-Finfish |
| 330 | Sweetlips              | <i>Plectorhinchus picus</i>         | Other CRE-Finfish |
| 342 | Triggerfish (misc.)    | Balistidae                          | Other CRE-Finfish |
| 163 | Trumpetfish            | <i>Aulostomus chinensis</i>         | Other CRE-Finfish |
| 344 | Wedge Trigger          | <i>Rhinecanthus rectangulus</i>     | Other CRE-Finfish |
| 312 | Squirrelfish           | Holocentridae                       | Squirrelfish      |

|     |                           |                             |                          |
|-----|---------------------------|-----------------------------|--------------------------|
| 313 | Soldierfish (misc.)       | Holocentridae               | Squirrelfish             |
| 302 | Wrasse                    | Labridae                    | Wrasse                   |
| 390 | Tripletail Wrasse         | <i>Cheilinus trilobatus</i> | Wrasse                   |
| 309 | Rudderfish (guilli)       | <i>Kyphosus</i> sp.         | Rudderfish               |
| 373 | Highfin Rudderfish Silver | <i>Kyphosus cinerascens</i> | Rudderfish               |
| 374 | Highfin Rudderfish Brown  | <i>Kyphosus</i> sp.         | Rudderfish               |
| 200 | Bottomfish (misc)         | n/a                         | Misc. Bottomfish         |
| 300 | Reef fish (misc)          | n/a                         | Misc. Reef Fish          |
|     | Shallow bottom            | n/a                         | Misc. Shallow bottomfish |
| 501 | Crabs (misc)              | n/a                         | Crustaceans              |
| 503 | Coconut Crab              | <i>Birgus latro</i>         | Crustaceans              |
| 500 | Invertebrates             | n/a                         | Other Invertebrates      |
| 514 | Sea Cucumber              | Cucumariidae                | Other Invertebrates      |
| 600 | Seaweeds                  | n/a                         | Algae                    |
| 602 | Lemu                      | n/a                         | Algae                    |

**GUAM****1. Bottomfish Multi-species Stock Complex (FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b>             | <b>Scientific Name</b>             |
|--|---------------------------------|------------------------------------|
| 32302                                      | red snapper, silvermouth (lehi) | <i>Aphareus rutilans</i>           |
| 32303                                      | grey snapper, jobfish           | <i>Aprion virescens</i>            |
| 31404                                      | giant trevally, jack            | <i>Caranx ignoblis</i>             |
| 31405                                      | black trevally, jack            | <i>Caranx lugubris</i>             |
| 28919                                      | blacktip grouper                | <i>Epinephelus fasciatus</i>       |
| 28941                                      | lunartail (lyretail) grouper    | <i>Variola lauti</i>               |
| 32304                                      | red snapper (ehu)               | <i>Etelis carbunculus</i>          |
| 32305                                      | red snapper (onaga)             | <i>Etelis coruscans</i>            |
| 32818                                      | ambon emperor                   | <i>Lethrinus amboinensis</i>       |
| 32809                                      | redgill emperor                 | <i>Lethrinus rubrioperculatus</i>  |
| 32310                                      | blueline snapper                | <i>Lutjanus kasmira</i>            |
| 32317                                      | yellowtail snapper              | <i>Pristipomoides auricilla</i>    |
| 32318                                      | pink snapper (paka)             | <i>Pristipomoides filamentosus</i> |
| 32319                                      | yelloweye snapper               | <i>Pristipomoides flavipinnis</i>  |
| 32320                                      | pink snapper (kalekale)         | <i>Pristipomoides seiboldi</i>     |
| 32321                                      | snapper (gindai)                | <i>Pristipomoides zonatus</i>      |
| 31414                                      | amberjack                       | <i>Seriola dumerili</i>            |

**2. Crustacean deep-water shrimp complex (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>   |
|--|---------------------|--------------------------|
| 67600                                      | deepwater shrimp    | <i>Heterocarpus</i> spp. |
| 67601                                      | deepwater shrimp    | <i>Pandalus unid</i> sp. |
| 67602                                      | deepwater shrimp    | Pandalidae               |
| 67603                                      | deepwater shrimp    | Pandalidae               |

**3. Crustacean spiny lobster complex (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>        |
|--|---------------------|-------------------------------|
| 67913                                      | spiny lobster       | <i>Panulirus marginatus</i>   |
| 67915                                      | spiny lobster       | <i>Panulirus penicillatus</i> |

**4. Crustacean slipper lobster complex (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b> |
|--|---------------------|------------------------|
| 67954                                      | slipper lobster     | Scyllaridae            |
| 67955                                      | slipper lobster     | Scyllaridae            |

**5. Crustacean Kona crab complex (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b> |
|--|---------------------|------------------------|
| 69150                                      | Kona crab           | <i>Ranina ranina</i>   |

**6. Precious coral black coral complex (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>      |
|--|---------------------|-----------------------------|
| none                                       | Black Coral         | <i>Anitpathes dichotoma</i> |
| none                                       | Black Coral         | <i>Antipathes grandis</i>   |
| none                                       | Black Coral         | <i>Antipathes ulex</i>      |

**7. Exploratory area precious coral (except black coral) (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b> | <b>Scientific Name</b>      |
|--|---------------------|-----------------------------|
| none                                       | Pink coral          | <i>Corallium secundum</i>   |
| none                                       | Pink coral          | <i>Corallium regale</i>     |
| none                                       | Pink coral          | <i>Corallium laauense</i>   |
| none                                       | Bamboo coral        | <i>Lepidisis olapa</i>      |
| none                                       | Bamboo coral        | <i>Acanella</i> spp.        |
| none                                       | Gold Coral          | <i>Gerardia</i> spp.        |
| none                                       | Gold Coral          | <i>Callogorgia gilberti</i> |
| none                                       | Gold Coral          | <i>Narella</i> spp.         |
| none                                       | Gold Coral          | <i>Calyptrophora</i> spp.   |

**8. Coral reef ecosystem (non-FSSI)**

| <b>DAWR<br/>Creel<br/>Species<br/>Code</b> | <b>Species Name</b>        | <b>Scientific Name</b>         | <b>Species grouping</b> |
|--|----------------------------|--------------------------------|-------------------------|
| 41201                                      | Achilles tang              | <i>Acanthurus achilles</i>     | Acanthuridae            |
| 41232                                      | Bariene's surgeonfish      | <i>Acanthurus bariene</i>      | Acanthuridae            |
| 41207                                      | Ringtail surgeonfish       | <i>Acanthurus blochii</i>      | Acanthuridae            |
| 41234                                      | Chronixis surgeonfish      | <i>Acanthurus chronixis</i>    | Acanthuridae            |
| 41202                                      | Eye-striped surgeonfish    | <i>Acanthurus dussumieri</i>   | Acanthuridae            |
| 41204                                      | Whitespotted surgeonfish   | <i>Acanthurus guttatus</i>     | Acanthuridae            |
| 41239                                      | Whitebar surgeonfish       | <i>Acanthurus leucocheilus</i> | Acanthuridae            |
| 41205                                      | Palelipped surgeonfish     | <i>Acanthurus leucopareius</i> | Acanthuridae            |
| 41206                                      | Blue-banded surgeonfish    | <i>Acanthurus lineatus</i>     | Acanthuridae            |
| 41235                                      | White-Freckled surgeonfish | <i>Acanthurus maculiceps</i>   | Acanthuridae            |
| 41233                                      | Elongate surgeonfish       | <i>Acanthurus mata</i>         | Acanthuridae            |
| 41203                                      | Whitecheek surgeonfish     | <i>Acanthurus nigricans</i>    | Acanthuridae            |
| 41208                                      | Blackstreak surgeonfish    | <i>Acanthurus nigricauda</i>   | Acanthuridae            |
| 41209                                      | Brown surgeonfish          | <i>Acanthurus nigrofuscus</i>  | Acanthuridae            |
| 41210                                      | Bluelined surgeonfish      | <i>Acanthurus nigroris</i>     | Acanthuridae            |
| 41240                                      | Surgeonfish                | <i>Acanthurus nubilus</i>      | Acanthuridae            |
| 41211                                      | Orangeband surgeonfish     | <i>Acanthurus olivaceus</i>    | Acanthuridae            |
| 41212                                      | Mimic surgeonfish          | <i>Acanthurus pyroferus</i>    | Acanthuridae            |

|       |                           |                                     |              |
|-------|---------------------------|-------------------------------------|--------------|
| 41243 | Surgeonfishes/tangs       | Acanthuridae                        | Acanthuridae |
| 41200 | Surgeonfishes/tangs       | Acanthuridae                        | Acanthuridae |
| 41213 | Thomson's surgeonfish     | <i>Acanthurus thompsoni</i>         | Acanthuridae |
| 41214 | Convict tang              | <i>Acanthurus triostegus</i>        | Acanthuridae |
| 41215 | Yellowfin surgeonfish     | <i>Acanthurus xanthopterus</i>      | Acanthuridae |
| 41216 | Twospot bristletooth      | <i>Ctenochaetus binotatus</i>       | Acanthuridae |
| 41217 | Black surgeonfish         | <i>Ctenochaetus hawaiiensis</i>     | Acanthuridae |
| 41236 | Blue-spotted Bristletooth | <i>Ctenochaetus marginatus</i>      | Acanthuridae |
| 41218 | Striped bristletooth      | <i>Ctenochaetus striatus</i>        | Acanthuridae |
| 41231 | Yellow-eyed bristletooth  | <i>Ctenochaetus strigosus</i>       | Acanthuridae |
| 41237 | Tomini's surgeonfish      | <i>Ctenochaetus tominiensis</i>     | Acanthuridae |
| 41219 | Whitemargin unicornfish   | <i>Naso annulatus</i>               | Acanthuridae |
| 41220 | Humpback unicornfish      | <i>Naso brachycentron</i>           | Acanthuridae |
| 41221 | Spotted unicornfish       | <i>Naso brevirostris</i>            | Acanthuridae |
| 41241 | Gray unicornfish          | <i>Naso caesius</i>                 | Acanthuridae |
| 41222 | Black tongue unicornfish  | <i>Naso hexacanthus</i>             | Acanthuridae |
| 41223 | Orangespine unicornfish   | <i>Naso lituratus</i>               | Acanthuridae |
| 41238 | Naso tang                 | <i>Naso lopezi</i>                  | Acanthuridae |
| 41242 | Barred unicornfish        | <i>Naso thynnoides</i>              | Acanthuridae |
| 41224 | Humpnose unicornfish      | <i>Naso tuberosus</i>               | Acanthuridae |
| 41225 | Bluespine unicornfish     | <i>Naso unicornis</i>               | Acanthuridae |
| 41226 | Bignose unicornfish       | <i>Naso vlamingii</i>               | Acanthuridae |
| 41227 | Hepatus tang              | <i>Paracanthurus hepatus</i>        | Acanthuridae |
| 41228 | Yellow tang               | <i>Zebrasoma flavescens</i>         | Acanthuridae |
| 41229 | Brown tang                | <i>Zebrasoma scopas</i>             | Acanthuridae |
| 41230 | Pacific sailfin tang      | <i>Zebrasoma veliferum</i>          | Acanthuridae |
| 31401 | Pennantfish/threadfin     | <i>Alectis ciliaris</i>             | Carangidae   |
| 31402 | Malabar Trevally          | <i>Alectis indicus</i>              | Carangidae   |
| 31400 | Jack (misc)               | Carangidae                          | Carangidae   |
| 31420 |                           | <i>Carangini</i>                    | Carangidae   |
| 31419 | Blue kingfish trevally    | <i>Carangoides caeruleopinnatus</i> | Carangidae   |
| 31431 | Shadow kingfish           | <i>Carangoides dinema</i>           | Carangidae   |
| 31422 | Bar jack                  | <i>Carangoides ferdau</i>           | Carangidae   |
| 31433 | Yellow dotted trevally    | <i>Carangoides fulvoguttatus</i>    | Carangidae   |
| 31438 | Headnotch trevally        | <i>Carangoides hedlandensis</i>     | Carangidae   |
| 31403 | Goldspot trevally         | <i>Carangoides orthogrammus</i>     | Carangidae   |
| 31424 | Barcheek trevally         | <i>Carangoides plagiotaenia</i>     | Carangidae   |
| 31425 | Jacks (misc)              | <i>Carangoides talamparoides</i>    | Carangidae   |
| 31437 | Trevally                  | <i>Carangoides uii</i>              | Carangidae   |
| 31429 | Trevally                  | <i>Caranx i'e'</i>                  | Carangidae   |

|       |                            |                                  |             |
|-------|----------------------------|----------------------------------|-------------|
| 31406 | Bluefin trevally           | <i>Caranx melampygu</i>          | Carangidae  |
| 31428 | Brassy trevally            | <i>Caranx papuensis</i>          | Carangidae  |
| 31407 | Bigeye trevally            | <i>Caranx sexfasciatus</i>       | Carangidae  |
| 31408 | Mackerel scad              | <i>Decapterus macarellus</i>     | Carangidae  |
| 31423 | Mackerel scad              | <i>Decapterus macrosoma</i>      | Carangidae  |
| 31421 | Round scad                 | <i>Decapterus maruadsi</i>       | Carangidae  |
| 31430 | Round scad                 | <i>Decapterus russelli</i>       | Carangidae  |
| 31409 | Rainbow runner             | <i>Elagatis bipinnulatus</i>     | Carangidae  |
| 31410 | Golden trevally            | <i>Gnathanodon speciosus</i>     | Carangidae  |
| 31439 |                            | <i>Megalaspis cordyla</i>        | Carangidae  |
| 31435 | Pilotfish                  | <i>Naucrates ductor</i>          | Carangidae  |
| 31440 | Elagatis, Scomberoides     | <i>Naucratini</i>                | Carangidae  |
| 31412 | Leatherback                | <i>Scomberoides lysan</i>        | Carangidae  |
| 31415 | Almaco jack                | <i>Seriola rivoliana</i>         | Carangidae  |
| 31416 | Small spotted pompano      | <i>Trachinotus bailloni</i>      | Carangidae  |
| 31417 | Silver or Snubnose pompano | <i>Trachinotus blochii</i>       | Carangidae  |
| 31432 | Mandibular kingfish        | <i>Ulua mandibularis</i>         | Carangidae  |
| 31418 | Kingfish                   | <i>Uraspis helvola</i>           | Carangidae  |
| 31436 | Deep trevally              | <i>Uraspis secunda</i>           | Carangidae  |
| 31434 | Whitemouth trevally        | <i>Uraspis uraspis</i>           | Carangidae  |
| 31413 | Atulai                     | <i>Selar crumenophthalmus</i>    | Atulai      |
| 31426 | Atulai                     | <i>Atule mate</i>                | Atulai      |
| 31427 | Atulai                     | <i>Selar boops</i>               | Atulai      |
| 32800 | Emperors                   | Lethrinidae                      | Lethrinidae |
| 32801 | Yellow-Spot Emperor        | <i>Gnathodentex aurolineatus</i> | Lethrinidae |
| 32802 | Grey Bream                 | <i>Gymnocranius griseus</i>      | Lethrinidae |
| 32804 | Thumbprint Emperor         | <i>Lethrinus harak</i>           | Lethrinidae |
| 32805 | Yellowtail Emperor         | <i>Lethrinus atkinsoni</i>       | Lethrinidae |
| 32806 | Longface Emperor           | <i>Lethrinus olivaceus</i>       | Lethrinidae |
| 32807 | Ornate Emperor             | <i>Lethrinus ornatus</i>         | Lethrinidae |
| 32808 | Orange-Striped Emperor     | <i>Lethrinus obsoletus</i>       | Lethrinidae |
| 32810 | Black-Blotch Emperor       | <i>Lethrinus semicinctus</i>     | Lethrinidae |
| 32811 | Yellowlip Emperor          | <i>Lethrinus xanthochilus</i>    | Lethrinidae |
| 32812 | Bigeye Emperor             | <i>Monotaxis grandoculus</i>     | Lethrinidae |
| 32813 | Japanese Bream             | <i>Gymnocranius euanus</i>       | Lethrinidae |
| 32814 | Orange-Spotted Emperor     | <i>Lethrinus erythracanthus</i>  | Lethrinidae |
| 32815 | Large-Eye Bream            | <i>Wattsia mossambica</i>        | Lethrinidae |
| 32816 | Stout Emperor              | <i>Gymnocranius sp</i>           | Lethrinidae |
| 32817 | Smtoothed Emperor          | <i>Lethrinus microdon</i>        | Lethrinidae |

|       |                        |                                 |             |
|-------|------------------------|---------------------------------|-------------|
| 32819 | Longspine Emperor      | <i>Lethrinus genivittatus</i>   | Lethrinidae |
| 32820 | Pinkear Emperor        | <i>Lethrinus lentjan</i>        | Lethrinidae |
| 32821 | Blue-Spotted Bream     | <i>Gymnocranius microdon</i>    | Lethrinidae |
| 32822 | Longfin Emperor        | <i>Lethrinus erythropterus</i>  | Lethrinidae |
| 32823 | Blue-Lined Bream       | <i>Gymnocranius grandoculus</i> | Lethrinidae |
| 32824 | Slender Emperor        | <i>Lethrinus variegatus</i>     | Lethrinidae |
| 36402 | Bucktooth Parrotfish   | <i>Calotomus carolinus</i>      | Scaridae    |
| 36420 | Spineytooth Parrotfish | <i>Calotomus spinidens</i>      | Scaridae    |
| 36403 | Bicolor Parrotfish     | <i>Cetoscarus bicolor</i>       | Scaridae    |
| 36422 | Parrotfish             | <i>Chlorurus bleekeri</i>       | Scaridae    |
| 36431 | Parrotfish             | <i>Chlorurus bowersi</i>        | Scaridae    |
| 36408 | Tan-Faced Parrotfish   | <i>Chlorurus frontalis</i>      | Scaridae    |
| 36410 | Steephead Parrotfish   | <i>Chlorurus microrhinos</i>    | Scaridae    |
| 36433 | Parrotfish             | <i>Chlorurus pyrrhurus</i>      | Scaridae    |
| 36416 | Bullethead Parrotfish  | <i>Chlorurus sordidus</i>       | Scaridae    |
| 36404 | Parrotfish             | <i>Hipposcarus longiceps</i>    | Scaridae    |
| 36405 | Seagrass Parrotfish    | <i>Leptoscarus vaigiensis</i>   | Scaridae    |
| 36400 | Parrotfishes           | Scaridae                        | Scaridae    |
| 36406 | Fil-Finned Parrotfish  | <i>Scarus altipinnis</i>        | Scaridae    |
| 36429 | Parrotfish             | <i>Scarus chameleon</i>         | Scaridae    |
| 36423 | Parrotfish             | <i>Scarus dimidiatus</i>        | Scaridae    |
| 36419 | Parrotfish             | <i>Scarus festivus</i>          | Scaridae    |
| 36434 | Yellowfin Parrotfish   | <i>Scarus flavipectoralis</i>   | Scaridae    |
| 36417 | Tricolor Parrotfish    | <i>Scarus forsteni</i>          | Scaridae    |
| 36407 | Vermiculate Parrotfish | <i>Scarus frenatus</i>          | Scaridae    |
| 36409 | Blue-Barred Parrotfish | <i>Scarus ghobban</i>           | Scaridae    |
| 36411 | Parrotfish             | <i>Scarus globiceps</i>         | Scaridae    |
| 36424 | Java Parrotfish        | <i>Scarus hypselosoma</i>       | Scaridae    |
| 36418 | Parrotfish             | <i>Scarus sp.</i>               | Scaridae    |
| 36432 | Black Parrotfish       | <i>Scarus niger</i>             | Scaridae    |
| 36412 | Parrotfish             | <i>Scarus oviceps</i>           | Scaridae    |
| 36425 | Greenthroat Parrotfish | <i>Scarus prasiognathos</i>     | Scaridae    |
| 36413 | Pale Nose Parrotfish   | <i>Scarus psittacus</i>         | Scaridae    |
| 36426 | Parrotfish             | <i>Scarus quoyi</i>             | Scaridae    |
| 36427 | Parrotfish             | <i>Scarus rivulatus</i>         | Scaridae    |
| 36414 | Parrotfish             | <i>Scarus rubroviolaceus</i>    | Scaridae    |
| 36415 | Chevron Parrotfish     | <i>Scarus schlegeli</i>         | Scaridae    |
| 36428 | Parrotfish             | <i>Scarus spinus</i>            | Scaridae    |
| 36435 | Tricolor Parrotfish    | <i>Scarus tricolor</i>          | Scaridae    |



|       |                         |                                     |          |
|-------|-------------------------|-------------------------------------|----------|
| 36421 | Parrotfish              | <i>Scarus xanthopleura</i>          | Scaridae |
| 33200 | Goatfishes              | Mullidae                            | Mullidae |
| 33201 | Yellowstriped Goatfish  | <i>Mulloidichthys flavolineatus</i> | Mullidae |
| 33202 | Orange Goatfish         | <i>Mulloidichthys pflugeri</i>      | Mullidae |
| 33219 | Juvenile Goatfish       | <i>Mulloidichthys ti'ao</i>         | Mullidae |
| 33203 | Yellowfin Goatfish      | <i>Mulloidichthys vanicolensis</i>  | Mullidae |
| 33216 |                         | <i>Parupeneus barberinoides</i>     | Mullidae |
| 33204 | Dash And Dot Goatfish   | <i>Parupeneus barberinus</i>        | Mullidae |
| 33205 |                         | <i>Parupeneus bifasciatus</i>       | Mullidae |
| 33210 | White-Lined Goatfish    | <i>Parupeneus ciliatus</i>          | Mullidae |
| 33206 | Yellow Goatfish         | <i>Parupeneus cyclostomus</i>       | Mullidae |
| 33208 | Redspot Goatfish        | <i>Parupeneus heptacanthus</i>      | Mullidae |
| 33214 | Indian Goatfish         | <i>Parupeneus indicus</i>           | Mullidae |
| 33211 | Multibarred Goatfish    | <i>Parupeneus multifasciatus</i>    | Mullidae |
| 33209 | Sidespot Goatfish       | <i>Parupeneus pleurostigma</i>      | Mullidae |
| 33217 | Goatfish                | <i>Parupeneus</i> sp.               | Mullidae |
| 33218 | Goatfish                | <i>Upeneus arge</i>                 | Mullidae |
| 33212 | Band-Tailed Goatfish    | <i>Upeneus taeniopterus</i>         | Mullidae |
| 33215 | Blackstriped Goatfish   | <i>Upeneus tragula</i>              | Mullidae |
| 33213 | Yellowbanded Goatfish   | <i>Upeneus vittatus</i>             | Mullidae |
| 54501 | Spiney Chiton           | <i>Acanthopleura spinosa</i>        | Mollusks |
| 54410 | Bubble Shells,Sea Hares | Acteonidae                          | Mollusks |
| 54603 | Antique Ark             | <i>Anadara antiquata</i>            | Mollusks |
| 54602 | Indo-Pacific Ark        | <i>Arca navicularis</i>             | Mollusks |
| 54601 | Ventricose Ark          | <i>Arca ventricosa</i>              | Mollusks |
| 54600 | Ark Shells              | Arcidae                             | Mollusks |
| 57742 | Common Paper Nautilus   | <i>Argonauta argo</i>               | Mollusks |
| 57745 | Gruner'S Paper Nautilus | <i>Argonauta gruneri</i>            | Mollusks |
| 57741 | Brown Paper Nautilus    | <i>Argonauta hians</i>              | Mollusks |
| 57743 | Nodose Paper Nautilus   | <i>Argonauta nodosa</i>             | Mollusks |
| 57744 | Noury'S Paper Nautilus  | <i>Argonauta nouri</i>              | Mollusks |
| 57740 | Paper Nautilus          | Argonautidae                        | Mollusks |
| 56896 | Pacific Sand Clam       | <i>Asaphis violescens</i>           | Mollusks |
| 56891 | Gaudy Sand Clam         | <i>Asaphis deflorata</i>            | Mollusks |
| 51751 | Peron'S Sea Butterfly   | <i>Atlanta peroni</i>               | Mollusks |
| 51750 |                         | Atlantidae                          | Mollusks |
| 54424 | Wh Pacific Aty          | <i>Atya naucum</i>                  | Mollusks |
| 54604 | Almond Ark              | <i>Babatia amygdalumtostum</i>      | Mollusks |
| 50840 | Goblets,Dwarf Tritons   | Buccinidae                          | Mollusks |

|       |                        |                               |          |
|-------|------------------------|-------------------------------|----------|
| 54421 | Ampule Bubble          | <i>Bulla ampulla</i>          | Mollusks |
| 54420 | Bubble Shells          | Bullidae                      | Mollusks |
| 54422 | Lined Bubble           | <i>Bullina lineata</i>        | Mollusks |
| 50796 | Giant Frog Shell       | <i>Bursa bubo</i>             | Mollusks |
| 50791 | Warty Frog Shell       | <i>Bursa bufonia</i>          | Mollusks |
| 50792 | Blood-Stain Frog Shell | <i>Bursa cruentata</i>        | Mollusks |
| 50793 | Granulate Frog Shell   | <i>Bursa granularis</i>       | Mollusks |
| 50799 | Lamarck'S Frog Shell   | <i>Bursa lamarcki</i>         | Mollusks |
| 50798 | Red-Mth Frog Shell     | <i>Bursa lissostoma</i>       | Mollusks |
| 50794 | Udder Frog Shell       | <i>Bursa mammata</i>          | Mollusks |
| 50797 | Ruddy Frog Shell       | <i>Bursa rebeta</i>           | Mollusks |
| 50795 | Wine-Mth Frog Shell    | <i>Bursa rhodostoma</i>       | Mollusks |
| 50790 | Frog Shells            | Bursidae                      | Mollusks |
| 50751 | Umbilicate Ovula       | <i>Calpurnus verrucosus</i>   | Mollusks |
| 50878 | File Miter             | <i>Cancilla filaris</i>       | Mollusks |
| 50842 | Smoky Goblet           | <i>Cantharus fumosus</i>      | Mollusks |
| 50841 | Waved Goblet           | <i>Cantharus undosus</i>      | Mollusks |
| 56721 | Varitated Cardita      | <i>Cardita variegata</i>      | Mollusks |
| 56720 | Carditid Clams         | Carditidae                    | Mollusks |
| 50767 | Vibex Bonnet           | <i>Casmaria erinaceus</i>     | Mollusks |
| 50768 | Heavy Bonnet           | <i>Casmaria ponderosa</i>     | Mollusks |
| 50765 | Helmet Shells          | Cassidae                      | Mollusks |
| 50766 | Horned Helmet          | <i>Cassius cornuta</i>        | Mollusks |
| 55022 | 3-Toothed Cavoline     | <i>Cavolina tridentata</i>    | Mollusks |
| 55023 | Unicate Cavoline       | <i>Cavolina uncinata</i>      | Mollusks |
| 55021 | Sea Butterfly          | <i>Cavolinia cf globulosa</i> | Mollusks |
| 55020 | Sea Butterflies        | Cavolinidae                   | Mollusks |
| 50650 | Turret, Worm-Shells    | Cerithiidae                   | Mollusks |
| 50654 | Column Certh           | <i>Cerithium columna</i>      | Mollusks |
| 50651 | Giant Knobbed Certh    | <i>Cerithium nodulosum</i>    | Mollusks |
| 56711 | Lazarus Jewel Box      | <i>Chama lazarus</i>          | Mollusks |
| 56710 | Jewel Boxes            | Chamidae                      | Mollusks |
| 50781 | Triton Trumpet         | <i>Charonia tritonis</i>      | Mollusks |
| 50812 | Ramose Murex           | <i>Chicoreus ramosus</i>      | Mollusks |
| 54500 | Chitons                | Chitonidae                    | Mollusks |
| 56623 | Cook'S Scallop         | <i>Chlamys cooki</i>          | Mollusks |
| 56621 | Squamose Scallop       | <i>Chlamys squamosa</i>       | Mollusks |
| 56500 | Bivalves               | Class Bivalvia                | Mollusks |
| 55027 | Pyramid Clio           | <i>Clio cuspidata</i>         | Mollusks |

|       |                        |                             |          |
|-------|------------------------|-----------------------------|----------|
| 55026 | Irregular Urchins      | <i>Clio pyramidata</i>      | Mollusks |
| 50652 | Morus Certh            | <i>Clypeomorus concisus</i> | Mollusks |
| 56706 | Punctate Lucina        | <i>Codakia punctata</i>     | Mollusks |
| 50847 | Maculated Dwarf Triton | <i>Columbraria muricata</i> | Mollusks |
| 50845 | Shiny Dwarf Triton     | <i>Columbraria nitidula</i> | Mollusks |
| 50846 | Twisted Dwarf Triton   | <i>Columbraria tortuosa</i> | Mollusks |
| 50920 | Cone Shells            | Conidae                     | Mollusks |
| 50952 | Sand-Dusted Cone       | <i>Conus arenatus</i>       | Mollusks |
| 50963 | Princely Cone          | <i>Conus aulicus</i>        | Mollusks |
| 50968 | Aureus Cone            | <i>Conus aureus</i>         | Mollusks |
| 50969 | Gold-Leaf Cone         | <i>Conus auricomus</i>      | Mollusks |
| 50947 | Banded Marble-Cone     | <i>Conus bandanus</i>       | Mollusks |
| 50971 | Bubble Cone            | <i>Conus bullatus</i>       | Mollusks |
| 50942 | Captain Cone           | <i>Conus capitaneus</i>     | Mollusks |
| 50932 | Cat Cone               | <i>Conus catus</i>          | Mollusks |
| 50924 | Chaldean Cone          | <i>Conus chaldeus</i>       | Mollusks |
| 50972 | Comma Cone             | <i>Conus connectens</i>     | Mollusks |
| 50922 | Crowned Cone           | <i>Conus coronatus</i>      | Mollusks |
| 50970 | Cylindrical Cone       | <i>Conus cylandraceus</i>   | Mollusks |
| 50926 | Distantly-Lined Cone   | <i>Conus distans</i>        | Mollusks |
| 50923 | Hebrew Cone            | <i>Conus ebraeus</i>        | Mollusks |
| 50936 | Ivory Cone             | <i>Conus eburneus</i>       | Mollusks |
| 50965 | Episcopus Cone         | <i>Conus episcopus</i>      | Mollusks |
| 50927 | Pacific Yellow Cone    | <i>Conus flavidus</i>       | Mollusks |
| 50928 | Frigid Cone            | <i>Conus frigidus</i>       | Mollusks |
| 50945 | General Cone           | <i>Conus generalis</i>      | Mollusks |
| 50961 | Geography Cone         | <i>Conus geographus</i>     | Mollusks |
| 50955 | Acorn Cone             | <i>Conus glans</i>          | Mollusks |
| 50946 | Imperial Cone          | <i>Conus imperialis</i>     | Mollusks |
| 50964 | Ambassador Cone        | <i>Conus legatus</i>        | Mollusks |
| 50938 | Leopard Cone           | <i>Conus leopardus</i>      | Mollusks |
| 50951 | Lithography Cone       | <i>Conus lithoglyphus</i>   | Mollusks |
| 50937 | Lettered Cone          | <i>Conus litteratus</i>     | Mollusks |
| 50929 | Livid Cone             | <i>Conus lividus</i>        | Mollusks |
| 50958 | Luteus Cone            | <i>Conus luteus</i>         | Mollusks |
| 50966 | Dignified Cone         | <i>Conus magnificus</i>     | Mollusks |
| 50930 | Soldier Cone           | <i>Conus miles</i>          | Mollusks |
| 50939 | 1000-Spot Cone         | <i>Conus miliaris</i>       | Mollusks |
| 50935 | Morelet'S Cone         | <i>Conus moreleti</i>       | Mollusks |

|       |                       |                                  |          |
|-------|-----------------------|----------------------------------|----------|
| 50934 | Muricate Cone         | <i>Conus muriculatus</i>         | Mollusks |
| 50940 | Music Cone            | <i>Conus musicus</i>             | Mollusks |
| 50943 | Weasel Cone           | <i>Conus mustelinus</i>          | Mollusks |
| 50954 | Obscure Cone          | <i>Conus obscurus</i>            | Mollusks |
| 50959 | Pertusus Cone         | <i>Conus pertusus</i>            | Mollusks |
| 50921 | Flea-Bite Cone        | <i>Conus pulicarius</i>          | Mollusks |
| 50931 | Rat Cone              | <i>Conus rattus</i>              | Mollusks |
| 50967 | Netted Cone           | <i>Conus retifer</i>             | Mollusks |
| 50933 | Blood-Stained Cone    | <i>Conus sanguinolentus</i>      | Mollusks |
| 50957 | Leaden Cone           | <i>Conus scabriusculus</i>       | Mollusks |
| 50925 | Marriage Cone         | <i>Conus sponsalis</i>           | Mollusks |
| 50950 | Striatellus Cone      | <i>Conus striatellus</i>         | Mollusks |
| 50948 | Striated Cone         | <i>Conus striatus</i>            | Mollusks |
| 50956 | Terebra Cone          | <i>Conus terebra</i>             | Mollusks |
| 50944 | Checkered Cone        | <i>Conus tessellatus</i>         | Mollusks |
| 50953 | Textile Cone          | <i>Conus textile</i>             | Mollusks |
| 50962 | Tulip Cone            | <i>Conus tulipa</i>              | Mollusks |
| 50960 | Varius Cone           | <i>Conus varius</i>              | Mollusks |
| 50941 | Flag Cone             | <i>Conus vexillum</i>            | Mollusks |
| 50949 | Calf Cone             | <i>Conus vitulinus</i>           | Mollusks |
| 50832 | Eroded Coral Shell    | <i>Coralliophila erosa</i>       | Mollusks |
| 50831 | Violet Coral Shell    | <i>Coralliophila neritodidea</i> | Mollusks |
| 50830 | Coral Shells          | <i>Coralliophilidae</i>          | Mollusks |
| 56662 | Giant Oyster          | <i>Crassostrea gigas</i>         | Mollusks |
| 56661 | Mangrove Oyster       | <i>Crassostrea mordax</i>        | Mollusks |
| 50813 | Bionic Rock Shell     | <i>Cronia biconica</i>           | Mollusks |
| 56624 | Speciosus Scallop     | <i>Cryptopecten speciosum</i>    | Mollusks |
| 55025 | Cigar Pteropod        | <i>Cuvierina columnella</i>      | Mollusks |
| 50770 | Tritons               | Cymatiidae                       | Mollusks |
| 50784 | Clandestine Triton    | <i>Cymatium clandestinum</i>     | Mollusks |
| 50773 | Jeweled Triton        | <i>Cymatium gemmatum</i>         | Mollusks |
| 50776 | Liver Triton          | <i>Cymatium hepaticum</i>        | Mollusks |
| 50786 | Wide-Lipped Triton    | <i>Cymatium labiosum</i>         | Mollusks |
| 50782 | Black-Spotted Triton  | <i>Cymatium lotorium</i>         | Mollusks |
| 50774 | Short-Neck Triton     | <i>Cymatium muricinum</i>        | Mollusks |
| 50772 | Nicobar Hairy Triton  | <i>Cymatium nicobaricum</i>      | Mollusks |
| 50779 | Common Hairy Triton   | <i>Cymatium pileare</i>          | Mollusks |
| 50771 | Aquatile Hairy Triton | <i>Cymatium pileare aquatile</i> | Mollusks |
| 50783 | Pear Triton           | <i>Cymatium pyrum</i>            | Mollusks |

|       |                    |                               |          |
|-------|--------------------|-------------------------------|----------|
| 50775 | Red Triton         | <i>Cymatium rubeculum</i>     | Mollusks |
| 50785 | Dwarf Hairy Triton | <i>Cymatium vespaceum</i>     | Mollusks |
| 50703 | Gold-Ringer Cowry  | <i>Cypraea annulus</i>        | Mollusks |
| 50726 | Arabian Cowry      | <i>Cypraea arabica</i>        | Mollusks |
| 50734 | Eyed Cowry         | <i>Cypraea argus</i>          | Mollusks |
| 50739 | Golden Cowry       | <i>Cypraea aurantium</i>      | Mollusks |
| 50738 | Beck'S Cowry       | <i>Cypraea beckii</i>         | Mollusks |
| 50733 | Bistro Cowry       | <i>Cypraea bistronatata</i>   | Mollusks |
| 50702 | Snake'S Head Cowry | <i>Cypraea caputserpentis</i> | Mollusks |
| 50710 | Carnelian Cowry    | <i>Cypraea carneola</i>       | Mollusks |
| 50740 | Chinese Cowry      | <i>Cypraea chinensis</i>      | Mollusks |
| 50732 | Chick-Pea Cowry    | <i>Cypraea cicercula</i>      | Mollusks |
| 50721 | Clandestine Cowry  | <i>Cypraea clandestina</i>    | Mollusks |
| 50715 | Sieve Cowry        | <i>Cypraea cribaria</i>       | Mollusks |
| 50713 | Sowerby'S Cowry    | <i>Cypraea cylindrica</i>     | Mollusks |
| 50717 | Depressed Cowry    | <i>Cypraea depressa</i>       | Mollusks |
| 50743 | Dillwyn'S Cowry    | <i>Cypraea dillywini</i>      | Mollusks |
| 50706 | Eglantine Cowry    | <i>Cypraea eglantina</i>      | Mollusks |
| 50708 | Eroded Cowry       | <i>Cypraea erosa</i>          | Mollusks |
| 50736 | Globular Cowry     | <i>Cypraea globulus</i>       | Mollusks |
| 50711 | Honey Cowry        | <i>Cypraea helvola</i>        | Mollusks |
| 50730 | Swallow Cowry      | <i>Cypraea hirundo</i>        | Mollusks |
| 50742 | Humphrey'S Cowry   | <i>Cypraea humphreysi</i>     | Mollusks |
| 50707 | Isabelle Cowry     | <i>Cypraea isabella</i>       | Mollusks |
| 50731 | Lined-Lip Cowry    | <i>Cypraea labrolineata</i>   | Mollusks |
| 50741 | Limacina Cowry     | <i>Cypraea limicina</i>       | Mollusks |
| 50704 | Lynx Cowry         | <i>Cypraea lynx</i>           | Mollusks |
| 50716 | Reticulated Cowry  | <i>Cypraea maculifera</i>     | Mollusks |
| 50705 | Map Cowry          | <i>Cypraea mappa</i>          | Mollusks |
| 50737 | Marie'S Cowry      | <i>Cypraea mariae</i>         | Mollusks |
| 50725 | Humpback Cowry     | <i>Cypraea mauritiana</i>     | Mollusks |
| 50723 | Microdon Cowry     | <i>Cypraea microdon</i>       | Mollusks |
| 50701 | Money Cowry        | <i>Cypraea moneta</i>         | Mollusks |
| 50722 | Nuclear Cowry      | <i>Cypraea nucleus</i>        | Mollusks |
| 50709 | Porus Cowry        | <i>Cypraea poraria</i>        | Mollusks |
| 50714 | Punctata Cowry     | <i>Cypraea punctata</i>       | Mollusks |
| 50729 | Jester Cowry       | <i>Cypraea scurra</i>         | Mollusks |
| 50712 | Grape Cowry        | <i>Cypraea staphlea</i>       | Mollusks |
| 50724 | Stolid Cowry       | <i>Cypraea stolid</i>         | Mollusks |

|       |                        |                                    |          |
|-------|------------------------|------------------------------------|----------|
| 50720 | Mole Cowry             | <i>Cypraea talpa</i>               | Mollusks |
| 50728 | Teres Cowry            | <i>Cypraea teres</i>               | Mollusks |
| 50718 | Tiger Cowry            | <i>Cypraea tigris</i>              | Mollusks |
| 50727 | Ventral Cowry          | <i>Cypraea ventriculus</i>         | Mollusks |
| 50719 | Pacific Deer Cowry     | <i>Cypraea vitellus</i>            | Mollusks |
| 50735 | Undulating Cowry       | <i>Cypraea ziczac</i>              | Mollusks |
| 50700 | Cowrys                 | Cypraeidae                         | Mollusks |
| 55024 | 3-Spined Cavoline      | <i>Diacria trispinosa</i>          | Mollusks |
| 50778 | Anal Triton            | <i>Distorso anus</i>               | Mollusks |
| 55100 | Dorid Nudibranchs      | Doridae                            | Mollusks |
| 50823 | Clatherate Drupe       | <i>Drupa clathrata</i>             | Mollusks |
| 50821 | Elegant Pacific Drupe  | <i>Drupa elegans</i>               | Mollusks |
| 50820 | Digitate Pacific Drupe | <i>Drupa grossularia</i>           | Mollusks |
| 50819 | Purple Pacific Drupe   | <i>Drupa morum</i>                 | Mollusks |
| 50818 | Prickley Pacific Drupe | <i>Drupa ricinus</i>               | Mollusks |
| 50822 | Strawberry Drupe       | <i>Drupa rubusidacaeus</i>         | Mollusks |
| 56622 | Spectacular Scallop    | <i>Excellichlamys spectiabilis</i> | Mollusks |
| 50850 | Spindles               | Fascioliariidae                    | Mollusks |
| 56722 | Pac Strawberry Cockle  | <i>Fragum fragum</i>               | Mollusks |
| 56908 | Tumid Venus            | <i>Gafrarium tumidum</i>           | Mollusks |
| 50777 | Rosy Gyre Triton       | <i>Gyrineum roseum</i>             | Mollusks |
| 50780 | Purple Gyre Triton     | <i>Gyrinium pusillum</i>           | Mollusks |
| 50911 | Little Love Harp       | <i>Harpa amouretta</i>             | Mollusks |
| 50913 | True Harp              | <i>Harpa harpa</i>                 | Mollusks |
| 50912 | Major Harp             | <i>Harpa major</i>                 | Mollusks |
| 50910 | Harp Shells            | Harpidae                           | Mollusks |
| 50989 | Lance Auger            | <i>Hastula lanceata</i>            | Mollusks |
| 50988 | Pencil Auger           | <i>Hastula penicillata</i>         | Mollusks |
| 55101 | Spanish Dancer         | <i>Hexabranthus sanguineus</i>     | Mollusks |
| 56881 | Giant Clam             | <i>Hippopus hippopus</i>           | Mollusks |
| 50806 | Anatomical Murex       | <i>Homalocanthia anatomica</i>     | Mollusks |
| 54423 | Gr-Lined Paber Bubble  | <i>Hydratina physis</i>            | Mollusks |
| 50875 | Cone-Like Miter        | <i>Imbricaria conularis</i>        | Mollusks |
| 50873 | Olive-Shaped Miter     | <i>Imbricaria olivaeformis</i>     | Mollusks |
| 50874 | Bonelike Miter         | <i>Imbricaria punctata</i>         | Mollusks |
| 56611 | Saddle Tree Oyster     | <i>Isognomon ephippium</i>         | Mollusks |
| 56610 | Tree Oysters           | Isognomonidae                      | Mollusks |
| 54351 | Janthina Snail         | <i>Janthina janthina</i>           | Mollusks |
| 54350 | Pelagic Snails         | Janthinidae                        | Mollusks |

|       |                       |                                 |          |
|-------|-----------------------|---------------------------------|----------|
| 50682 | Chiragra Spider Conch | <i>Lambis chiragra</i>          | Mollusks |
| 50685 | Ormouth Spider Conch  | <i>Lambis crocota</i>           | Mollusks |
| 50681 | Common Spider Conch   | <i>Lambis lambis</i>            | Mollusks |
| 50684 | Scorpio Conch         | <i>Lambis scorpius scorpius</i> | Mollusks |
| 50680 | Spider Conch          | <i>Lambis</i> sp.               | Mollusks |
| 50683 | Giant Spider Conch    | <i>Lambis truncata</i>          | Mollusks |
| 50851 | Nobby Spindle         | <i>Latirus nodatus</i>          | Mollusks |
| 50852 | Spindle               | <i>Latirus rudis</i>            | Mollusks |
| 56681 | Fragile Lima          | <i>Lima fragilis</i>            | Mollusks |
| 56682 | Indo-Pac Spiny Lima   | <i>Lima vulgaris</i>            | Mollusks |
| 56680 | Limas                 | Limidae                         | Mollusks |
| 56904 | Camp Pitar Venus      | <i>Lioconcha castrensis</i>     | Mollusks |
| 56906 | Hieroglyphic Venus    | <i>Lioconcha hieroglyphica</i>  | Mollusks |
| 56905 | Ornate Pitar Venus    | <i>Lioconcha ornata</i>         | Mollusks |
| 50642 | Scabra Periwinkle     | <i>Littorina scabra</i>         | Mollusks |
| 50641 | Undulate Periwinkle   | <i>Littorina undulata</i>       | Mollusks |
| 50640 | Periwinkles           | <i>Littorinidae</i>             | Mollusks |
| 56705 | Lucinas               | Lucinidae                       | Mollusks |
| 50762 | Apple Tun             | <i>Malea pomum</i>              | Mollusks |
| 50811 | Pinnacle Murex        | <i>Marchia bipinnatus</i>       | Mollusks |
| 50809 | Fenestrate Murex      | <i>Marchia martinetana</i>      | Mollusks |
| 54430 | Melampus Shells       | Melampidae                      | Mollusks |
| 54431 | Yellow Melampus       | <i>Melampus luteus</i>          | Mollusks |
| 57401 | Flamboyant Cuttlefish | <i>Metasepia pfefferi</i>       | Mollusks |
| 54425 | Mini Lined-Bubble     | <i>Micromelo undatus</i>        | Mollusks |
| 54411 | Ventricose Milda      | <i>Milda ventricosa</i>         | Mollusks |
| 56626 | Miraculous Scallop    | <i>Mirapekten mirificus</i>     | Mollusks |
| 50897 | Imperial Miter        | <i>Mitra imperialis</i>         | Mollusks |
| 50899 | Acuminate Miter       | <i>Mitra acuminata</i>          | Mollusks |
| 50890 | Cardinal Miter        | <i>Mitra cardinalis</i>         | Mollusks |
| 50893 | Chrysalis Miter       | <i>Mitra chrysalis</i>          | Mollusks |
| 50895 | Gold-Mth Miter        | <i>Mitra chrysostoma</i>        | Mollusks |
| 50889 | Coffee Miter          | <i>Mitra coffea</i>             | Mollusks |
| 50898 | Contracted Miter      | <i>Mitra contracta</i>          | Mollusks |
| 50892 | Kettle Miter          | <i>Mitra cucumaria</i>          | Mollusks |
| 50876 | Rusty Miter           | <i>Mitra ferruginea</i>         | Mollusks |
| 50891 | Strawberry Miter      | <i>Mitra fraga</i>              | Mollusks |
| 50888 | Tesselate Miter       | <i>Mitra incompta</i>           | Mollusks |
| 50872 | Episcopal Miter       | <i>Mitra mitra</i>              | Mollusks |

|       |                    |                                 |          |
|-------|--------------------|---------------------------------|----------|
| 50883 | Papal Miter        | <i>Mitra papalis</i>            | Mollusks |
| 50894 | Red-Painted Miter  | <i>Mitra rubitincta</i>         | Mollusks |
| 50871 | Pontifical Miter   | <i>Mitra stictica</i>           | Mollusks |
| 50870 | Miter Shells       | Mitridae                        | Mollusks |
| 50000 | Mollusca           | Mollusca                        | Mollusks |
| 50801 | Burnt Murex        | <i>Murex burneus</i>            | Mollusks |
| 50800 | Murex Shells       | <i>Muricidae</i>                | Mollusks |
| 56505 | Mussels            | <i>Mytilidae</i>                | Mollusks |
| 50804 | Tragonula Murex    | <i>Naquetia trigonulus</i>      | Mollusks |
| 50803 | Triquetra Murex    | <i>Naquetia triquetra</i>       | Mollusks |
| 50817 | Francolina Jopas   | <i>Nassa francolina</i>         | Mollusks |
| 50855 | Nassa Mud Snails   | Nassariidae                     | Mollusks |
| 50858 | Granulated Nassa   | <i>Nassarius graniferus</i>     | Mollusks |
| 50857 | Margarite Nassa    | <i>Nassarius margaritiferus</i> | Mollusks |
| 50856 | Pimpled Basket     | <i>Nassarius papillosus</i>     | Mollusks |
| 50755 | Moon Shells        | Naticidae                       | Mollusks |
| 57300 | Nautilus           | Nautilidae                      | Mollusks |
| 57301 | Chambered Nautilus | <i>Nautilus pompilius</i>       | Mollusks |
| 50884 | Clathrus Miter     | <i>Neocancilla clathrus</i>     | Mollusks |
| 50896 | Flecked Miter      | <i>Neocancilla granitina</i>    | Mollusks |
| 50901 | Butterfly Miter    | <i>Neocancilla papilio</i>      | Mollusks |
| 50633 | Ox-Palate Nerite   | <i>Nerita albicilla</i>         | Mollusks |
| 50631 | Plicate Nerite     | <i>Nerita plicata</i>           | Mollusks |
| 50632 | Polished Nerite    | <i>Nerita polita</i>            | Mollusks |
| 50634 | Reticulate Nerite  | <i>Nerita signata</i>           | Mollusks |
| 50630 | Nerites            | Neritidae                       | Mollusks |
| 50600 | Diotocardia        | O Archaeogastropoda             | Mollusks |
| 57700 | Octopus            | Octopodidae                     | Mollusks |
| 57735 | Common Octopus     | <i>Octopus cyanea</i>           | Mollusks |
| 57734 | Red Octopus        | <i>Octopus luteus</i>           | Mollusks |
| 57736 | Ornate Octopus     | <i>Octopus ornatus</i>          | Mollusks |
| 57730 | Octopus            | <i>Octopus sp.</i>              | Mollusks |
| 57732 | Pelagic Octopus    | <i>Octopus sp.</i>              | Mollusks |
| 57733 | Long-Armed Octopus | <i>Octopus sp.</i>              | Mollusks |
| 57731 | Elongate Octopus   | <i>Octopus teuthoides</i>       | Mollusks |
| 50861 | Amethyst Olive     | <i>Oliva annulata</i>           | Mollusks |
| 50863 | Carnelian Olive    | <i>Oliva carneola</i>           | Mollusks |
| 50862 | Red-Mth Olive      | <i>Oliva miniacea</i>           | Mollusks |
| 50864 | Peg Olive          | <i>Oliva paxillus</i>           | Mollusks |



|       |                      |                                |          |
|-------|----------------------|--------------------------------|----------|
| 50860 | Olive Shells         | Olividae                       | Mollusks |
| 57500 | Squids               | <i>Order Teuthoidea</i>        | Mollusks |
| 56660 | True Oysters         | Ostreidae                      | Mollusks |
| 54412 | Cat'S Ear Otopleura  | <i>Otopleura auriscati</i>     | Mollusks |
| 50753 | Common Egg Cowry     | <i>Ovula ovum</i>              | Mollusks |
| 50750 | Egg Shells           | Ovulidae                       | Mollusks |
| 56620 | Scallops             | Pectinidae                     | Mollusks |
| 56902 | Crispate Venus       | <i>Periglypta crispata</i>     | Mollusks |
| 56903 | Youthful Venus       | <i>Periglypta puerpera</i>     | Mollusks |
| 56901 | Reticulate Venus     | <i>Periglypta reticulata</i>   | Mollusks |
| 56601 | Pearl Oyster         | <i>Pinctada margaritifera</i>  | Mollusks |
| 56531 | Bicolor Pen Shell    | <i>Pinna bicolor</i>           | Mollusks |
| 56530 | Pen Shells           | Pinnidae                       | Mollusks |
| 50756 | Breast-Shaped Moon   | <i>Polinices mamatus</i>       | Mollusks |
| 50757 | Pear-Shaped Moon     | <i>Polinices tumidus</i>       | Mollusks |
| 50844 | Strawberry Goblet    | <i>Pollia fragaria</i>         | Mollusks |
| 50843 | Beautiful Goblet     | <i>Pollia pulchra</i>          | Mollusks |
| 50752 | Fruit Ovula          | <i>Prionovula fruticum</i>     | Mollusks |
| 56600 | Pearl Oysters        | Pteriidae                      | Mollusks |
| 50904 | Crenulate Miter      | <i>Pterygia crenulata</i>      | Mollusks |
| 50907 | Fenestrate Miter     | <i>Pterygia fenestrata</i>     | Mollusks |
| 50905 | Nut Miter            | <i>Pterygia nucea</i>          | Mollusks |
| 50902 | Rough Miter          | <i>Pterygia scabricula</i>     | Mollusks |
| 50810 | Club Murex           | <i>Pterynotus elongatus</i>    | Mollusks |
| 50807 | Fluted Murex         | <i>Pterynotus laqueatus</i>    | Mollusks |
| 50808 | 3-Winged Murex       | <i>Pterynotus tripterus</i>    | Mollusks |
| 54413 | Solid Pupa           | <i>Pupa solidula</i>           | Mollusks |
| 50816 | Perssian Purpura     | <i>Purpura persica</i>         | Mollusks |
| 54401 | Sulcate Pyram        | <i>Pyramidella sulcata</i>     | Mollusks |
| 54400 | Pyram Shells         | Pyramidellidae                 | Mollusks |
| 50833 | Quoy'S Coral Shell   | <i>Quoyula madreporarum</i>    | Mollusks |
| 50834 | Rapa Snail           | <i>Rapa rapa</i>               | Mollusks |
| 50653 | Rough Vertigus       | <i>Rhinoclavis aspera</i>      | Mollusks |
| 50655 | Obelisk Vertigus     | <i>Rhinoclavis sinensis</i>    | Mollusks |
| 50900 | Chaste Miter         | <i>Sabricula casta</i>         | Mollusks |
| 56625 | Tiger Scallop        | <i>Semipallium tigris</i>      | Mollusks |
| 57403 | Broadclub Cuttlefish | <i>Sepia latimanus</i>         | Mollusks |
| 57402 | Cuttlefish           | <i>Sepia</i> sp.               | Mollusks |
| 57594 | Bigfin Reef Squid    | <i>Sepioteuthis lessoniana</i> | Mollusks |

|       |                      |                              |          |
|-------|----------------------|------------------------------|----------|
| 56511 | Box Mussel           | <i>Septifer bilocularis</i>  | Mollusks |
| 50805 | Lacy Murex           | <i>Siratus laciniatus</i>    | Mollusks |
| 56670 | Thorny Oysters       | Spondylidae                  | Mollusks |
| 56671 | Ducal Thorny Oyster  | <i>Spondylus squamosus</i>   | Mollusks |
| 56532 | Baggy Pen Shell      | <i>Streptopinna saccata</i>  | Mollusks |
| 50660 | True Conchs          | Strombidae                   | Mollusks |
| 50665 | Samar Conch          | <i>Strombus dentatus</i>     | Mollusks |
| 50666 | Fragile Conch        | <i>Strombus fragilis</i>     | Mollusks |
| 50663 | Gibbose Conch        | <i>Strombus gibberulus</i>   | Mollusks |
| 50669 | Lavender-Mouth Conch | <i>Strombus haemastoma</i>   | Mollusks |
| 50667 | Silver-Lip Conch     | <i>Strombus lentiginosus</i> | Mollusks |
| 50662 | Red-Lip Conch        | <i>Strombus luhuanus</i>     | Mollusks |
| 50664 | Micro Conch          | <i>Strombus microurceus</i>  | Mollusks |
| 50661 | Mutable Conch        | <i>Strombus mutabilis</i>    | Mollusks |
| 50672 | Pretty Conch         | <i>Strombus plicatus</i>     | Mollusks |
| 50670 | Lacinate Conch       | <i>Strombus sinuatus</i>     | Mollusks |
| 50671 | Bull Conch           | <i>Strombus taurus</i>       | Mollusks |
| 50612 | Pyramid Top          | <i>Tectus pyramis</i>        | Mollusks |
| 56894 | Box-Like Tellin      | <i>Tellina capsoides</i>     | Mollusks |
| 56892 | Cat'S Tongue Tellin  | <i>Tellina linguafelis</i>   | Mollusks |
| 56895 | Remie'S Tellin       | <i>Tellina remies</i>        | Mollusks |
| 56893 | Rasp Tellin          | <i>Tellina scobinata</i>     | Mollusks |
| 56890 | Tellin Clams         | Tellinidae                   | Mollusks |
| 50668 | Terebellum Conch     | <i>Terebellum terebellum</i> | Mollusks |
| 50985 | Similar Auger        | <i>Terebra affinis</i>       | Mollusks |
| 50997 | Fly-Spotted Auger    | <i>Terebra areolata</i>      | Mollusks |
| 50996 | Eyed Auger           | <i>Terebra argus</i>         | Mollusks |
| 50987 | Babylonian Auger     | <i>Terebra babylonica</i>    | Mollusks |
| 50990 | Certhlike Auger      | <i>Terebra cerithiana</i>    | Mollusks |
| 50995 | Short Auger          | <i>Terebra chlorata</i>      | Mollusks |
| 50984 | Crenulated Auger     | <i>Terebra crenulata</i>     | Mollusks |
| 50982 | Dimidiate Auger      | <i>Terebra dimidiata</i>     | Mollusks |
| 50994 | Tiger Auger          | <i>Terebra felina</i>        | Mollusks |
| 50991 | Funnel Auger         | <i>Terebra funiculata</i>    | Mollusks |
| 50993 | Spotted Auger        | <i>Terebra gutatta</i>       | Mollusks |
| 50981 | Marlinspike Auger    | <i>Terebra maculata</i>      | Mollusks |
| 50986 | Cloud Auger          | <i>Terebra nubulosa</i>      | Mollusks |
| 50983 | Subulate Auger       | <i>Terebra subulata</i>      | Mollusks |
| 50992 | Undulate Auger       | <i>Terebra undulata</i>      | Mollusks |

|       |                               |                                 |           |
|-------|-------------------------------|---------------------------------|-----------|
| 50980 | Auger Shells                  | Terebridae                      | Mollusks  |
| 50815 | Belligerent Rock Shell        | <i>Thais armigera</i>           | Mollusks  |
| 50814 | Tuberose Rock Shell           | <i>Thais tuberosa</i>           | Mollusks  |
| 50761 | Partridge Tun                 | <i>Tonna perdix</i>             | Mollusks  |
| 50760 | Tun Shells                    | Tonnidae                        | Mollusks  |
| 56723 | Angulate Cockle               | <i>Trachycardium angulatum</i>  | Mollusks  |
| 56882 | Giant Clam                    | <i>Tridacna crocea</i>          | Mollusks  |
| 56883 | Lagoon Giant Clam             | <i>Tridacna derasa</i>          | Mollusks  |
| 56884 | Giant Clam                    | <i>Tridacna gigas</i>           | Mollusks  |
| 56885 | Common Giant Clam             | <i>Tridacna maxima</i>          | Mollusks  |
| 56886 | Fluted Giant Clam             | <i>Tridacna squamosa</i>        | Mollusks  |
| 56880 | Giant Clams                   | Tridacnidae                     | Mollusks  |
| 50610 | Top Shells                    | Trochidae                       | Mollusks  |
| 50611 | Top Shell                     | <i>Trochus niloticus</i>        | Mollusks  |
| 50613 | Radiate Top                   | <i>Trochus radiatus</i>         | Mollusks  |
| 50865 | Vases                         | Turbinellidae                   | Mollusks  |
| 50620 | Turban Shell                  | Turbinidae                      | Mollusks  |
| 50622 | Silver-Mouth Turbin           | <i>Turbo argyrostoma</i>        | Mollusks  |
| 50623 | Tapestry Turbin               | <i>Turbo petholatus</i>         | Mollusks  |
| 50621 | Rough Turbin                  | <i>Turbo setosus</i>            | Mollusks  |
| 50867 | Ceramic Vase                  | <i>Vasum ceramicum</i>          | Mollusks  |
| 50866 | Common Pacific Vase           | <i>Vasum turbinellus</i>        | Mollusks  |
| 56900 | Venus Shells                  | Veneridae                       | Mollusks  |
| 50887 | Bernhard'S Miter              | <i>Vexillum bernhardiana</i>    | Mollusks  |
| 50882 | Cancellaria Miter             | <i>Vexillum cancellarioides</i> | Mollusks  |
| 50880 | Saffron Miter                 | <i>Vexillum crocatum</i>        | Mollusks  |
| 50879 | Roughened Miter               | <i>Vexillum exasperatum</i>     | Mollusks  |
| 50885 | Patriarchal Miter             | <i>Vexillum patriarchalis</i>   | Mollusks  |
| 50881 | Half-Banded Miter             | <i>Vexillum semifasciatum</i>   | Mollusks  |
| 50903 | Specious Miter                | <i>Vexillum speciosum</i>       | Mollusks  |
| 50886 | Bumpy Miter                   | <i>Vexillum tuberosum</i>       | Mollusks  |
| 50906 | Turbin Miter                  | <i>Vexillum turbin</i>          | Mollusks  |
| 50877 | Decorated Miter               | <i>Vexillum unifasciatum</i>    | Mollusks  |
| 50802 | Spotted Vitularia             | <i>Vitularia miliaris</i>       | Mollusks  |
| 41316 | Manahak (Forktail Rabbitfish) | <i>Manahak lessor'</i>          | Siganidae |
| 41318 | Manahak                       | <i>Manahak</i> sp.              | Siganidae |
| 41300 | Rabbitfish                    | Siganidae                       | Siganidae |
| 41301 | Fork-Tail Rabbitfish          | <i>Siganus argenteus</i>        | Siganidae |
| 41307 | Seagrass Rabbitfish           | <i>Siganus canaliculatus</i>    | Siganidae |

|       |                            |                                    |            |
|-------|----------------------------|------------------------------------|------------|
| 41308 | Coral Rabbitfish           | <i>Siganus corallinus</i>          | Siganidae  |
| 41302 | Pencil-Streaked Rabbitfish | <i>Siganus doliatus</i>            | Siganidae  |
| 41303 | Fuscescens Rabbitfish      | <i>Siganus fuscescens</i>          | Siganidae  |
| 41309 | Golden Rabbitfish          | <i>Siganus guttatus</i>            | Siganidae  |
| 41311 | Lined Rabbitfish           | <i>Siganus lineatus</i>            | Siganidae  |
| 41313 | White-Spotted Rabbitfish   | <i>Siganus oramin</i>              | Siganidae  |
| 41310 | Masked Rabbitfish          | <i>Siganus puellus</i>             | Siganidae  |
| 41314 | Peppered Rabbitfish        | <i>Siganus punctatissimus</i>      | Siganidae  |
| 41304 | Gold-Spotted Rabbitfish    | <i>Siganus punctatus</i>           | Siganidae  |
| 41315 | Randal'S Rabbitfish        | <i>Siganus randalli</i>            | Siganidae  |
| 41305 | Scribbled Rabbitfish       | <i>Siganus spinus</i>              | Siganidae  |
| 41306 | Vermiculated Rabbitfish    | <i>Siganus vermiculatus</i>        | Siganidae  |
| 41312 | Rabbitfish                 | <i>Siganus vulpinus</i>            | Siganidae  |
| 32301 | Silvermouth/Jobfish        | <i>Aphareus furca</i>              | Lutjanidae |
| 32300 | Snappers                   | Lutjanidae                         | Lutjanidae |
| 32306 | River Snapper              | <i>Lutjanus argentimaculatus</i>   | Lutjanidae |
| 32325 | Two-Spot Snapper           | <i>Lutjanus biguttatus</i>         | Lutjanidae |
| 32307 | Red Snapper                | <i>Lutjanus bohar</i>              | Lutjanidae |
| 32334 | Snapper                    | <i>Lutjanus bouton</i>             | Lutjanidae |
| 32326 | Checkered Snapper          | <i>Lutjanus decussatus</i>         | Lutjanidae |
| 32327 | Blackspot Snapper          | <i>Lutjanus ehrenbergi</i>         | Lutjanidae |
| 32335 | Snapper                    | <i>Lutjanus fulviflamma</i>        | Lutjanidae |
| 32308 | Flametail Snapper          | <i>Lutjanus fulvus</i>             | Lutjanidae |
| 32309 | Humpback Snapper           | <i>Lutjanus gibbus</i>             | Lutjanidae |
| 32328 | Malabar Snapper            | <i>Lutjanus malabaricus</i>        | Lutjanidae |
| 32312 | Onespot Snapper            | <i>Lutjanus monostigma</i>         | Lutjanidae |
| 32311 | Scribbled Snapper          | <i>Lutjanus rivulatus</i>          | Lutjanidae |
| 32333 | Snapper                    | <i>Lutjanus sebae</i>              | Lutjanidae |
| 32329 | 1/2-Barred Snapper         | <i>Lutjanus semicinctus</i>        | Lutjanidae |
| 32330 | One-Lined Snapper          | <i>Lutjanus vitta</i>              | Lutjanidae |
| 32332 | Bl And Wh Snapper          | <i>Macolor macularis</i>           | Lutjanidae |
| 32313 | Black Snapper              | <i>Macolor niger</i>               | Lutjanidae |
| 32314 | Fusilier                   | <i>Paracaesio sordidus</i>         | Lutjanidae |
| 32315 | Yellowtail Fusilier        | <i>Paracaesio xanthurus</i>        | Lutjanidae |
| 32322 | Deepwater Snapper          | <i>Randallichthys filamentosus</i> | Lutjanidae |
| 49130 | Shallow Snappers           | Shallow Snappers                   | Lutjanidae |
| 32331 | Sailfin Snapper            | <i>Symphoricichthys spilurus</i>   | Lutjanidae |
| 28901 | Red-Flushed Grouper        | <i>Aethaloperca rogae</i>          | Serranidae |
| 28956 | Grouper                    | <i>Anyperodon leucogrammicus</i>   | Serranidae |

|       |                              |  |            |
|-------|------------------------------|--|------------|
| 28908 | Orange Grouper               | <i>Cephalopholis analis</i>              | Serranidae |
| 28907 | Peacock Grouper              | <i>Cephalopholis argus</i>               | Serranidae |
| 28911 | Brownbarred Grouper          | <i>Cephalopholis boenack</i>             | Serranidae |
| 28909 | Ybanded Grouper              | <i>Cephalopholis igarashiensis</i>       | Serranidae |
| 28910 | Leopard Grouper              | <i>Cephalopholis leopardus</i>           | Serranidae |
| 28945 | Coral Grouper                | <i>Cephalopholis miniata</i>             | Serranidae |
| 28929 | Harlequin Grouper            | <i>Cephalopholis polleni</i>             | Serranidae |
| 28913 | 6-Banded Grouper             | <i>Cephalopholis sexmaculata</i>         | Serranidae |
| 28912 | Tomato Grouper               | <i>Cephalopholis sonnerati</i>           | Serranidae |
| 28903 | Grouper                      | <i>Cephalopholis sp</i>                  | Serranidae |
| 28906 | Pygmy Grouper                | <i>Cephalopholis spiloparaea</i>         | Serranidae |
| 28914 | Flag-Tailed Grouper          | <i>Cephalopholis urodeta</i>             | Serranidae |
| 28915 | Grouper                      | <i>Cromileptes altivelis</i>             | Serranidae |
| 28947 | Orange Grouper               | <i>Epinephelus<br/>caeruleopunctatus</i> | Serranidae |
| 28948 | Brown-Spotted Grouper        | <i>Epinephelus chlorostigma</i>          | Serranidae |
| 28960 | Orange Spot Grouper          | <i>Epinephelus coioides</i>              | Serranidae |
| 28957 | Grouper                      | <i>Epinephelus corallicola</i>           | Serranidae |
| 28946 | Grouper                      | <i>Epinephelus cyanopodus</i>            | Serranidae |
| 28920 | Blotchy Grouper              | <i>Epinephelus fuscoguttatus</i>         | Serranidae |
| 28921 | Hexagon Grouper              | <i>Epinephelus hexagonatus</i>           | Serranidae |
| 28918 | Grouper                      | <i>Epinephelus howlandi</i>              | Serranidae |
| 28922 | Giant Grouper                | <i>Epinephelus lanceolatus</i>           | Serranidae |
| 28958 | Grouper                      | <i>Epinephelus macrospilos</i>           | Serranidae |
| 28923 | Highfin Grouper              | <i>Epinephelus maculatus</i>             | Serranidae |
| 28950 | Malabar Grouper              | <i>Epinephelus malabaricus</i>           | Serranidae |
| 28949 | Bl-Spot Honeycomb<br>Grouper | <i>Epinephelus melanostigma</i>          | Serranidae |
| 28925 | Honeycomb Grouper            | <i>Epinephelus merra</i>                 | Serranidae |
| 28942 | Grouper                      | <i>Epinephelus miliaris</i>              | Serranidae |
| 28916 | Grouper                      | <i>Epinephelus morrhua</i>               | Serranidae |
| 28951 | Wavy-Lined Grouper           | <i>Epinephelus ongus</i>                 | Serranidae |
| 28926 | Marbled Grouper              | <i>Epinephelus polyphkadion</i>          | Serranidae |
| 28953 | Grouper                      | <i>Epinephelus retouti</i>               | Serranidae |
| 28930 | 7-Banded Grouper             | <i>Epinephelus<br/>septemfasciatus</i>   | Serranidae |
| 28924 | Tidepool Grouper             | <i>Epinephelus socialis</i>              | Serranidae |
| 28952 | 4-Saddle Grouper             | <i>Epinephelus spilotoceps</i>           | Serranidae |
| 28928 | Greasy Grouper               | <i>Epinephelus tauvina</i>               | Serranidae |
| 28902 | Truncated Grouper            | <i>Epinephelus truncatus</i>             | Serranidae |

|       |                           |                                  |                 |
|-------|---------------------------|----------------------------------|-----------------|
| 28943 | Wh-Margined Grouper       | <i>Gracila albomarginata</i>     | Serranidae      |
| 28938 | Squairetail Grouper       | <i>Plectropomus areolatus</i>    | Serranidae      |
| 28937 | Saddleback Grouper        | <i>Plectropomus laevis</i>       | Serranidae      |
| 28954 | Leopard Coral Trout       | <i>Plectropomus leopardus</i>    | Serranidae      |
| 28955 | Blue-Lined Coral Trout    | <i>Plectropomus oligacanthus</i> | Serranidae      |
| 28940 | Powell'S Grouper          | <i>Saloptia powelli</i>          | Serranidae      |
| 28900 | Sea Basses,Groupers       | Serranidae                       | Serranidae      |
| 28944 | Whmargin Lyretail Grouper | <i>Variola albimarginata</i>     | Serranidae      |
| 35902 | Fringelip Mullet          | <i>Crenimugil crenilabis</i>     | Mugilidae       |
| 35903 | Yellowtail Mullet         | <i>Ellochelon vaigiensis</i>     | Mugilidae       |
| 35901 | Engel'S Mullet            | <i>Moolgarda engeli</i>          | Mugilidae       |
| 35906 | Bluespot Mullet           | <i>Moolgarda seheli</i>          | Mugilidae       |
| 35904 | Gray Mullet               | <i>Mugil cephalus</i>            | Mugilidae       |
| 35900 | Mullets                   | Mugilidae                        | Mugilidae       |
| 35905 | Acute-Jawed Mullet        | <i>Neomyxus leuciscus</i>        | Mugilidae       |
| 33900 | Rudderfish                | Kyphosidae                       | Kyphosidae      |
| 33901 | Highfin Rudderfish        | <i>Kyphosus cinerascens</i>      | Kyphosidae      |
| 33902 | Lowfin Rudderfish         | <i>Kyphosus vaigiensis</i>       | Kyphosidae      |
| 33903 | Insular Rudderfish        | <i>Kyphosus bigibbus</i>         | Kyphosidae      |
| 69251 | Spider Crab               | <i>Achaeus japonicus</i>         | CRE-Crustaceans |
| 67500 | Snapping Shrimp           | Alpheidae                        | CRE-Crustaceans |
| 67501 | Snapping Shrimp           | <i>Alpheus bellulus</i>          | CRE-Crustaceans |
| 67502 | Snapping Shrimp           | <i>Alpheus paracrinitus</i>      | CRE-Crustaceans |
| 64999 | Anchylomerids             | <i>Anchylomeridae</i>            | CRE-Crustaceans |
| 67951 | Slipper Lobster           | <i>Arctides regalis</i>          | CRE-Crustaceans |
| 60101 | Acorn Barnacle            | <i>Balanus</i> sp                | CRE-Crustaceans |
| 62050 | Mantis Shrimp             | <i>Bathysquillidae</i>           | CRE-Crustaceans |
| 69201 | Box Crab                  | <i>Calappa bicornis</i>          | CRE-Crustaceans |
| 69202 | Box Crab                  | <i>Calappa calappa</i>           | CRE-Crustaceans |
| 69203 | Box Crab                  | <i>Calappa hepatica</i>          | CRE-Crustaceans |
| 69200 | Box Crabs                 | Calappidae                       | CRE-Crustaceans |
| 69252 | Decorator Crab            | <i>Camposcia retusa</i>          | CRE-Crustaceans |
| 69350 | Cancrids                  | Cancridae                        | CRE-Crustaceans |
| 69501 | 7-11 Crab                 | <i>Carpilius convexus</i>        | CRE-Crustaceans |
| 69502 | 7-11 Crab                 | <i>Carpilius maculatus</i>       | CRE-Crustaceans |
| 69401 | Red-Legged Sw Crab        | <i>Charybdis erythroductyla</i>  | CRE-Crustaceans |
| 69402 | Red Sw Crab               | <i>Charybdis hawaiiensis</i>     | CRE-Crustaceans |
| 69204 | Box Crab                  | <i>Cycloes granulosa</i>         | CRE-Crustaceans |
| 69301 | Elbow Crab                | <i>Daldorfia horrida</i>         | CRE-Crustaceans |

|       |                           |                                     |                 |
|-------|---------------------------|-------------------------------------|-----------------|
| 68202 | Marine Hermit Crab        | <i>Dardanus gemmatus</i>            | CRE-Crustaceans |
| 68204 | Marine Hermit Crab        | <i>Dardanus megistos</i>            | CRE-Crustaceans |
| 68203 | Marine Hermit Crab        | <i>Dardanus pendunculatus</i>       | CRE-Crustaceans |
| 68201 | Marine Hermit Crab        | <i>Dardanus</i> sp                  | CRE-Crustaceans |
| 67121 | Commensal Shrimp          | <i>Dasycaris zanzibarica</i>        | CRE-Crustaceans |
| 67000 | Decapod Crustaceans       | Decapoda                            | CRE-Crustaceans |
| 68200 | Marine Hermit Crabs       | Diogenidae                          | CRE-Crustaceans |
| 69161 | Dorippid Crab             | <i>Dorippe frascone</i>             | CRE-Crustaceans |
| 69171 | Sponge Crab               | <i>Dromia dormia</i>                | CRE-Crustaceans |
| 69170 | Sponge Crabs              | Dromiidae                           | CRE-Crustaceans |
| 68701 | Mole Crab                 | <i>Emerita pacifica</i>             | CRE-Crustaceans |
| 67851 | Soft Lobster              | <i>Enoplometopus debelius</i>       | CRE-Crustaceans |
| 67852 | Hairy Lobster             | <i>Enoplometopus occidentalis</i>   | CRE-Crustaceans |
| 69553 | Redeye Crab               | <i>Eriphia sebana</i>               | CRE-Crustaceans |
| 69554 | Red-Reef Crab             | <i>Etisus dentatus</i>              | CRE-Crustaceans |
| 69551 | Red-Reef Crab             | <i>Etisus splendidus</i>            | CRE-Crustaceans |
| 69555 | Brown-Reef Crab           | <i>Etisus utilis</i>                | CRE-Crustaceans |
| 62100 | Mantis Shrimp             | Eurysquillidae                      | CRE-Crustaceans |
| 68500 | Squat Lobsters            | Galatheidae                         | CRE-Crustaceans |
| 69850 | Gecarcinids               | Gecarcinidae                        | CRE-Crustaceans |
| 67220 | Bbee And Harlequin Shrimp | Gnathophyllidae                     | CRE-Crustaceans |
| 67221 | Bumblebee Shrimp          | <i>Gnathophylloides mineri</i>      | CRE-Crustaceans |
| 67222 | Bumblebee Shrimp          | <i>Gnathophyllum americanum</i>     | CRE-Crustaceans |
| 62203 | Mantis Shrimp             | <i>Gonodactylaceus mutatus</i>      | CRE-Crustaceans |
| 62201 | Mantis Shrimp             | <i>Gonodactylellus affinis</i>      | CRE-Crustaceans |
| 62200 | Mantis Shrimp             | Gonodactylidae                      | CRE-Crustaceans |
| 62202 | Mantis Shrimp             | <i>Gonodactylus chiragra</i>        | CRE-Crustaceans |
| 62204 | Mantis Shrimp             | <i>Gonodactylus platysoma</i>       | CRE-Crustaceans |
| 62205 | Mantis Shrimp             | <i>Gonodactylus smithii</i>         | CRE-Crustaceans |
| 69860 | Shore Crabs               | Grapsidae                           | CRE-Crustaceans |
| 69861 | Shore Crab                | <i>Grapsus albolineatus</i>         | CRE-Crustaceans |
| 69862 | Shore Crab                | <i>Grapsus grapsus tenuicrustat</i> | CRE-Crustaceans |
| 69950 | Hapalocarcinids           | Hapalocarcinidae                    | CRE-Crustaceans |
| 62550 | Mantis Shrimp             | Harposquillidae                     | CRE-Crustaceans |
| 62300 | Mantis Shrimp             | Hemisquillidae                      | CRE-Crustaceans |
| 67104 | Deepwater Shrimps         | <i>Heteropenaeus</i> sp             | CRE-Crustaceans |
| 67210 | Hump-Backed Shrimp        | Hippolytidae                        | CRE-Crustaceans |
| 69100 | Homolids                  | Homolidae                           | CRE-Crustaceans |

|       |                     |                                    |                 |
|-------|---------------------|------------------------------------|-----------------|
| 67853 | Soft Lobster        | <i>Hoplometopus holthuisi</i>      | CRE-Crustaceans |
| 67223 | Harlequin Shrimp    | <i>Hymenocera picta</i>            | CRE-Crustaceans |
| 64810 | Hyperid Amphipods   | <i>Hyperiidae</i>                  | CRE-Crustaceans |
| 67921 | Slipper Lobster     | <i>Ibacus sp</i>                   | CRE-Crustaceans |
| 69000 | True Crabs          | <i>Io Brachyura</i>                | CRE-Crustaceans |
| 67931 | Long-Handed Lobster | <i>Justitia longimanus</i>         | CRE-Crustaceans |
| 67211 | Hump-Backed Shrimp  | <i>Koror misticius</i>             | CRE-Crustaceans |
| 69302 | Elbow Crab          | <i>Lambrus longispinis</i>         | CRE-Crustaceans |
| 67111 | Palaemonid Shrimp   | <i>Leander plumosus</i>            | CRE-Crustaceans |
| 68300 | Lithodids           | <i>Lithodidae</i>                  | CRE-Crustaceans |
| 69421 | Swimming Crab       | <i>Lupocyclus grimquedentatus</i>  | CRE-Crustaceans |
| 64830 | Lycaeids            | <i>Lycaeidae</i>                   | CRE-Crustaceans |
| 69151 | 3-Toothed Frog Crab | <i>Lyreidus tridentatus</i>        | CRE-Crustaceans |
| 62800 | Mantis Shrimp       | <i>Lysiosquillidae</i>             | CRE-Crustaceans |
| 60100 | Barnacles           | <i>Lythoglyptidae</i>              | CRE-Crustaceans |
| 69901 | Telescope-Eye Crab  | <i>Macrophthalmus telescopicus</i> | CRE-Crustaceans |
| 69250 | Spider Crabs        | <i>Majidae</i>                     | CRE-Crustaceans |
| 67101 | Penaeid Prawn       | <i>Metapenaeopsis sp.1</i>         | CRE-Crustaceans |
| 67102 | Penaeid Prawn       | <i>Metapenaeopsis sp.2</i>         | CRE-Crustaceans |
| 67103 | Penaeid Prawn       | <i>Metapenaeopsis sp.3</i>         | CRE-Crustaceans |
| 69205 | Box Crab            | <i>Mursia spinimanus</i>           | CRE-Crustaceans |
| 62900 | Mantis Shrimp       | <i>Nannosquillidae</i>             | CRE-Crustaceans |
| 67850 | Soft Lobsters       | <i>Nephropidae</i>                 | CRE-Crustaceans |
| 69902 | Large Ghost Crab    | <i>Ocypode ceratophthalma</i>      | CRE-Crustaceans |
| 69903 | Ghost Crab          | <i>Ocypode cordimana</i>           | CRE-Crustaceans |
| 69904 | Ghost Crab          | <i>Ocypode saratum</i>             | CRE-Crustaceans |
| 69900 | Ocypodids           | <i>Ocypodidae</i>                  | CRE-Crustaceans |
| 62350 | Mantis Shrimp       | <i>Odontodactylidae</i>            | CRE-Crustaceans |
| 62351 | Mantis Shrimp       | <i>Odontodactylus brevirostris</i> | CRE-Crustaceans |
| 62352 | Mantis Shrimp       | <i>Odontodactylus scyallarus</i>   | CRE-Crustaceans |
| 62701 | Mantis Shrimp       | <i>Oratosquilla oratoria</i>       | CRE-Crustaceans |
| 62700 | Mantis Shrimp       | <i>Oratosquillidae</i>             | CRE-Crustaceans |
| 68400 | Soldier Hermit Crab | <i>Paguridae</i>                   | CRE-Crustaceans |
| 68401 | Coral Hermit Crab   | <i>Paguritta gracilipes</i>        | CRE-Crustaceans |
| 68402 | Coral Hermit Crab   | <i>Paguritta harmsi</i>            | CRE-Crustaceans |
| 67110 | Palaemonid Shrimp   | <i>Palaemonidae</i>                | CRE-Crustaceans |
| 67917 | Mole Lobster        | <i>Palinurellus wieneckii</i>      | CRE-Crustaceans |
| 67918 | Painted Crayfish    | <i>Panulirus albiflagellum</i>     | CRE-Crustaceans |



|       |                         |  |                 |
|-------|-------------------------|--|-----------------|
| 67911 | Painted Crayfish        | <i>Panulirus homarus</i>                   | CRE-Crustaceans |
| 67912 | Painted Crayfish        | <i>Panulirus longipes</i>                  | CRE-Crustaceans |
| 67914 | Painted Crayfish        | <i>Panulirus ornatus</i>                   | CRE-Crustaceans |
| 67910 | Painted Crayfish        | <i>Panulirus sp</i>                        | CRE-Crustaceans |
| 67916 | Painted Crayfish        | <i>Panulirus versicolor</i>                | CRE-Crustaceans |
| 69300 | Elbow Crabs             | <i>Parthenopidae</i>                       | CRE-Crustaceans |
| 67100 | Panaeid Prawns          | <i>Penaeidae</i>                           | CRE-Crustaceans |
| 67106 | Penaeid Prawn           | <i>Penaeus latisulcatus</i>                | CRE-Crustaceans |
| 67105 | Penaeid Prawn           | <i>Penaeus monodon</i>                     | CRE-Crustaceans |
| 69864 | Flat Rock Crab          | <i>Percnon planissimum</i>                 | CRE-Crustaceans |
| 67122 | Commensal Shrimp        | <i>Periclimenes amboinensis</i>            | CRE-Crustaceans |
| 67123 | Commensal Shrimp        | <i>Periclimenes brevicarpalis</i>          | CRE-Crustaceans |
| 67124 | Commensal Shrimp        | <i>Periclimenes cf<br/>ceratophthalmus</i> | CRE-Crustaceans |
| 67125 | Commensal Shrimp        | <i>Periclimenes holthuisi</i>              | CRE-Crustaceans |
| 67126 | Commensal Shrimp        | <i>Periclimenes imperator</i>              | CRE-Crustaceans |
| 67127 | Commensal Shrimp        | <i>Periclimenes inornatus</i>              | CRE-Crustaceans |
| 67128 | Commensal Shrimp        | <i>Periclimenes kororensis</i>             | CRE-Crustaceans |
| 67129 | Commensal Shrimp        | <i>Periclimenes ornatus</i>                | CRE-Crustaceans |
| 67130 | Commensal Shrimp        | <i>Periclimenes psamathe</i>               | CRE-Crustaceans |
| 67131 | Commensal Shrimp        | <i>Periclimenes soror</i>                  | CRE-Crustaceans |
| 67132 | Commensal Shrimp        | <i>Periclimenes tenuipes</i>               | CRE-Crustaceans |
| 67133 | Commensal Shrimp        | <i>Periclimenes venustus</i>               | CRE-Crustaceans |
| 68601 | Porcelain Crab          | <i>Petrolisthes lamarkii</i>               | CRE-Crustaceans |
| 64820 | Phronimids              | <i>Phronimidae</i>                         | CRE-Crustaceans |
| 69863 | Shore Crab              | <i>Plagusia depressa<br/>tuberculata</i>   | CRE-Crustaceans |
| 64840 | Platyscelids            | <i>Platyscelidae</i>                       | CRE-Crustaceans |
| 67134 | Commensal Shrimp        | <i>Pliopotonia furtiva</i>                 | CRE-Crustaceans |
| 69461 | Long-Eyed Swimming Crab | <i>Podophthalmus vigil</i>                 | CRE-Crustaceans |
| 67135 | Commensal Shrimp        | <i>Pontonides uncigar</i>                  | CRE-Crustaceans |
| 67120 | Commensal Shrimp        | <i>Pontoniidae</i>                         | CRE-Crustaceans |
| 68600 | Porcellanid Crabs       | <i>Porcellanidae</i>                       | CRE-Crustaceans |
| 69400 | Swimming Crabs          | <i>Portunidae</i>                          | CRE-Crustaceans |
| 69432 | Blue Swimming Crab      | <i>Portunus pelagicus</i>                  | CRE-Crustaceans |
| 69431 | Swimming Crab           | <i>Portunus sanguinolentus</i>             | CRE-Crustaceans |
| 62400 | Mantis Shrimp           | <i>Protosquillidae</i>                     | CRE-Crustaceans |
| 62501 | Mantis Shrimp           | <i>Pseudosquilla ciliata</i>               | CRE-Crustaceans |
| 62500 | Mantis Shrimp           | <i>Pseudosquillidae</i>                    | CRE-Crustaceans |
| 67231 | Hingebeak Prawn         | <i>Rhynchocinetes hiatti</i>               | CRE-Crustaceans |

|       |                            |                                  |                 |
|-------|----------------------------|----------------------------------|-----------------|
| 67230 | Hinge-Beaked Prawns        | <i>Rhynchocinetidae</i>          | CRE-Crustaceans |
| 69471 | Mangrove Crab              | <i>Scylla serrata</i>            | CRE-Crustaceans |
| 67604 | Solenocerids               | <i>Solenoceridae</i>             | CRE-Crustaceans |
| 62600 | Mantis Shrimp              | <i>Squilla</i>                   | CRE-Crustaceans |
| 67136 | Commensal Shrimp           | <i>Stegopontonia commensalis</i> | CRE-Crustaceans |
| 67200 | Cleaner Shrimp             | <i>Stenopodidae</i>              | CRE-Crustaceans |
| 67201 | Banded Coral Shrimp        | <i>Stenopus hispidus</i>         | CRE-Crustaceans |
| 62000 | Mantis Shrimps             | <i>Stomatopoda</i>               | CRE-Crustaceans |
| 67503 | Snapping Shrimp            | <i>Synalpheus carinatus</i>      | CRE-Crustaceans |
| 60102 | Acorn Barnacle             | <i>Tetraclitella divisa</i>      | CRE-Crustaceans |
| 69481 | Swimming Crab              | <i>Thalamita crenata</i>         | CRE-Crustaceans |
| 67212 | Ambonian Shrimp            | <i>Thor amboinensis</i>          | CRE-Crustaceans |
| 69598 | Xanthid Crab               | <i>Unid Megalops</i>             | CRE-Crustaceans |
| 69499 | Portunid Crab              | <i>Unid sp.1</i>                 | CRE-Crustaceans |
| 69599 | Xanthid Crab               | <i>Unid sp.1</i>                 | CRE-Crustaceans |
| 69498 | Portunid Crab              | <i>Unid sp.2</i>                 | CRE-Crustaceans |
| 69597 | Xanthid Crab               | <i>Unid sp.2</i>                 | CRE-Crustaceans |
| 67112 | Palaemonid Shrimp          | <i>Urocaridella antonbruunii</i> | CRE-Crustaceans |
| 69500 | Dark-Finger Coral Crabs    | <i>Xanthidae</i>                 | CRE-Crustaceans |
| 69870 | Urchin Crab                | <i>Zebrida adamsii</i>           | CRE-Crustaceans |
| 69552 | Shallow Reef Crab          | <i>Zosymus aeneus</i>            | CRE-Crustaceans |
| 24300 | Squirrel,Soldierfishes     | <i>Holocentridae</i>             | Holocentridae   |
| 24398 | Squirrelfishes             | <i>Holocentrinae</i>             | Holocentridae   |
| 24399 | Soldierfishes              | <i>Myripristinae</i>             | Holocentridae   |
| 24313 | Bronze Soldierfish         | <i>Myripristis adusta</i>        | Holocentridae   |
| 24314 | Brick Soilderfish          | <i>Myripristis amaena</i>        | Holocentridae   |
| 24331 | Doubletooth Soldierfish    | <i>Myripristis amaena</i>        | Holocentridae   |
| 24315 | Bigscale Soldierfish       | <i>Myripristis berndti</i>       | Holocentridae   |
| 24324 | Yellowfin Soldierfish      | <i>Myripristis chryseres</i>     | Holocentridae   |
| 24317 | Pearly Soldierfish         | <i>Myripristis kuntee</i>        | Holocentridae   |
| 24318 | Red Soldierfish            | <i>Myripristis murdjan</i>       | Holocentridae   |
| 24322 | Scarlet Soldierfish        | <i>Myripristis pralinia</i>      | Holocentridae   |
| 24319 | Violet Soldierfish         | <i>Myripristis violacea</i>      | Holocentridae   |
| 24320 | White-Tipped Soldierfish   | <i>Myripristis vittata</i>       | Holocentridae   |
| 24326 | White-Spot Soldierfish     | <i>Myripristis woodsi</i>        | Holocentridae   |
| 24309 | Clearfin Squirrelfish      | <i>Neoniphon argenteus</i>       | Holocentridae   |
| 24312 | Yellowstriped Squirrelfish | <i>Neoniphon aurolineatus</i>    | Holocentridae   |
| 24310 | Blackfin Squirrlefish      | <i>Neoniphon opercularis</i>     | Holocentridae   |
| 24311 | Bloodspot Squirrelfish     | <i>Neoniphon sammara</i>         | Holocentridae   |

|       |                           |                                    |               |
|-------|---------------------------|------------------------------------|---------------|
| 24340 | Deepwater Soldierfish     | <i>Ostichthys brachygnathus</i>    | Holocentridae |
| 24323 | Deepwater Soldierfish     | <i>Ostichthys kaianus</i>          | Holocentridae |
| 24321 | Cardinal Squirrelfish     | <i>Plectrypops lima</i>            | Holocentridae |
| 24301 | Tailspot Squirrelfish     | <i>Sargocentron caudimaculatum</i> | Holocentridae |
| 24332 | 3-Spot Squirrelfish       | <i>Sargocentron cornutum</i>       | Holocentridae |
| 24302 | Crown Squirrelfish        | <i>Sargocentron diadema</i>        | Holocentridae |
| 24330 | Spotfin Squirrelfish      | <i>Sargocentron dorsomaculatum</i> | Holocentridae |
| 24334 | Furcate Squirrelfish      | <i>Sargocentron furcatum</i>       | Holocentridae |
| 24327 | Samurai Squirrelfish      | <i>Sargocentron ittodai</i>        | Holocentridae |
| 24333 | Squirrelfish              | <i>Sargocentron lepros</i>         | Holocentridae |
| 24328 | Blackspot Squirrelfish    | <i>Sargocentron melanospilos</i>   | Holocentridae |
| 24304 | Finelined Squirrelfish    | <i>Sargocentron microstoma</i>     | Holocentridae |
| 24305 | Dark-Striped Squirrelfish | <i>Sargocentron praslin</i>        | Holocentridae |
| 24303 | Speckled Squirrelfish     | <i>Sargocentron punctatissimum</i> | Holocentridae |
| 24306 | Long-Jawed Squirrelfish   | <i>Sargocentron spiniferum</i>     | Holocentridae |
| 24307 | Blue-Lined Squirrelfish   | <i>Sargocentron tiere</i>          | Holocentridae |
| 24308 | Pink Squirrelfish         | <i>Sargocentron tieroides</i>      | Holocentridae |
| 24329 | Violet Squirrelfish       | <i>Sargocentron violaceum</i>      | Holocentridae |
| 92102 | Algae                     | <i>Enteromorpha clathrata</i>      | Algae         |
| 92200 | Algae                     | <i>Caulerpaceae</i>                | Algae         |
| 92217 | Algae                     | <i>Caulerpa racemosa</i>           | Algae         |
| 93602 | Algae                     | <i>Sargassum polycystum</i>        | Algae         |
| 93604 | Algae                     | <i>Turbinaria ornata</i>           | Algae         |
| 95000 | Algae                     | <i>Div Anthophyta</i>              | Algae         |
| 95003 | Algae                     | <i>Halodule uninervis</i>          | Algae         |
| 36201 | Chiseltooth Wrasse        | <i>Anampses caeruleopunctatus</i>  | Labridae      |
| 36297 | Geographic Wrasse         | <i>Anampses geographicus</i>       | Labridae      |
| 36268 | Wrasse                    | <i>Anampses melanurus</i>          | Labridae      |
| 36202 | Yellowtail Wrasse         | <i>Anampses meleagrides</i>        | Labridae      |
| 36203 | Yellowbreasted Wrasse     | <i>Anampses twisti</i>             | Labridae      |
| 36205 | Lyretail Hogfish          | <i>Bodianus anthioides</i>         | Labridae      |
| 36206 | Axilspot Hogfish          | <i>Bodianus axillaris</i>          | Labridae      |
| 36288 | 2-Spot Slender Hogfish    | <i>Bodianus bimaculatus</i>        | Labridae      |
| 36269 | Diana'S Hogfish           | <i>Bodianus diana</i>              | Labridae      |
| 36270 | Blackfin Hogfish          | <i>Bodianus loxozonus</i>          | Labridae      |
| 36271 | Mesothorax Hogfish        | <i>Bodianus mesothorax</i>         | Labridae      |

|       |                          |                                     |          |
|-------|--------------------------|-------------------------------------|----------|
| 36243 | Hogfish                  | <i>Bodianus tanyokidus</i>          | Labridae |
| 36209 | Floral Wrasse            | <i>Cheilinus chlorourus</i>         | Labridae |
| 36210 | Red-Breasted Wrasse      | <i>Cheilinus fasciatus</i>          | Labridae |
| 36211 | Snooty Wrasse            | <i>Cheilinus oxycephalus</i>        | Labridae |
| 36213 | Tripletail Wrasse        | <i>Cheilinus trilobatus</i>         | Labridae |
| 36216 | Cigar Wrasse             | <i>Cheilio inermis</i>              | Labridae |
| 36217 | Yel-Cheeked Tuskfish     | <i>Choerodon anchorago</i>          | Labridae |
| 36313 | Harlequin Tuskfish       | <i>Choerodon fasciatus</i>          | Labridae |
| 36305 | Wrasse                   | <i>Cirrhilabrus balteatus</i>       | Labridae |
| 36272 | Wrasse                   | <i>Cirrhilabrus cyanopleura</i>     | Labridae |
| 36273 | Exquisite Wrasse         | <i>Cirrhilabrus exquisitus</i>      | Labridae |
| 36306 | Johnson'S Wrasse         | <i>Cirrhilabrus johnsoni</i>        | Labridae |
| 36218 | Wrasse                   | <i>Cirrhilabrus katherinae</i>      | Labridae |
| 36274 | Yellowband Wrasse        | <i>Cirrhilabrus luteovittatus</i>   | Labridae |
| 36307 | Rhomboid Wrasse          | <i>Cirrhilabrus rhomboidalis</i>    | Labridae |
| 36309 | Red-Margined Wrasse      | <i>Cirrhilabrus rubrimarginatus</i> | Labridae |
| 36219 | Clown Coris              | <i>Coris aygula</i>                 | Labridae |
| 36275 | Dapple Coris             | <i>Coris batuensis</i>              | Labridae |
| 36314 | Pale-Barred Coris        | <i>Coris dorsomacula</i>            | Labridae |
| 36220 | Yellowtailed Coris       | <i>Coris gaimardi</i>               | Labridae |
| 36221 | Knife Razorfish          | <i>Cymolutes praetextatus</i>       | Labridae |
| 36291 | Finescale Razorfish      | <i>Cymolutes torquatus</i>          | Labridae |
| 36300 | Wandering Cleaner Wrasse | <i>Diproctacanthus xanthurus</i>    | Labridae |
| 36222 | Sling-Jawed Wrasse       | <i>Epibulus insidiator</i>          | Labridae |
| 36276 | Sling-Jawed Wrasse       | <i>Epibulus n sp</i>                | Labridae |
| 36223 | Bird Wrasse              | <i>Gomphosus varius</i>             | Labridae |
| 36224 | 2-Spotted Wrasse         | <i>Halichoeres biocellatus</i>      | Labridae |
| 36277 | Drab Wrasse              | <i>Halichoeres chloropterus</i>     | Labridae |
| 36278 | Canary Wrasse            | <i>Halichoeres chrysus</i>          | Labridae |
| 36318 | Wrasse                   | <i>Halichoeres dussumieri</i>       | Labridae |
| 36226 | Checkerboard Wrasse      | <i>Halichoeres hortulanus</i>       | Labridae |
| 36227 | Weedy Surge Wrasse       | <i>Halichoeres margaritaceus</i>    | Labridae |
| 36228 | Dusky Wrasse             | <i>Halichoeres marginatus</i>       | Labridae |
| 36279 | Pinstriped Wrasse        | <i>Halichoeres melanurus</i>        | Labridae |
| 36229 | Black-Ear Wrasse         | <i>Halichoeres melasmapomus</i>     | Labridae |
| 36311 | Ornate Wrasse            | <i>Halichoeres ornatissimus</i>     | Labridae |
| 36315 | Seagrass Wrasse          | <i>Halichoeres papilionaceus</i>    | Labridae |
| 36298 | Wrasse                   | <i>Halichoeres prosopeion</i>       | Labridae |
| 36304 | Wrasse                   | <i>Halichoeres purpurascens</i>     | Labridae |

|       |                           |                                      |          |
|-------|---------------------------|--------------------------------------|----------|
| 36280 | Richmond'S Wrasse         | <i>Halichoeres richmondi</i>         | Labridae |
| 36281 | Zigzag Wrasse             | <i>Halichoeres scapularis</i>        | Labridae |
| 36312 | Shwartz Wrasse            | <i>Halichoeres shwartzi</i>          | Labridae |
| 36282 | Wrasse                    | <i>Halichoeres sp</i>                | Labridae |
| 36230 | 3-Spot Wrasse             | <i>Halichoeres trimaculatus</i>      | Labridae |
| 36225 | Wrasse                    | <i>Halichoeres zeylonicus</i>        | Labridae |
| 36231 | Striped Clown Wrasse      | <i>Hemigymnus fasciatus</i>          | Labridae |
| 36232 | 1/2 & 1/2 Wrasse          | <i>Hemigymnus melapterus</i>         | Labridae |
| 36303 | Wrasse                    | <i>Hologymnosus annulatus</i>        | Labridae |
| 36233 | Ring Wrasse               | <i>Hologymnosus doliatus</i>         | Labridae |
| 36234 | Tubelip Wrasse            | <i>Labrichthys unilineatus</i>       | Labridae |
| 36200 | Wrasse                    | Labridae                             | Labridae |
| 36235 | Bicolor Cleaner Wrasse    | <i>Labroides bicolor</i>             | Labridae |
| 36266 | Bluestreak Cleaner Wrasse | <i>Labroides dimidiatus</i>          | Labridae |
| 36237 | Black-Spot Cleaner Wrasse | <i>Labroides pectoralis</i>          | Labridae |
| 36283 | Allen'S Wrasse            | <i>Labropsis alleni</i>              | Labridae |
| 36238 | Micronesian Wrasse        | <i>Labropsis micronesica</i>         | Labridae |
| 36239 | Wedge-Tailed Wrasse       | <i>Labropsis xanthonota</i>          | Labridae |
| 36240 | Leopard Wrasse            | <i>Macropharyngodon meleagris</i>    | Labridae |
| 36284 | Negros Wrasse             | <i>Macropharyngodon negrosensis</i>  | Labridae |
| 36241 | Seagrass Razorfish        | <i>Novaculichthys macrolepidotus</i> | Labridae |
| 36242 | Dragon Wrasse             | <i>Novaculichthys taeniourus</i>     | Labridae |
| 36207 | Arenatus Wrasse           | <i>Oxycheilinus arenatus</i>         | Labridae |
| 36264 | 2-Spot Wrasse             | <i>Oxycheilinus bimaculatus</i>      | Labridae |
| 36208 | Celebes Wrasse            | <i>Oxycheilinus celebecus</i>        | Labridae |
| 36263 | Bandcheek Wrasse          | <i>Oxycheilinus digrammus</i>        | Labridae |
| 36215 | Oriental Wrasse           | <i>Oxycheilinus orientalis</i>       | Labridae |
| 36212 | Ringtail Wrasse           | <i>Oxycheilinus unifasciatus</i>     | Labridae |
| 36292 | Wrasse                    | <i>Paracheilinus bellae</i>          | Labridae |
| 36293 | Wrasse                    | <i>Paracheilinus sp.</i>             | Labridae |
| 36265 | Wrasse                    | <i>Polylepion russelli</i>           | Labridae |
| 36294 | Wrasse                    | <i>Pseudocheilinops ataenia</i>      | Labridae |
| 36244 | Striated Wrasse           | <i>Pseudocheilinus evanidus</i>      | Labridae |
| 36245 | 6 Line Wrasse             | <i>Pseudocheilinus hexataenia</i>    | Labridae |
| 36246 | 8 Line Wrasse             | <i>Pseudocheilinus octotaenia</i>    | Labridae |
| 36285 | Line Wrasse               | <i>Pseudocheilinus sp</i>            | Labridae |
| 36247 | 4 Line Wrasse             | <i>Pseudocheilinus tetrataenia</i>   | Labridae |

|       |                        |                                     |                |
|-------|------------------------|-------------------------------------|----------------|
| 36316 | Rust-Banded Wrasse     | <i>Pseudocoris aurantiofasciata</i> | Labridae       |
| 36317 | Torpedo Wrasse         | <i>Pseudocoris heteroptera</i>      | Labridae       |
| 36286 | Yamashiro'S Wrasse     | <i>Pseudocoris yamashiroi</i>       | Labridae       |
| 36267 | Chiseltooth Wrasse     | <i>Pseudodax moluccanus</i>         | Labridae       |
| 36248 | Polynesian Wrasse      | <i>Pseudojuloides atavai</i>        | Labridae       |
| 36249 | Smalltail Wrasse       | <i>Pseudojuloides cerasinus</i>     | Labridae       |
| 36250 | Wrasse                 | <i>Pterogogus cryptus</i>           | Labridae       |
| 36296 | Wrasse                 | <i>Pterogogus guttatus</i>          | Labridae       |
| 36251 | Red-Shoulder Wrasse    | <i>Stethojulis bandanensis</i>      | Labridae       |
| 36252 | Wrasse                 | <i>Stethojulis strigiventor</i>     | Labridae       |
| 36299 | Wrasse                 | <i>Stethojulis trilineata</i>       | Labridae       |
| 36253 | 2 Tone Wrasse          | <i>Thalassoma amblycephalum</i>     | Labridae       |
| 36255 | 6 Bar Wrasse           | <i>Thalassoma hardwickii</i>        | Labridae       |
| 36262 | Jansen'S Wrasse        | <i>Thalassoma janseni</i>           | Labridae       |
| 36287 | Crescent Wrasse        | <i>Thalassoma lunare</i>            | Labridae       |
| 36256 | Sunset Wrasse          | <i>Thalassoma lutescens</i>         | Labridae       |
| 36257 | Surge Wrasse           | <i>Thalassoma purpureum</i>         | Labridae       |
| 36258 | 5-Stripe Surge Wrasse  | <i>Thalassoma quinquevittatum</i>   | Labridae       |
| 36254 | Xmas Wrasse            | <i>Thalassoma trilobatum</i>        | Labridae       |
| 36289 | Wh-Barred Pygmy Wrasse | <i>Wetmorella albofasciata</i>      | Labridae       |
| 36259 | Bl-Spot Pygmy Wrasse   | <i>Wetmorella nigropinnata</i>      | Labridae       |
| 36290 | Wrasse                 | <i>Xiphocheilus</i> sp.             | Labridae       |
| 36261 | Yblotch Razorfish      | <i>Xyrichtys aneitensis</i>         | Labridae       |
| 36301 | Celebe'S Razorfish     | <i>Xyrichtys celebecus</i>          | Labridae       |
| 36302 | Razorfish              | <i>Xyrichtys geisha</i>             | Labridae       |
| 36308 | Yellowpatch Razorfish  | <i>Xyrichtys melanopus</i>          | Labridae       |
| 36260 | Blue Razorfish         | <i>Xyrichtys pavo</i>               | Labridae       |
| 36401 | Bumphead parrotfish    | <i>Bolbometopon muricatum</i>       | Labridae       |
| 36214 | Napolean wrasse        | <i>Cheilius undulatus</i>           | Labridae       |
| 1101  | Carcharhinidae         | <i>Carcharhinus albimarginatus</i>  | Carcharhinidae |
| 1102  | Carcharhinidae         | <i>Carcharhinus amblyrhynchos</i>   | Carcharhinidae |
| 1104  | Carcharhinidae         | <i>Carcharhinus galapagensis</i>    | Carcharhinidae |
| 1106  | Carcharhinidae         | <i>Carcharhinus melanopterus</i>    | Carcharhinidae |
| 1201  | Hammerhead             | <i>Sphyrna lewini</i>               | Carcharhinidae |
| 1202  | Hammerhead             | <i>Sphyrna mokorran</i>             | Carcharhinidae |
| 1200  | Hammerhead             | Sphyrnidae                          | Carcharhinidae |
| 44518 | Starry Triggerfish     | <i>Abalistes stellatus</i>          | Other          |

|       |                       |                                     |       |
|-------|-----------------------|-------------------------------------|-------|
| 20701 | Barred Needlefish     | <i>Ablennes hians</i>               | Other |
| 35050 | Blackspot Sergeant    | <i>Abudefduf lorenzi</i>            | Other |
| 35051 | Yellowtail Sergeant   | <i>Abudefduf notatus</i>            | Other |
| 35001 | Banded Sergeant       | <i>Abudefduf septemfasciatus</i>    | Other |
| 35002 | Scis-Tail Sgt Major   | <i>Abudefduf sexfasciatus</i>       | Other |
| 35003 | Black Spot Sergeant   | <i>Abudefduf sordidus</i>           | Other |
| 35004 | Sergeant-Major        | <i>Abudefduf vaigiensis</i>         | Other |
| 29150 | Spiney Basslets       | Acanthoclinidae                     | Other |
| 29151 | Hiatt'S Basslet       | <i>Acanthoplesiops hiatti</i>       | Other |
| 40537 | Goby                  | <i>Acentrogobius bonti</i>          | Other |
| 44566 | Seagrass Filefish     | <i>Acreichthys tomentosus</i>       | Other |
| 25601 | Shrimpfish            | <i>Aeoliscus strigatus</i>          | Other |
| 2201  | Spotted Eagle Ray     | <i>Aetobatis narinari</i>           | Other |
| 2202  | Eagle Ray             | <i>Aetomyleaus maculatus</i>        | Other |
| 4801  | Indo-Pacific Bonefish | <i>Albula glossodonta</i>           | Other |
| 4802  | Bonefish              | <i>Albula neoguinaica</i>           | Other |
| 4800  | Bonefish              | Albulidae                           | Other |
| 17100 | Lancetfishes          | Alepisauidae                        | Other |
| 17101 | Lancetfish            | <i>Alepisaurus ferox</i>            | Other |
| 40711 | Dorothea'S Wriggler   | <i>Allomicrodesmis dorotheae</i>    | Other |
| 39202 | Blenny                | <i>Alticus arnoldorum</i>           | Other |
| 44558 | Unicorn Filefish      | <i>Aluterus monoceros</i>           | Other |
| 44551 | Filefish              | <i>Aluterus scriptus</i>            | Other |
| 44552 | Filefish              | <i>Amanses scopas</i>               | Other |
| 28700 | Glass Perch           | Ambassidae                          | Other |
| 28701 | Glassie               | <i>Ambassis buruensis</i>           | Other |
| 28702 | Glassie               | <i>Ambassis interrupta</i>          | Other |
| 35201 | 2-Spot Hawkfish       | <i>Amblycirrhitus bimacula</i>      | Other |
| 40501 | Goby                  | <i>Amblyeleotris faciata</i>        | Other |
| 40502 | Goby                  | <i>Amblyeleotris fontaseni</i>      | Other |
| 40503 | Goby                  | <i>Amblyeleotris guttata</i>        | Other |
| 40506 | Goby                  | <i>Amblyeleotris randalli</i>       | Other |
| 40505 | Brown-Barred Goby     | <i>Amblyeleotris steinitzi</i>      | Other |
| 40507 | Bluespotted Goby      | <i>Amblyeleotris wheeleri</i>       | Other |
| 4306  | Blue Pilchard         | <i>Amblygaster clupeoides</i>       | Other |
| 4307  | Spotted Pilchard      | <i>Amblygaster sirm</i>             | Other |
| 35005 | Damsel                | <i>Amblygliphidodon aureus</i>      | Other |
| 35006 | Staghorn Damsel       | <i>Amblygliphidodon curacao</i>     | Other |
| 35052 | White-Belly Damsel    | <i>Amblygliphidodon leucogaster</i> | Other |

|       |                         |                                     |       |
|-------|-------------------------|-------------------------------------|-------|
| 35053 | Ternate Damsel          | <i>Amblygliphidodon ternatensis</i> | Other |
| 40523 | Goby                    | <i>Amblygobius decussatus</i>       | Other |
| 40524 | Goby                    | <i>Amblygobius hectori</i>          | Other |
| 40670 |                         | <i>Amblygobius linki</i>            | Other |
| 40525 | Goby                    | <i>Amblygobius nocturnus</i>        | Other |
| 40526 | Goby                    | <i>Amblygobius phalaena</i>         | Other |
| 40527 | Goby                    | <i>Amblygobius rainfordi</i>        | Other |
| 40662 | Goby                    | <i>Amblygobius sp</i>               | Other |
| 44816 | Evileye Puffer          | <i>Amblyrhynchotus honckenii</i>    | Other |
| 40504 | Prawn Goby              | <i>Amlbyeleotris periophthalma</i>  | Other |
| 35007 | Org-Fin Anemonefish     | <i>Amphiprion chrysopterus</i>      | Other |
| 35008 | Clark'S Anemonefish     | <i>Amphiprion clarkii</i>           | Other |
| 35095 | Tomato Anemonefish      | <i>Amphiprion frenatus</i>          | Other |
| 35009 | Dusky Anemonefish       | <i>Amphiprion melanopus</i>         | Other |
| 35096 | False Clown Anemonefish | <i>Amphiprion ocellaris</i>         | Other |
| 35010 | Pink Anemofish          | <i>Amphiprion peridaeraion</i>      | Other |
| 35097 | 3-Banded Anemonefish    | <i>Amphiprion tricinctus</i>        | Other |
| 43507 | Dragonet                | <i>Anaora tentaculata</i>           | Other |
| 5601  | Allardice'S Moray       | <i>Anarchias allardicei</i>         | Other |
| 5646  | Canton Island Moray     | <i>Anarchias cantonensis</i>        | Other |
| 5602  | Seychelles Moray        | <i>Anarchias seychellensis</i>      | Other |
| 4901  | Freshwater Eel          | <i>Anguilla bicolor</i>             | Other |
| 4902  | Freshwater Eel          | <i>Anguilla marmorata</i>           | Other |
| 4900  | Freshwater Eel          | Anguillidae                         | Other |
| 24250 | Flashlightfish          | Anomalopidae                        | Other |
| 24251 | Flashlightfish          | <i>Anomalops katoptron</i>          | Other |
| 19200 | Anglerfish              | Antenariidae                        | Other |
| 19201 | Pigmy Frogfish          | <i>Antennarius analis</i>           | Other |
| 19202 | Frogfish                | <i>Antennarius biocellatus</i>      | Other |
| 19203 | Freckled Frogfish       | <i>Antennarius coccineus</i>        | Other |
| 19204 | Giant Frogfish          | <i>Antennarius commersonii</i>      | Other |
| 19205 | Bandtail Frogfish       | <i>Antennarius dorehensis</i>       | Other |
| 19206 | Sargassumfish           | <i>Antennarius maculatus</i>        | Other |
| 19207 | Spotfin Frogfish        | <i>Antennarius nummifer</i>         | Other |
| 19208 | Painted Frogfish        | <i>Antennarius pictus</i>           | Other |
| 19209 | Randall'S Frogfish      | <i>Antennarius randalli</i>         | Other |
| 19210 | Spiney-Tufted Frogfish  | <i>Antennarius rosaceus</i>         | Other |
| 19211 | Bandfin Frogfish        | <i>Antennatus tuberosus</i>         | Other |



|       |                            |                                |       |
|-------|----------------------------|--------------------------------|-------|
| 25201 | Boarfish                   | <i>Antigonia malayana</i>      | Other |
| 26460 | Velvetfishes               | <i>Aploactinidae</i>           | Other |
| 30435 | Cardinalfish               | <i>Apogon amboinensis</i>      | Other |
| 30401 | Broad-Striped Cardinalfish | <i>Apogon angustatus</i>       | Other |
| 30402 | Bigeye Cardinalfish        | <i>Apogon bandanensis</i>      | Other |
| 30403 | Cryptic Cardinalfish       | <i>Apogon coccineus</i>        | Other |
| 30436 | Ohcre-Striped Cardinalfish | <i>Apogon compressus</i>       | Other |
| 30437 | Redspot Cardinalfish       | <i>Apogon dispar</i>           | Other |
| 30438 | Longspine Cardinalfish     | <i>Apogon doryssa</i>          | Other |
| 30455 | Elliot'S Cardinalfish      | <i>Apogon ellioiti</i>         | Other |
| 30462 | Cardinalfish               | <i>Apogon eremeia</i>          | Other |
| 30439 | Evermann'S Cardinalfish    | <i>Apogon evermanni</i>        | Other |
| 30404 | Eyeshadow Cardinalfish     | <i>Apogon exostigma</i>        | Other |
| 30405 | Bridled Cardinalfish       | <i>Apogon fraenatus</i>        | Other |
| 30441 | Cardinalfish               | <i>Apogon fragilis</i>         | Other |
| 30440 | Gilbert'S Cardinalfish     | <i>Apogon gilberti</i>         | Other |
| 30406 | Guam Cardinalfish          | <i>Apogon guamensis</i>        | Other |
| 30468 |                            | <i>Apogon hartzfeldii</i>      | Other |
| 30407 | Iridescent Cardinalfish    | <i>Apogon kallopterus</i>      | Other |
| 30408 | Inshore Cardinalfish       | <i>Apogon lateralis</i>        | Other |
| 30409 | Bluestreak Cardinalfish    | <i>Apogon leptacanthus</i>     | Other |
| 30457 | Black Cardinalfish         | <i>Apogon melas</i>            | Other |
| 30463 | Cardinalfish               | <i>Apogon nigripinnis</i>      | Other |
| 30412 | Black-Striped Cardinalfish | <i>Apogon nigrofasciatus</i>   | Other |
| 30464 | Cardinalfish               | <i>Apogon notatus</i>          | Other |
| 30413 | 7-Lined Cardinalfish       | <i>Apogon novemfasciatus</i>   | Other |
| 30442 | Pearly Cardinalfish        | <i>Apogon perlitus</i>         | Other |
| 30465 | Cardinalfish               | <i>Apogon rhodopterus</i>      | Other |
| 30443 | Sangi Cardinalfish         | <i>Apogon sangiensis</i>       | Other |
| 30415 | Gray Cardinalfish          | <i>Apogon savayensis</i>       | Other |
| 30456 | Seale'S Cardinalfish       | <i>Apogon sealei</i>           | Other |
| 30417 | Cardinalfish               | <i>Apogon sp</i>               | Other |
| 30414 | Bandfin Cardinalfish       | <i>Apogon taeniophorus</i>     | Other |
| 30410 | Bandfin Cardinalfish       | <i>Apogon taeniopterus</i>     | Other |
| 30416 | 3-Spot Cardinalfish        | <i>Apogon trimaculatus</i>     | Other |
| 30418 | Ocellated Cardinalfish     | <i>Apogonichthys ocellatus</i> | Other |
| 30444 | Perdix Cardinalfish        | <i>Apogonichthys perdix</i>    | Other |
| 30400 | Cardinalfishes             | Apogonidae                     | Other |
| 34377 | Angelfish                  | <i>Apolemichthys griffisi</i>  | Other |

|       |                           |                                       |       |
|-------|---------------------------|---------------------------------------|-------|
| 34351 | Flagfin Angelfish         | <i>Apolemichthys trimaculatus</i>     | Other |
| 34376 | Angelfish                 | <i>Apolemichthys xanthopunctatus</i>  | Other |
| 29201 | 2-Lined Soapfish          | <i>Aporops bilinearis</i>             | Other |
| 6619  | Snake Eel                 | <i>Apterichtus klazingai</i>          | Other |
| 30419 | Twinspot Cardinalfish     | <i>Archamia biguttata</i>             | Other |
| 30420 | Orange-Lined Cardinalfish | <i>Archamia fucata</i>                | Other |
| 30445 | Blackbelted Cardinalfish  | <i>Archamia zosterophora</i>          | Other |
| 6206  | Scheele'S Conger          | <i>Ariosoma scheelei</i>              | Other |
| 43903 | Flounder                  | <i>Arnoglossus intermedius</i>        | Other |
| 44801 | Brown Puffer              | <i>Arothron hispidus</i>              | Other |
| 44802 | Puffer                    | <i>Arothron manilensis</i>            | Other |
| 44803 | Puffer                    | <i>Arothron mappa</i>                 | Other |
| 44804 | White-Spot Puffer         | <i>Arothron meleagris</i>             | Other |
| 44805 | Black-Spotted Puffer      | <i>Arothron nigropunctatus</i>        | Other |
| 44806 | Star Puffer               | <i>Arothron stellatus</i>             | Other |
| 44102 | Black Spotted Sole        | <i>Aseraggodes melanostictus</i>      | Other |
| 44103 | Smith'S Sole              | <i>Aseraggodes smithi</i>             | Other |
| 44104 | Whitaker'S Sole           | <i>Aseraggodes whitakeri</i>          | Other |
| 39257 | Lance Blenny              | <i>Aspidontus dussumieri</i>          | Other |
| 39203 | Cleaner Mimic             | <i>Aspidontus taeniatus</i>           | Other |
| 40539 |                           | <i>Asteropteryx semipunctatus</i>     | Other |
| 43905 | Intermediate Flounder     | <i>Asterorhombus intermedius</i>      | Other |
| 40538 | Goby                      | <i>Asterropteryx ensiferus</i>        | Other |
| 21800 | Silverside                | Atherinidae                           | Other |
| 21805 | Tropical Silverside       | <i>Atherinomorus duodecimalis</i>     | Other |
| 21806 | Striped Silverside        | <i>Atherinomorus endrachtensis</i>    | Other |
| 21803 | Silverside                | <i>Atherinomorus lacunosus</i>        | Other |
| 21804 | Hardyhead Silverside      | <i>Atherinomorus lacunosus</i>        | Other |
| 21801 | Bearded Silverside        | <i>Atherion elymus</i>                | Other |
| 39240 | Blenny                    | <i>Atrosalariaus fuscus holomelas</i> | Other |
| 25300 | Trumpetfish               | Aulostomidae                          | Other |
| 25301 | Trumpetfish               | <i>Aulostomus chinensis</i>           | Other |
| 40540 | Goby                      | <i>Austrolethops wardi</i>            | Other |
| 40541 | Goby                      | <i>Awaous grammepomus</i>             | Other |
| 40542 | Goby                      | <i>Awaous guamensis</i>               | Other |
| 44501 | Undulate Triggerfish      | <i>Balistapus undulatus</i>           | Other |

|       |                            |                                  |       |
|-------|----------------------------|----------------------------------|-------|
| 44500 | Triggerfishes              | Balistidae                       | Other |
| 44502 | Clown Triggerfish          | <i>Balistoides conspicillum</i>  | Other |
| 44503 | Titan Triggerfish          | <i>Balistoides viridescens</i>   | Other |
| 40543 | Goby                       | <i>Bathygobius cocosensis</i>    | Other |
| 40544 | Goby                       | <i>Bathygobius cotticeps</i>     | Other |
| 40545 | Goby                       | <i>Bathygobius fuscus</i>        | Other |
| 20700 | Needlefish                 | Belonidae                        | Other |
| 29001 | Soapfish                   | <i>Belonoperca chaubanaudi</i>   | Other |
| 24200 | Lantern-Eye Fish           | Berycidae                        | Other |
| 24201 | Flashlightfish             | <i>Beryx decadactylus</i>        | Other |
| 25818 | Pipefish                   | <i>Bhanotia nuda</i>             | Other |
| 6205  | Conger Eel                 | <i>Blachea xenobranchialis</i>   | Other |
| 39218 | Blenny                     | <i>Blenniella cyanostigma</i>    | Other |
| 39222 | Blenny                     | <i>Blenniella gibbifrons</i>     | Other |
| 39239 |                            | <i>Blenniella paula</i>          | Other |
| 39221 | Blenny                     | <i>Blenniella periophthalmus</i> | Other |
| 39200 | Blennies                   | Blenniidae                       | Other |
| 43900 | Flounders                  | Bothidae                         | Other |
| 43901 | Peacock Flounder           | <i>Bothus mancus</i>             | Other |
| 43902 | Leopard Flounder           | <i>Bothus pantherinus</i>        | Other |
| 44559 | Taylor'S Inflater Filefish | <i>Brachaluteres taylori</i>     | Other |
| 6601  | Snake Eel                  | <i>Brachysomophis sauropsis</i>  | Other |
| 18201 | Codlet                     | <i>Bregmaceros nectabanus</i>    | Other |
| 18200 | Codlets                    | <i>Bregmacerotidae</i>           | Other |
| 18651 | Free-Tailed Brotula        | <i>Brosomphyciops pautzkei</i>   | Other |
| 18601 | Reef Cusk Eel              | <i>Brotula multibarbata</i>      | Other |
| 18602 | Townsend'S Cusk Eel        | <i>Brotula townsendi</i>         | Other |
| 40546 | Goby                       | <i>Bryaninops amplus</i>         | Other |
| 40547 | Goby                       | <i>Bryaninops erythrops</i>      | Other |
| 40548 | Goby                       | <i>Bryaninops natans</i>         | Other |
| 40549 | Goby                       | <i>Bryaninops ridens</i>         | Other |
| 40550 | Goby                       | <i>Bryaninops youngei</i>        | Other |
| 25819 | Pipefish                   | <i>Bulbonaricus brauni</i>       | Other |
| 40402 | Gudgeon                    | <i>Butis amboinensis</i>         | Other |
| 18650 | Livebearing Brotulas       | Bythitidae                       | Other |
| 40551 | Goby                       | <i>Cabillus tongarevae</i>       | Other |
| 6602  | Snake Eel                  | <i>Caecula polyophthalma</i>     | Other |
| 32351 | Scissor-Tailed Fusilier    | <i>Caesio caeruleaurea</i>       | Other |
| 32355 | Fusilier                   | <i>Caesio cuning</i>             | Other |

|       |                        |                                   |       |
|-------|------------------------|-----------------------------------|-------|
| 32356 | Lunar Fusilier         | <i>Caesio lunaris</i>             | Other |
| 32352 | Yellowback Caesio      | <i>Caesio teres</i>               | Other |
| 32350 | Fusilier               | <i>Caesionidae</i>                | Other |
| 29050 | Goldies                | <i>Callanthiidae</i>              | Other |
| 6603  | Snake Eel              | <i>Callechelys marmorata</i>      | Other |
| 6604  | Snake Eel              | <i>Callechelys melanotaenia</i>   | Other |
| 43500 | Dragonets              | <i>Callionymidae</i>              | Other |
| 43508 | Delicate Dragonet      | <i>Callionymus delicatulus</i>    | Other |
| 43501 | Mangrove Dragonet      | <i>Callionymus enneactis</i>      | Other |
| 43502 | Simple-Spined Dragonet | <i>Callionymus simplicicornis</i> | Other |
| 40559 | Goby                   | <i>Callogobius sp</i>             | Other |
| 40552 | Goby                   | <i>Callogobius bauchotae</i>      | Other |
| 40553 | Goby                   | <i>Callogobius centrolepis</i>    | Other |
| 40554 | Goby                   | <i>Callogobius hasselti</i>       | Other |
| 40555 | Goby                   | <i>Callogobius maculipinnis</i>   | Other |
| 40556 | Goby                   | <i>Callogobius okinawae</i>       | Other |
| 40557 | Goby                   | <i>Callogobius plumatus</i>       | Other |
| 40558 | Goby                   | <i>Callogobius sclateri</i>       | Other |
| 29401 | Longfin                | <i>Callopleysiops altivelis</i>   | Other |
| 40403 | Sleeper                | <i>Calumia godeffroyi</i>         | Other |
| 44553 | Gray Leatherjacket     | <i>Cantherhines dumerilii</i>     | Other |
| 44565 | Specktaled Filefish    | <i>Cantherhines fronticinctus</i> | Other |
| 44554 | Honeycomb Filefish     | <i>Cantherhines pardalis</i>      | Other |
| 44504 | Rough Triggerfish      | <i>Canthidermis maculatus</i>     | Other |
| 44807 | Puffer                 | <i>Canthigaster amboinensis</i>   | Other |
| 44808 | Puffer                 | <i>Canthigaster bennetti</i>      | Other |
| 44815 | Puffer                 | <i>Canthigaster compressa</i>     | Other |
| 44809 | Sharp Back Puffer      | <i>Canthigaster coronata</i>      | Other |
| 44810 | Puffer                 | <i>Canthigaster epilampra</i>     | Other |
| 44811 | Puffer                 | <i>Canthigaster janthinoptera</i> | Other |
| 44812 | Puffer                 | <i>Canthigaster leoparda</i>      | Other |
| 44819 | Circle-Barred Toby     | <i>Canthigaster ocellicineta</i>  | Other |
| 44820 | Papuan Toby            | <i>Canthigaster papua</i>         | Other |
| 44813 | Sharpnose Puffer       | <i>Canthigaster solandri</i>      | Other |
| 44814 | Saddle Shpns Puffer    | <i>Canthigaster valentini</i>     | Other |
| 25200 | Boarfishes             | <i>Caproidae</i>                  | Other |
| 26700 | Coral Crouchers        | <i>Caracanthidae</i>              | Other |
| 26701 | Velvetfish             | <i>Caracanthus maculatus</i>      | Other |
| 26702 | Velvetfish             | <i>Caracanthus unipinna</i>       | Other |

|       |                            |                                  |       |
|-------|----------------------------|----------------------------------|-------|
| 18700 | Pearlfish                  | <i>Carapodidae</i>               | Other |
| 18702 | Pearlfish                  | <i>Carapus mourlani</i>          | Other |
| 1109  | Blackfin Shark             | <i>Carcharhinus limbatus</i>     | Other |
| 902   | Great White Shark          | <i>Carcharodon carcharius</i>    | Other |
| 25600 | Shrimpfishes               | <i>Centriscidae</i>              | Other |
| 34379 | Golden Angelfish           | <i>Centropyge aurantia</i>       | Other |
| 34352 | Bicolor Angelfish          | <i>Centropyge bicolor</i>        | Other |
| 34353 | Dusky Angelfish            | <i>Centropyge bispinosus</i>     | Other |
| 34354 | Colin'S Angelfish          | <i>Centropyge colini</i>         | Other |
| 34367 | White-Tail Angelfish       | <i>Centropyge flavicauda</i>     | Other |
| 34355 | Lemonpeel Angelfish        | <i>Centropyge flavissimus</i>    | Other |
| 34356 | Herald'S Angelfish         | <i>Centropyge heraldi</i>        | Other |
| 34357 | Flame Angelfish            | <i>Centropyge loriculus</i>      | Other |
| 34368 | Multicolor Angelfish       | <i>Centropyge multicolor</i>     | Other |
| 34358 | Multibarred Angelfish      | <i>Centropyge multifasciatus</i> | Other |
| 34359 | Black-Spot Angelfish       | <i>Centropyge nigriocellus</i>   | Other |
| 34378 | Midnight Angelfish         | <i>Centropyge nox</i>            | Other |
| 34360 | Shepard'S Angelfish        | <i>Centropyge shepardi</i>       | Other |
| 34369 | Keyhole Angelfish          | <i>Centropyge tibicen</i>        | Other |
| 34361 | Pearlscale Angelfish       | <i>Centropyge vrolicki</i>       | Other |
| 28959 | Grouper                    | <i>Cephalopholis cyanostigma</i> | Other |
| 39008 | Triplefin                  | <i>Ceratobregma helenae</i>      | Other |
| 34301 | Threadfin Butterflyfish    | <i>Chaetodon auriga</i>          | Other |
| 34330 | E Triangular Butterflyfish | <i>Chaetodon barronessa</i>      | Other |
| 34302 | Bennetts Butterflyfish     | <i>Chaetodon bennetti</i>        | Other |
| 34331 | Burgess' Butterflyfish     | <i>Chaetodon burgessi</i>        | Other |
| 34303 | Speckled Butterflyfish     | <i>Chaetodon citrinellus</i>     | Other |
| 34304 | Saddleback Butterflyfish   | <i>Chaetodon ephippium</i>       | Other |
| 34305 | Ylw-Crn Butterflyfish      | <i>Chaetodon flavocoronatus</i>  | Other |
| 34306 | Kleins Butterflyfish       | <i>Chaetodon kleinii</i>         | Other |
| 34307 | Lined Butterflyfish        | <i>Chaetodon lineolatus</i>      | Other |
| 34308 | Racoon Butterflyfish       | <i>Chaetodon lunula</i>          | Other |
| 34316 | Redfinned Butterflyfish    | <i>Chaetodon lunulatus</i>       | Other |
| 34309 | Black-Back Butterflyfish   | <i>Chaetodon melannotus</i>      | Other |
| 34310 | Mertens Butterflyfish      | <i>Chaetodon mertensii</i>       | Other |
| 34332 | Meyer'S Butterflyfish      | <i>Chaetodon meyeri</i>          | Other |
| 34311 | Butterflyfish              | <i>Chaetodon modestus</i>        | Other |
| 34333 | Spot-Tail Butterflyfish    | <i>Chaetodon ocellicaudus</i>    | Other |
| 34334 | 8-Banded Butterflyfish     | <i>Chaetodon octofasciatus</i>   | Other |

|       |                           |                                      |       |
|-------|---------------------------|--------------------------------------|-------|
| 34312 | Ornate Butterflyfish      | <i>Chaetodon ornatissimus</i>        | Other |
| 34335 | Spot-Nape Butterflyfish   | <i>Chaetodon oxycephalus</i>         | Other |
| 34313 | Spotbnded Butterflyfish   | <i>Chaetodon punctatofasciatus</i>   | Other |
| 34314 | 4-Spotted Butterflyfish   | <i>Chaetodon quadrimaculatus</i>     | Other |
| 34336 | Latticed Butterflyfish    | <i>Chaetodon rafflesii</i>           | Other |
| 34315 | Retculted Butterflyfish   | <i>Chaetodon reticulatus</i>         | Other |
| 34337 | Dotted Butterflyfish      | <i>Chaetodon semeion</i>             | Other |
| 34338 | Oval-Spot Butterflyfish   | <i>Chaetodon speculum</i>            | Other |
| 34340 | Tinker'S Butterflyfish    | <i>Chaetodon tinkeri</i>             | Other |
| 34329 | Chevron Butterflyfish     | <i>Chaetodon trifascialis</i>        | Other |
| 34317 | Pac Dblsddl Butterflyfish | <i>Chaetodon ulietensis</i>          | Other |
| 34318 | Teardrop Butterflyfish    | <i>Chaetodon unimaculatus</i>        | Other |
| 34319 | Vagabond Butterflyfish    | <i>Chaetodon vagabundus</i>          | Other |
| 34300 | Butterflyfish             | <i>Chaetodontidae</i>                | Other |
| 34370 | Vermiculated Angelfish    | <i>Chaetodontoplus mesoleucus</i>    | Other |
| 37401 | Saddled Sandburrer        | <i>Chalixodytes tauensis</i>         | Other |
| 36701 | Gaper                     | <i>Champsodon vorax</i>              | Other |
| 36700 | Gapers                    | <i>Champsodontidae</i>               | Other |
| 9800  | Milkfish                  | <i>Chanidae</i>                      | Other |
| 5647  | Long-Jawed Moray          | <i>Channomuraena vittata</i>         | Other |
| 9801  | Milkfish                  | <i>Chanos chanos</i>                 | Other |
| 30458 | Lined Cardinalfish        | <i>Cheilodipterus artus</i>          | Other |
| 30466 | Intermediate Cardinalfish | <i>Cheilodipterus intermedius</i>    | Other |
| 30446 | Cardinalfish              | <i>Cheilodipterus isostigma</i>      | Other |
| 30422 | Lg-Toothed Cardinalfish   | <i>Cheilodipterus macrodon</i>       | Other |
| 30423 | 5-Lined Cardinalfish      | <i>Cheilodipterus quinquelineata</i> | Other |
| 30421 | Truncate Cardinalfish     | <i>Cheilodipterus singaporensis</i>  | Other |
| 20601 | Flying Fish               | <i>Cheilopogon spilonopterus</i>     | Other |
| 20602 | Flying Fish               | <i>Cheilopogon spilopterus</i>       | Other |
| 20603 | Flying Fish               | <i>Cheilopogon unicolor</i>          | Other |
| 35089 | Minstrel Fish             | <i>Cheiloprion labiatus</i>          | Other |
| 35907 | Ceram Mullet              | <i>Chelon macrolepis</i>             | Other |
| 5400  | False Moray Eel           | Chlopsidae                           | Other |
| 25802 | Pipefish                  | <i>Choeroichthys brachysoma</i>      | Other |
| 25801 | Pipefish                  | <i>Choeroichthys sculptus</i>        | Other |
| 37001 | Duckbill                  | <i>Chrionema squamiceps</i>          | Other |

|       |                      |                                    |       |
|-------|----------------------|------------------------------------|-------|
| 35011 | Midget Chromis       | <i>Chromis acares</i>              | Other |
| 35012 | Bronze Reef Chromis  | <i>Chromis agilis</i>              | Other |
| 35022 | Yel-Speckled Chromis | <i>Chromis alpha</i>               | Other |
| 35013 | Ambon Chromis        | <i>Chromis amboinensis</i>         | Other |
| 35014 | Yellow Chromis       | <i>Chromis analis</i>              | Other |
| 35015 | Black-Axil Chromis   | <i>Chromis atripectoralis</i>      | Other |
| 35054 | Dark-Fin Chromis     | <i>Chromis atripes</i>             | Other |
| 35059 | Blue-Axil Chromis    | <i>Chromis caudalis</i>            | Other |
| 35060 | Deep Reef Chromis    | <i>Chromis delta</i>               | Other |
| 35017 | Twin-Spot Chromis    | <i>Chromis elerae</i>              | Other |
| 35018 | Scaly Chromis        | <i>Chromis lepidolepis</i>         | Other |
| 35055 | Lined Chromis        | <i>Chromis lineata</i>             | Other |
| 35019 | Bicolor Chromis      | <i>Chromis margaritifer</i>        | Other |
| 35056 | Black-Bar Chromis    | <i>Chromis retrofasciata</i>       | Other |
| 35049 | Ternate Chromis      | <i>Chromis ternatensis</i>         | Other |
| 35020 | Vanderbilt'S Chromis | <i>Chromis vanderbilti</i>         | Other |
| 35016 | Blue-Green Chromis   | <i>Chromis viridis</i>             | Other |
| 35057 | Weber'S Chromis      | <i>Chromis weberi</i>              | Other |
| 35058 | Yel-Axil Chromis     | <i>Chromis xanthochir</i>          | Other |
| 35021 | Black Chromis        | <i>Chromis xanthura</i>            | Other |
| 35024 | 2-Spot Demoiselle    | <i>Chrysiptera biocellata</i>      | Other |
| 35027 | Surge Demoiselle     | <i>Chrysiptera brownriggii</i>     | Other |
| 35025 | Blue-Line Demoiselle | <i>Chrysiptera caeruleolineata</i> | Other |
| 35062 | Blue Devil           | <i>Chrysiptera cyanea</i>          | Other |
| 35026 | Gray Demoiselle      | <i>Chrysiptera glauca</i>          | Other |
| 35090 | Blue-Spot Demoiselle | <i>Chrysiptera oxycephala</i>      | Other |
| 35064 | King Demoiselle      | <i>Chrysiptera rex</i>             | Other |
| 35065 | Talbot'S Demoiselle  | <i>Chrysiptera talboti</i>         | Other |
| 35028 | Tracey'S Demoiselle  | <i>Chrysiptera traceyi</i>         | Other |
| 35091 | 1-Spot Demoiselle    | <i>Chrysiptera unimaculata</i>     | Other |
| 34610 | Peacock Bass         | <i>Cichla ocellaris</i>            | Other |
| 34600 | Cichlids             | Cichlidae                          | Other |
| 35211 | Threadfin Hawkfish   | <i>Cirrhichthys aprinus</i>        | Other |
| 35202 | Falco'S Hawkfish     | <i>Cirrhichthys falco</i>          | Other |
| 35203 | Pixy Hawkfish        | <i>Cirrhichthys oxycephalus</i>    | Other |
| 35200 | Hawkfish             | Cirrhitidae                        | Other |
| 35204 | Stocky Hawkfish      | <i>Cirrhitus pinnulatus</i>        | Other |
| 6620  | Fringelip Snake Eel  | <i>Cirricaecula johnsoni</i>       | Other |
| 39242 | Chestnut Blenny      | <i>Cirripectes castaneus</i>       | Other |

|       |                         |  |       |
|-------|-------------------------|--|-------|
| 39204 | Spotted Blenny          | <i>Cirripectes fuscoguttatus</i>       | Other |
| 39243 | Blenny                  | <i>Cirripectes perustus</i>            | Other |
| 39206 | Barred Blenny           | <i>Cirripectes polyzona</i>            | Other |
| 39205 | Squiggly Blenny         | <i>Cirripectes quagga</i>              | Other |
| 39244 | Red-Streaked Blenny     | <i>Cirripectes stigmaticus</i>         | Other |
| 39207 | Red-Speckled Blenny     | <i>Cirripectes variolosus</i>          | Other |
| 14802 | Air-Breath Catfish      | <i>Clarias batrachus</i>               | Other |
| 14801 | Air-Breath Catfish      | <i>Clarias macrocephalus</i>           | Other |
| 14800 | Air-Breath Catfish      | Clariidae                              | Other |
| 4300  | Herring,Sprat,Sardines  | Clupeidae                              | Other |
| 26461 | Velvetfish              | <i>Cocotropis larvatus</i>             | Other |
| 6201  | White Eel               | <i>Conger cinereus cinereus</i>        | Other |
| 6202  | Conger Eel              | <i>Conger oligoporus</i>               | Other |
| 6208  | Conger Eel              | <i>Conger sp</i>                       | Other |
| 6200  | White,Conger,Garden Eel | Congridae                              | Other |
| 30306 | Deepwater Glasseye      | <i>Cookeolus boops</i>                 | Other |
| 30304 | Bulleye                 | <i>Cookeolus japonicus</i>             | Other |
| 34339 | Orangebanded Coralfish  | <i>Coradion chrysozonus</i>            | Other |
| 40590 | Goby                    | <i>Coryphopterus signipinnis</i>       | Other |
| 25803 | Network Pipefish        | <i>Corythoichthys flavofasciatus</i>   | Other |
| 25820 | Pipefish                | <i>Corythoichthys haematopterus</i>    | Other |
| 25804 | Reef Pipefish           | <i>Corythoichthys intestinalis</i>     | Other |
| 25805 | Bl-Breasted Pipefish    | <i>Corythoichthys nigripectus</i>      | Other |
| 25821 | Ocellated Pipefish      | <i>Corythoichthys ocellatus</i>        | Other |
| 25822 | Many-Spotted Pipefish   | <i>Corythoichthys polynotatus</i>      | Other |
| 25823 | Guided Pipefish         | <i>Corythoichthys schultzi</i>         | Other |
| 25824 | Roughridge Pipefish     | <i>Cosmocampus banneri</i>             | Other |
| 25806 | D'Arros Pipefish        | <i>Cosmocampus darrosanus</i>          | Other |
| 25825 | Maxweber'S Pipefish     | <i>Cosmocampus maxweberi</i>           | Other |
| 37400 | Sand Burrowers          | Creedidae                              | Other |
| 35911 | Mullet                  | <i>Crenimugil heterochilos</i>         | Other |
| 40560 | Goby                    | <i>Cristagobius sp</i>                 | Other |
| 40508 | Goby                    | <i>Cryptocentroides insignis</i>       | Other |
| 40511 | Goby                    | <i>Cryptocentrus cauruleomaculatus</i> | Other |
| 40509 | Goby                    | <i>Cryptocentrus cinctus</i>           | Other |
| 40510 | Goby                    | <i>Cryptocentrus koumansi</i>          | Other |
| 40512 | Goby                    | <i>Cryptocentrus</i>                   | Other |



|       |                        |   |       |
|-------|------------------------|---|-------|
|       |                        | <i>leptocephalus</i>                      |       |
| 40514 | Goby                   | <i>Cryptocentrus sp.A</i>                 | Other |
| 40513 | Goby                   | <i>Cryptocentrus strigiliceps</i>         | Other |
| 40515 | Goby                   | <i>Ctenogobiops aurocingulus</i>          | Other |
| 40516 | Goby                   | <i>Ctenogobiops feroculus</i>             | Other |
| 40517 | Goby                   | <i>Ctenogobiops pomastictus</i>           | Other |
| 40518 | Long-Finned Prwn Goby  | <i>Ctenogobiops tangarorai</i>            | Other |
| 27304 | Flathead               | <i>Cymbacephalus beauforti</i>            | Other |
| 35212 | Swallowtail Hawkfish   | <i>Cyprinocirrhites polyactis</i>         | Other |
| 20604 | Flying Fish            | <i>Cypselurus angusticeps</i>             | Other |
| 20605 | Flying Fish            | <i>Cypselurus poecilopterus</i>           | Other |
| 20606 | Flying Fish            | <i>Cypselurus speculiger</i>              | Other |
| 28501 | Flying Gurnard         | <i>Dactyloptena orientalis</i>            | Other |
| 28502 | Flying Gurnard         | <i>Dactyloptena petersoni</i>             | Other |
| 28500 | Flying Gurnard         | <i>Dactylopteridae</i>                    | Other |
| 35029 | Humbug Dascyllus       | <i>Dascyllus aruanus</i>                  | Other |
| 35066 | Black-Tail Dascyllus   | <i>Dascyllus melanurus</i>                | Other |
| 35030 | Reticulated Dascyllus  | <i>Dascyllus reticulatus</i>              | Other |
| 35031 | 3-Spot Dascyllus       | <i>Dascyllus trimaculatus</i>             | Other |
| 2000  | Stingray               | Dasyatidae                                | Other |
| 2001  | Blue-Spotted Sting Ray | <i>Dasyatis kuhlii</i>                    | Other |
| 26401 | Scorpionfish           | <i>Dendrochirus biocellatus</i>           | Other |
| 26402 | Scorpionfish           | <i>Dendrochirus brachypterus</i>          | Other |
| 26427 | Zebra Lionfish         | <i>Dendrochirus zebra</i>                 | Other |
| 32701 | Slaty Sweetlips        | <i>Diagramma pictum</i>                   | Other |
| 16701 | Lanternfish            | <i>Diaphus schmidti</i>                   | Other |
| 18652 | Bythitid               | <i>Dinematichthys<br/>iluocoetenoides</i> | Other |
| 44903 | Porcupinefish          | <i>Diodon eydouxi</i>                     | Other |
| 44901 | Porcupinefish          | <i>Diodon hystrix</i>                     | Other |
| 44902 | Porcupinefish          | <i>Diodon liturosus</i>                   | Other |
| 44900 | Porcupinefish          | Diodontidae                               | Other |
| 43503 | Dragonet               | <i>Diplogrammus goramensis</i>            | Other |
| 8801  | Bristlemouth           | <i>Diplophos sp</i>                       | Other |
| 35067 | White-Spot Damsel      | <i>Dischistodus chrysopoecilus</i>        | Other |
| 35068 | Black-Vent Damsel      | <i>Dischistodus melanotus</i>             | Other |
| 35032 | White Damsel           | <i>Dischistodus perspicillatus</i>        | Other |
| 25808 | Banded Pipefish        | <i>Doryramphus<br/>dactyliophorus</i>     | Other |
| 25807 | Bluestripe Pipefish    | <i>Doryramphus excisus</i>                | Other |

|       |                      |  |       |
|-------|----------------------|--|-------|
| 25826 | Janss' Pipefish      | <i>Doryramphus janssi</i>                | Other |
| 25827 | Negros Pipefish      | <i>Doryramphus negrosensis negrsensi</i> | Other |
| 4303  | Sprat                | <i>Dussumieria elopsoides</i>            | Other |
| 4302  | Sprats               | <i>Dussumieria sp.B</i>                  | Other |
| 31300 | Diskfishes           | Echeneidae                               | Other |
| 31304 | Remora               | <i>Echeneis naucrates</i>                | Other |
| 5603  | Whiteface Moray      | <i>Echidna leucotaenia</i>               | Other |
| 5604  | Snowflake Moray      | <i>Echidna nebulosa</i>                  | Other |
| 5605  | Girdled Moray Eel    | <i>Echidna polyzona</i>                  | Other |
| 5606  | Unicolor Moray       | <i>Echidna unicolor</i>                  | Other |
| 1350  | Bramble Shark        | <i>Echinorhinidae</i>                    | Other |
| 1351  | Bramble Shark        | <i>Echinorhinus brucus</i>               | Other |
| 1352  | Bramble Shark        | <i>Echinorhinus cookei</i>               | Other |
| 39264 | Banda Clown Blenny   | <i>Ecsenius bandanus</i>                 | Other |
| 39208 | Blenny               | <i>Ecsenius bicolor</i>                  | Other |
| 39209 | Blenny               | <i>Ecsenius opsifrontalis</i>            | Other |
| 39245 | Blenny               | <i>Ecsenius sellifer</i>                 | Other |
| 39246 | Blenny               | <i>Ecsenius yaeyamaensis</i>             | Other |
| 6621  | Snake Eel            | <i>Elapsopsis versicolor</i>             | Other |
| 40400 | Sleepers             | Eleotrididae                             | Other |
| 40401 | Gudgeon              | <i>Eleotris fusca</i>                    | Other |
| 32201 | Bonnetmouth          | <i>Emmelichthys karnellai</i>            | Other |
| 32200 | Bonnet Mouths        | <i>Emmelichtyidae</i>                    | Other |
| 18703 | Pearlfish            | <i>Encheliophis boraboraensis</i>        | Other |
| 18705 | Pearlfish            | <i>Encheliophis gracilis</i>             | Other |
| 18701 | Pearlfish            | <i>Encheliophis homei</i>                | Other |
| 18704 | Pearlfish            | <i>Encheliophis vermicularis</i>         | Other |
| 5607  | Bayer'S Moray        | <i>Enchelycore bayeri</i>                | Other |
| 5608  | Bikini Atoll Moray   | <i>Enchelycore bikiniensis</i>           | Other |
| 5655  | Dark-Spotted Moray   | <i>Enchelycore kamara</i>                | Other |
| 5609  | White-Margined Moray | <i>Enchelycore schismatorhynchus</i>     | Other |
| 5610  | Viper Moray          | <i>Enchelynassa canina</i>               | Other |
| 39210 | Blenny               | <i>Enchelyurus kraussi</i>               | Other |
| 4406  | Gold Anchovy         | <i>Enchrasicholina devisi</i>            | Other |
| 4405  | Blue Anchovy         | <i>Enchrasicholina heterolobus</i>       | Other |
| 4401  | Oceanic Anchovy      | <i>Enchrasicholina punctifer</i>         | Other |
| 4400  | Anchovies            | Engraulidae                              | Other |

|       |                 |   |       |
|-------|-----------------|---|-------|
| 43904 | Flounder        | <i>Engyprosopon sp</i>                    | Other |
| 39001 | Triplefin       | <i>Enneapterygius hemimelas</i>           | Other |
| 39002 | Triplefin       | <i>Enneapterygius minutus</i>             | Other |
| 39003 | Triplefin       | <i>Enneapterygius nanus</i>               | Other |
| 39247 | Blenny          | <i>Entomacrodus caudofasciatus</i>        | Other |
| 39248 | Blenny          | <i>Entomacrodus cymatobiotus</i>          | Other |
| 39211 | Blenny          | <i>Entomacrodus decussatus</i>            | Other |
| 39212 | Blenny          | <i>Entomacrodus niuafoensis</i>           | Other |
| 39213 | Blenny          | <i>Entomacrodus sealei</i>                | Other |
| 39241 | Blenny          | <i>Entomacrodus stellifer</i>             | Other |
| 39214 | Blenny          | <i>Entomacrodus striatus</i>              | Other |
| 39215 | Blenny          | <i>Entomacrodus thalassinus thalassin</i> | Other |
| 34000 | Batfish         | Ephippidae                                | Other |
| 32202 | Bonnetmouth     | <i>Erythrocles scintillans</i>            | Other |
| 1301  | Spiny Dogfish   | <i>Etmopterus pusillus</i>                | Other |
| 20757 | Ribbon Halfbeak | <i>Euleptorhamphus viridis</i>            | Other |
| 28601 | Dragon Fish     | <i>Eurypegasmus draconis</i>              | Other |
| 4304  | Mantis Shrimp   | <i>Eutremus teres</i>                     | Other |
| 40561 | Kawakawa        | <i>Eviota afelei</i>                      | Other |
| 40562 | Herring         | <i>Eviota albolineata</i>                 | Other |
| 40563 | Goby            | <i>Eviota bifasciata</i>                  | Other |
| 40564 | Goby            | <i>Eviota cometa</i>                      | Other |
| 40565 | Goby            | <i>Eviota distigma</i>                    | Other |
| 40566 | Goby            | <i>Eviota fasciola</i>                    | Other |
| 40567 | Goby            | <i>Eviota herrei</i>                      | Other |
| 40568 | Goby            | <i>Eviota infulata</i>                    | Other |
| 40569 | Goby            | <i>Eviota lachdebrerei</i>                | Other |
| 40570 | Goby            | <i>Eviota latifasciata</i>                | Other |
| 40571 | Goby            | <i>Eviota melasma</i>                     | Other |
| 40572 | Goby            | <i>Eviota nebulosa</i>                    | Other |
| 40573 | Goby            | <i>Eviota pellucida</i>                   | Other |
| 40574 | Goby            | <i>Eviota prasina</i>                     | Other |
| 40575 | Goby            | <i>Eviota prasites</i>                    | Other |
| 40576 | Goby            | <i>Eviota punctulata</i>                  | Other |
| 40577 | Goby            | <i>Eviota queenslandica</i>               | Other |
| 40579 | Goby            | <i>Eviota saipanensis</i>                 | Other |
| 40578 | Goby            | <i>Eviota sebreei</i>                     | Other |
| 40580 | Goby            | <i>Eviota sigillata</i>                   | Other |

|       |                            |                                 |       |
|-------|----------------------------|---------------------------------|-------|
| 40581 | Goby                       | <i>Eviota smaragdus</i>         | Other |
| 40585 | Goby                       | <i>Eviota sp</i>                | Other |
| 40582 | Goby                       | <i>Eviota sparsa</i>            | Other |
| 40583 | Goby                       | <i>Eviota storthynx</i>         | Other |
| 40584 | Goby                       | <i>Eviota zonura</i>            | Other |
| 6622  | Snake Eel                  | <i>Evipes percinctus</i>        | Other |
| 39216 | Blenny                     | <i>Exalias brevis</i>           | Other |
| 20600 | Flying Fish                | <i>Exocoetidae</i>              | Other |
| 20611 | Flying Fish                | <i>Exocoetus volitans</i>       | Other |
| 40586 | Goby                       | <i>Exyrias belissimus</i>       | Other |
| 40587 | Goby                       | <i>Exyrias puntang</i>          | Other |
| 25401 | Cornetfish                 | <i>Fistularia commersoni</i>    | Other |
| 25400 | Cornetfish                 | Fistulariidae                   | Other |
| 30453 | Bay Cardinalfish           | <i>Foa brachygramma</i>         | Other |
| 30454 | Cardinalfish               | <i>Foa sp</i>                   | Other |
| 34320 | Longnosed Butterflyfish    | <i>Forcipiger flavissimus</i>   | Other |
| 34321 | Big Longnose Butterflyfish | <i>Forcipiger longirostris</i>  | Other |
| 30467 | Cardinalfish               | <i>Fowleria abocellata</i>      | Other |
| 30426 | Marbled Cardinalfish       | <i>Fowleria marmorata</i>       | Other |
| 30425 | Spotcheek Cardinalfish     | <i>Fowleria punctulata</i>      | Other |
| 30427 | Variegated Cardinalfish    | <i>Fowleria variegatus</i>      | Other |
| 40588 | Goby                       | <i>Fusigobius longispinus</i>   | Other |
| 40589 | Goby                       | <i>Fusigobius neophytus</i>     | Other |
| 1107  | Tiger Shark                | <i>Galeocerdo cuvier</i>        | Other |
| 31802 | Lg-Toothed Ponyfish        | <i>Gazza achlamys</i>           | Other |
| 31808 | Toothed Ponyfish           | <i>Gazza minuta</i>             | Other |
| 34362 | Ornate Angelfish           | <i>Genicanthus bellus</i>       | Other |
| 34371 | Black-Spot Angelfish       | <i>Genicanthus melanospilos</i> | Other |
| 34364 | Watanabe'S Angelfish       | <i>Genicanthus watanabei</i>    | Other |
| 32600 | Mojarras                   | Gerreidae                       | Other |
| 32602 | Deep-Bodied Mojarra        | <i>Gerres abbreviatus</i>       | Other |
| 32601 | Common Mojarra             | <i>Gerres acinaces</i>          | Other |
| 32604 | Filamentous Mojarra        | <i>Gerres filamentosus</i>      | Other |
| 32603 | Oblong Mojarra             | <i>Gerres oblongus</i>          | Other |
| 32605 | Oyena Mojarra              | <i>Gerres oyena</i>             | Other |
| 32606 | Mojarra                    | <i>Gerres punctatus</i>         | Other |
| 9200  | Telescopefish              | <i>Giganturidae</i>             | Other |
| 40591 | Goby                       | <i>Gladigobius ensifera</i>     | Other |
| 40592 | Goby                       | <i>Glossogobius biocellatus</i> | Other |

|       |                          |                                    |       |
|-------|--------------------------|------------------------------------|-------|
| 40593 | Goby                     | <i>Glossogobius celebius</i>       | Other |
| 40594 | Goby                     | <i>Glossogobius girus</i>          | Other |
| 39249 | Blenny                   | <i>Glyptoparus delicatulus</i>     | Other |
| 40595 | Goby                     | <i>Gnatholepis anjerensis</i>      | Other |
| 40601 |                          | <i>Gnatholepis caurensis</i>       | Other |
| 40596 | Goby                     | <i>Gnatholepis scapulostigma</i>   | Other |
| 40597 | Goby                     | <i>Gnatholepis sp.A</i>            | Other |
| 43400 | Clingfish                | Gobiesocidae                       | Other |
| 40500 | Goby                     | Gobiidae                           | Other |
| 40598 | Goby                     | <i>Gobiodon albofasciatus</i>      | Other |
| 40599 | Goby                     | <i>Gobiodon citrinus</i>           | Other |
| 40602 | Goby                     | <i>Gobiodon okinawae</i>           | Other |
| 40603 | Goby                     | <i>Gobiodon quinquestrigatus</i>   | Other |
| 40604 | Goby                     | <i>Gobiodon rivulatus</i>          | Other |
| 40605 | Goby                     | <i>Gobiopsis bravoii</i>           | Other |
| 8802  | Bristlemouth             | <i>Gonostoma atlanticum</i>        | Other |
| 8803  | Bristlemouth             | <i>Gonostoma ebelingi</i>          | Other |
| 8800  | Bristlemouths            | Gonostomatidae                     | Other |
| 6209  | Orange-Barred Garden Eel | <i>Gorgasia preclara</i>           | Other |
| 6203  | Conger Eel               | <i>Gorgasia sp</i>                 | Other |
| 29051 | Goldies                  | <i>Grammatonotus sp.1</i>          | Other |
| 29052 | Goldies                  | <i>Grammatonotus sp.2</i>          | Other |
| 41604 | 2-Lined Mackerel         | <i>Grammatorcynus bilineatus</i>   | Other |
| 29002 | Yellowstripe Soapfish    | <i>Grammistes sexlineatus</i>      | Other |
| 29000 | Soapfish                 | Grammistidae                       | Other |
| 29003 | Ocellate Soapfish        | <i>Grammistops ocellatus</i>       | Other |
| 41001 | Wormfish                 | <i>Gunnellichthys monostigma</i>   | Other |
| 41002 | Onestripe Wormfish       | <i>Gunnellichthys pleurotaenia</i> | Other |
| 41011 | Wormfish                 | <i>Gunnellichthys viridescens</i>  | Other |
| 30460 | Philippine Cardinalfish  | <i>Gymnapogon philippinus</i>      | Other |
| 30447 | Cardinalfish             | <i>Gymnapogon uros pilotus</i>     | Other |
| 32361 | Fusilier                 | <i>Gymnocaesio gymnopterus</i>     | Other |
| 5611  | Zebra Moray              | <i>Gymnomuraena zebra</i>          | Other |
| 5619  | Moray Eel                | <i>Gymnothorax berndti</i>         | Other |
| 5620  | Buro Moray               | <i>Gymnothorax buroensis</i>       | Other |
| 5624  | Moray Eel                | <i>Gymnothorax elegans</i>         | Other |
| 5635  | Enigmatic Moray          | <i>Gymnothorax enigmaticus</i>     | Other |
| 5621  | Fimbriated Moray         | <i>Gymnothorax fimbriatus</i>      | Other |
| 5622  | Yellow-Margined Moray    | <i>Gymnothorax flavimarginatus</i> | Other |

|       |                       |                                      |       |
|-------|-----------------------|--------------------------------------|-------|
| 5612  | Brown Spotted Moray   | <i>Gymnothorax fuscomaculatus</i>    | Other |
| 5623  | Graceful-Tailed Moray | <i>Gymnothorax gracilicaudus</i>     | Other |
| 5625  | Moray Eel             | <i>Gymnothorax hepaticus</i>         | Other |
| 5626  | Giant Moray           | <i>Gymnothorax javanicus</i>         | Other |
| 5627  | Blotch-Necked Moray   | <i>Gymnothorax margaritophorus</i>   | Other |
| 5613  | Marshall Isles Moray  | <i>Gymnothorax marshallensis</i>     | Other |
| 5614  | Dirty Yellow Moray    | <i>Gymnothorax melatremus</i>        | Other |
| 5628  | Whitemouth Moray      | <i>Gymnothorax meleagris</i>         | Other |
| 5648  | Monochrome Moray      | <i>Gymnothorax monochrous</i>        | Other |
| 5629  | 1-Spot Moray          | <i>Gymnothorax monostigmus</i>       | Other |
| 5630  | Moray Eel             | <i>Gymnothorax neglectus</i>         | Other |
| 5645  | Yellowmouth Moray     | <i>Gymnothorax nudivomer</i>         | Other |
| 5616  | Pinda Moray           | <i>Gymnothorax pindae</i>            | Other |
| 5649  | Moray Eel             | <i>Gymnothorax polyuranodon</i>      | Other |
| 5631  | Richardson'S Moray    | <i>Gymnothorax richardsoni</i>       | Other |
| 5632  | Yellow-Headed Moray   | <i>Gymnothorax rueppelliae</i>       | Other |
| 5618  | Moray Eel             | <i>Gymnothorax sp.cf Melatremus</i>  | Other |
| 5633  | Undulated Moray       | <i>Gymnothorax undulatus</i>         | Other |
| 5634  | Zonipectis Moray      | <i>Gymnothorax zonipectus</i>        | Other |
| 32700 | Sweetlips             | <i>Haemulidae</i>                    | Other |
| 25811 | Brock'S Pipefish      | <i>Halicampus brocki</i>             | Other |
| 25828 | Duncker'S Pipefish    | <i>Halicampus dunckeri</i>           | Other |
| 25812 | Samoan Pipefish       | <i>Halicampus mataafae</i>           | Other |
| 25829 | Glittering Pipefish   | <i>Halicampus nitidus</i>            | Other |
| 44301 | Spikefish             | <i>Halimochirurgus alcocki</i>       | Other |
| 39004 | Triplefin             | <i>Helcogramma capidata</i>          | Other |
| 39005 | Triplefin             | <i>Helcogramma chica</i>             | Other |
| 39006 | Triplefin             | <i>Helcogramma hudsoni</i>           | Other |
| 35069 | Damselfish            | <i>Hemiglyphidodon plagiometopon</i> | Other |
| 20751 | Halfbeak              | <i>Hemiramphus archipelagicus</i>    | Other |
| 20758 | Halfbeak              | <i>Hemiramphus far</i>               | Other |
| 20760 | Halfbeak              | <i>Hemiramphus lutkei</i>            | Other |
| 20750 | Halfbeak              | <i>Hemirhamphidae</i>                | Other |
| 34322 | Pyrimid Butterflyfish | <i>Hemitaurichthys polylepis</i>     | Other |
| 34323 | Butterflyfish         | <i>Hemitaurichthys thompsoni</i>     | Other |
| 34324 | Longfinned Bannerfish | <i>Heniochus acuminatus</i>          | Other |

|       |                        |  |       |
|-------|------------------------|--|-------|
| 34325 | Pennant Bannerfish     | <i>Heniochus chrysostomus</i>          | Other |
| 34341 | Bannerfish             | <i>Heniochus diphreutes</i>            | Other |
| 34326 | Masked Bannerfish      | <i>Heniochus monoceros</i>             | Other |
| 34327 | Singular Butterflyfish | <i>Heniochus singularis</i>            | Other |
| 34328 | Humphead Bannerfish    | <i>Heniochus varius</i>                | Other |
| 4308  | Gold Spot Herring      | <i>Herklotsichthys quadrimaculatus</i> | Other |
| 6204  | Conger Eel             | <i>Heteroconger hassi</i>              | Other |
| 40606 | Goby                   | <i>Heteroeleotris sp</i>               | Other |
| 30301 | Glasseye               | <i>Heteropriacanthus cruentatus</i>    | Other |
| 2006  | Whipray                | <i>Himantura fai</i>                   | Other |
| 2005  | Wh Tail Whipray        | <i>Himantura granulata</i>             | Other |
| 2003  | Leopard Ray            | <i>Himantura uarnak</i>                | Other |
| 25830 | Pipefish               | <i>Hippichthys cyanospilos</i>         | Other |
| 25831 | Pipefish               | <i>Hippichthys spicifer</i>            | Other |
| 25809 | Pipefish               | <i>Hippocampus histrix</i>             | Other |
| 25832 | Pipefish               | <i>Hippocampus kuda</i>                | Other |
| 19212 | Sargassum Fish         | <i>Histrion histrio</i>                | Other |
| 28965 | Fairy Basslet          | <i>Holanthias borbonius</i>            | Other |
| 28966 | Fairy Basslet          | <i>Holanthias katayamai</i>            | Other |
| 30801 | Tilefish               | <i>Hoplolatilus cuniculus</i>          | Other |
| 30802 | Tilefish               | <i>Hoplolatilus fronticinctus</i>      | Other |
| 30803 | Tilefish               | <i>Hoplolatilus starcki</i>            | Other |
| 21807 | Silverside             | <i>Hypoatherina barnesi</i>            | Other |
| 21808 | Silverside             | <i>Hypoatherina cylindrica</i>         | Other |
| 21802 | Silverside             | <i>Hypoatherina ovalaua</i>            | Other |
| 20753 | Halfbeak               | <i>Hyporhamphus acutus acutus</i>      | Other |
| 20754 | Halfbeak               | <i>Hyporhamphus affinis</i>            | Other |
| 20755 | Halfbeak               | <i>Hyporhamphus dussumieri</i>         | Other |
| 6623  | Snake Eel              | <i>Ichthyapus vulturus</i>             | Other |
| 26430 | Spiny Devilfish        | <i>Inimicus didactylus</i>             | Other |
| 21901 | Keeled Silverside      | <i>Iso hawaiiensis</i>                 | Other |
| 35210 | 6-Band Hawkfish        | <i>Isocirrhitis sexfasciatus</i>       | Other |
| 21900 | Keeled Silversides     | Isonidae                               | Other |
| 39265 | Beautiful Rockskipper  | <i>Istiblennius bellus</i>             | Other |
| 39217 | Blenny                 | <i>Istiblennius chrysospilos</i>       | Other |
| 39266 | Streaky Rockskipper    | <i>Istiblennius dussumieri</i>         | Other |
| 39219 | Blenny                 | <i>Istiblennius edentulus</i>          | Other |

|       |                         |                                      |       |
|-------|-------------------------|--------------------------------------|-------|
| 39267 | Interrupted Rockskipper | <i>Istiblennius interruptus</i>      | Other |
| 39220 | Blenny                  | <i>Istiblennius lineatus</i>         | Other |
| 40607 | Goby                    | <i>Istigobius decoratus</i>          | Other |
| 40608 | Goby                    | <i>Istigobius ornatus</i>            | Other |
| 40609 | Goby                    | <i>Istigobius rigilius</i>           | Other |
| 40610 | Goby                    | <i>Istigobius spence</i>             | Other |
| 41900 | Billfishes              | Istiophoridae                        | Other |
| 901   | Mackerel Shark          | <i>Isurus oxyrinchus</i>             | Other |
| 5402  | Bl-Nostril False Moray  | <i>Kaupichthys atronasmus</i>        | Other |
| 5403  | Shortfin False Moray    | <i>Kaupichthys brachychirus</i>      | Other |
| 5401  | Common False Moray      | <i>Kaupichthys hyoprорoides</i>      | Other |
| 40612 | Goby                    | <i>Kelloggella quindecimfasciata</i> | Other |
| 40611 | Goby                    | <i>Kelloggella cardinalis</i>        | Other |
| 40701 | Sand Dart               | <i>Kraemeria bryani</i>              | Other |
| 40702 | Sand Dart               | <i>Kraemeria cunicularia</i>         | Other |
| 40703 | Sand Dart               | <i>Kraemeria samoensis</i>           | Other |
| 40700 | Sand Darts              | Kraemeriidae                         | Other |
| 30103 | Dark-Margined Flagtail  | <i>Kuhlia marginata</i>              | Other |
| 30101 | Barred Flagtail         | <i>Kuhlia mugil</i>                  | Other |
| 30102 | River Flagtail          | <i>Kuhlia rupestris</i>              | Other |
| 30100 | Flagtails               | Kuhliidae                            | Other |
| 44601 | Longhorn Cowfish        | <i>Lactoria cornuta</i>              | Other |
| 44602 | Spiny Cowfish           | <i>Lactoria diaphana</i>             | Other |
| 44605 | Thornback Cowfish       | <i>Lactoria fornasini</i>            | Other |
| 44817 | Oceanic Blasop          | <i>Lagocephalus lagocephalus</i>     | Other |
| 44818 | Silverstripe Blasop     | <i>Lagocephalus scleratus</i>        | Other |
| 900   |                         |                                      |       |
| 6627  | Oriental Snake Eel      | <i>Lamnostoma orientalis</i>         | Other |
| 31800 | Ponyfishes              | Leiognathidae                        | Other |
| 31806 | Slipmouth               | <i>Leiognathus bindus</i>            | Other |
| 31804 | Slipmouth               | <i>Leiognathus elongatus</i>         | Other |
| 31801 | Common Slipmouth        | <i>Leiognathus equulus</i>           | Other |
| 31805 | Slipmouth               | <i>Leiognathus smithursti</i>        | Other |
| 31803 | Oblong Slipmouth        | <i>Leiognathus stercorarius</i>      | Other |
| 6605  | Saddled Snake Eel       | <i>Leiuranus semicinctus</i>         | Other |
| 43401 | Clingfish               | <i>Lepadichthys caritus</i>          | Other |
| 43402 | Clingfish               | <i>Lepadichthys minor</i>            | Other |
| 35048 | Fusilier Damsel         | <i>Lepidozygus tapienosoma</i>       | Other |
| 16901 | Barracudina             | <i>Lestidium nudun</i>               | Other |



|       |                                |   |       |
|-------|--------------------------------|---|-------|
| 37402 | Sand Burrower                  | <i>Limnichthys donaldsoni</i>             | Other |
| 43403 | Clingfish                      | <i>Liobranchia stria</i>                  | Other |
| 28991 | Swissguard Basslet             | <i>Liopropoma lunulatum</i>               | Other |
| 28997 | Swissguard Basslet             | <i>Liopropoma maculatum</i>               | Other |
| 28992 | Swissguard Basslet             | <i>Liopropoma mitratum</i>                | Other |
| 28993 | Swissguard Basslet             | <i>Liopropoma multilineatum</i>           | Other |
| 28994 | Pallid Basslet                 | <i>Liopropoma pallidum</i>                | Other |
| 28995 | Pinstripe Basslet              | <i>Liopropoma susumi</i>                  | Other |
| 28996 | Redstripe Basslet              | <i>Liopropoma tonstrinum</i>              | Other |
| 39251 | Blenny                         | <i>Litobranchus fowleri</i>               | Other |
| 35908 | Giantscale Mullet              | <i>Liza melinoptera</i>                   | Other |
| 32501 | Triplefin                      | <i>Lobotes surinamensis</i>               | Other |
| 32500 | Tripletails                    | Lobotidae                                 | Other |
| 40519 | Goby                           | <i>Lotilia graciliosa</i>                 | Other |
| 28981 | Magenta Slender Basslet        | <i>Luzonichthys waitei</i>                | Other |
| 28982 | Whitley'S Slender Basslet      | <i>Luzonichthys whitleyi</i>              | Other |
| 40613 | Goby                           | <i>Macrodontogobius wilburi</i>           | Other |
| 40520 | Goby                           | <i>Mahidolia mystacina</i>                | Other |
| 30800 | Tilefishes                     | Malacanthidae                             | Other |
| 30851 | Quakerfish                     | <i>Malacanthus brevirostris</i>           | Other |
| 30852 | Striped Blanquillo             | <i>Malacanthus latovittatus</i>           | Other |
| 2301  | Manta Ray                      | <i>Manta birostris</i>                    | Other |
| 45001 | Sharptail Sunfish              | <i>Masturus lanceolatus</i>               | Other |
| 4700  | Tarpons                        | Megalopidae                               | Other |
| 4701  | Indo-Pacific Tarpon            | <i>Megalops cyprinoides</i>               | Other |
| 39233 | Poison-Fang Blenny             | <i>Meiacanthus anema</i>                  | Other |
| 39223 | Poison-Fang Blenny             | <i>Meiacanthus atrodorsalis</i>           | Other |
| 39258 | 1-Stripe Poison-Fang Blenny    | <i>Meiacanthus ditrema</i>                | Other |
| 39259 | Striped Poison-Fang Blenny     | <i>Meiacanthus grammistes</i>             | Other |
| 44505 | Black Triggerfish              | <i>Melichthys niger</i>                   | Other |
| 44506 | Pinktail Triggerfish           | <i>Melichthys vidua</i>                   | Other |
| 18653 | Brotula                        | <i>Microbrotula</i> sp.                   | Other |
| 41000 | Wormfish                       | Microdesmidae                             | Other |
| 25817 | Anderson'S Shrt-Nosed Pipefish | <i>Micrognathus andersonii</i>            | Other |
| 25810 | Pygmy Short-Nosed Pipefish     | <i>Micrognathus brevirostris pygmaeus</i> | Other |
| 25833 | Pipefish                       | <i>Micropphis brachyurus brachyurus</i>   | Other |

|       |                     |                                  |       |
|-------|---------------------|----------------------------------|-------|
| 25834 | Pipefish            | <i>Microphis brevidorsalis</i>   | Other |
| 25835 | Pipefish            | <i>Microphis leiaspis</i>        | Other |
| 25836 | Pipefish            | <i>Microphis manadensis</i>      | Other |
| 25837 | Pipefish            | <i>Microphis retzii</i>          | Other |
| 25813 | Ventricose Milda    | <i>Minyichthys myersi</i>        | Other |
| 2300  | Myer'S Pipefish     | Mobulidae                        | Other |
| 45000 | Ocean Sunfishes     | Molidae                          | Other |
| 44550 | Filefishes          | Monacanthidae                    | Other |
| 33300 | Monos               | Monodactylidae                   | Other |
| 33301 | Mono                | <i>Monodactylus argenteus</i>    | Other |
| 18000 | Codlings            | Moridae                          | Other |
| 5103  | Rusty Spaghetti Eel | <i>Moringua ferruginea</i>       | Other |
| 5102  | Java Spaghetti Eel  | <i>Moringua javanica</i>         | Other |
| 5101  | Spaghetti Eel       | <i>Moringua microchir</i>        | Other |
| 5100  | Worm Eel            | Moringuidae                      | Other |
| 40614 | Goby                | <i>Mugilogobius tagala</i>       | Other |
| 40615 | Goby                | <i>Mugilogobius villa</i>        | Other |
| 6300  | Pike Eels           | Muraenesocidae                   | Other |
| 6301  | Pike Conger         | <i>Muraenesox cinereus</i>       | Other |
| 6612  | Snake Eel           | <i>Muraenichthys gymnotus</i>    | Other |
| 6606  | Snake Eel           | <i>Muraenichthys laticaudata</i> | Other |
| 6607  | Snake Eel           | <i>Muraenichthys macropterus</i> | Other |
| 6613  | Snake Eel           | <i>Muraenichthys schultzi</i>    | Other |
| 6614  | Snake Eel           | <i>Muraenichthys sibogae</i>     | Other |
| 5600  | Morays              | Muraenidae                       | Other |
| 16700 | Lanternfishes       | Myctophidae                      | Other |
| 16702 | Laternfish          | <i>Myctophum brachygnathos</i>   | Other |
| 2200  | Eagle Ray           | Myliobatidae                     | Other |
| 6624  | Snake Eel           | <i>Myrichthys bleekeri</i>       | Other |
| 6608  | Banded Snake Eel    | <i>Myrichthys colubrinus</i>     | Other |
| 6610  | Spotted Snake Eel   | <i>Myrichthys maculosus</i>      | Other |
| 6615  | Snake Eel           | <i>Myrophis uropterus</i>        | Other |
| 200   | Hagfish             | Myxinidae                        | Other |
| 201   | Hagfish             | <i>Eptaptretus carlhubbsi</i>    | Other |
| 39252 | Combtooth Blenny    | <i>Nannosalarius nativitatus</i> | Other |
| 701   | Nurse Shark         | <i>Nebrius ferrugineus</i>       | Other |
| 1110  | Lemon Shark         | <i>Negaprion acutidens</i>       | Other |
| 41010 | Decorated Dartfish  | <i>Nemateleotris decora</i>      | Other |
| 41003 | Helfrichs' Dartfish | <i>Nemateleotris helfrichi</i>   | Other |

|       |                           |  |       |
|-------|---------------------------|--|-------|
| 41004 | Fire Dartfish             | <i>Nemateleotris magnifica</i>                 | Other |
| 32400 | Threadfin Breams          | Nemipteridae                                   | Other |
| 32900 | Breams                    | Nemipteridae                                   | Other |
| 32412 | Forktail Bream            | <i>Nemipterus furcosus</i>                     | Other |
| 32409 | Butterfly Bream           | <i>Nemipterus hexadon</i>                      | Other |
| 32410 | Notched Butterfly Bream   | <i>Nemipterus peronii</i>                      | Other |
| 32411 | Butterfly Bream           | <i>Nemipterus tolu</i>                         | Other |
| 35205 | Flame Hawkfish            | <i>Neocirrhites armatus</i>                    | Other |
| 35072 | Royal Damsel              | <i>Neoglyphidodon melas</i>                    | Other |
| 35073 | Yellowfin Damsel          | <i>Neoglyphidodon nigroris</i>                 | Other |
| 35070 | Coral Demoiselle          | <i>Neopomacentrus nemurus</i>                  | Other |
| 35071 | Freshwater Demoiselle     | <i>Neopomacentrus taeniurus</i>                | Other |
| 35047 | Violet Demoiselle         | <i>Neopomacentrus violascens</i>               | Other |
| 42200 | Man-Of-War Fish           | Nomeidae                                       | Other |
| 39007 | Triplefin                 | <i>Norfolkia brachylepis</i>                   | Other |
| 44507 | Redtooth Triggerfish      | <i>Odonus niger</i>                            | Other |
| 35909 | Foldlip Mullet            | <i>Oedalechilus labiosus</i>                   | Other |
| 39263 | Mangrove Blenny           | <i>Omobranchus obliquus</i>                    | Other |
| 39224 | Blenny                    | <i>Omobranchus rotundiceps</i>                 | Other |
| 39256 | Blenny                    | <i>Omx biporos</i>                             | Other |
| 18706 | Bivalve Pearlfish         | <i>Onuxodon fowleri</i>                        | Other |
| 6600  | Snake Eel                 | Ophichthidae                                   | Other |
| 6611  | Dark-Shouldered Snake Eel | <i>Ophichthus cephalozona</i>                  | Other |
| 18600 | Cusk Eel                  | Ophidiidae                                     | Other |
| 40405 | Sleeper                   | <i>Ophieleotris aporos</i>                     | Other |
| 40406 | Sleeper                   | <i>Ophiocara porocephala</i>                   | Other |
| 36600 | Jawfishes                 | Opisthognathidae                               | Other |
| 36601 | Variable Jawfish          | <i>Opisthognathus</i> sp. A                    | Other |
| 36602 | Wass' Jawfish             | <i>Opisthognathus</i> sp. B                    | Other |
| 34700 | Knifejaws                 | Oplegnathidae                                  | Other |
| 34701 | Spotted Knifejaw          | <i>Oplegnathus punctatus</i>                   | Other |
| 40528 | Goby                      | <i>Oplopomops diacanthus</i>                   | Other |
| 40529 | Goby                      | <i>Oplopomus oplopomus</i>                     | Other |
| 40616 | Goby                      | <i>Opua nephodes</i>                           | Other |
| 700   | Nurse,Zebra,Carpet Sharks | Orectolobidae                                  | Other |
| 34601 | Tilapia                   | <i>Oreochromis mossambicus</i>                 | Other |
| 44600 | Boxfish, Cowfish          | Ostraciidae                                    | Other |
| 44603 | Cube Trunkfish            | <i>Ostracion cubicus</i>                       | Other |
| 44604 | Spotted Trunkfish         | <i>Ostracion meleagris</i><br><i>meleagris</i> | Other |

|       |                         |  |       |
|-------|-------------------------|--|-------|
| 44606 | Reticulate Boxfish      | <i>Ostracion solorensis</i>                  | Other |
| 35206 | Longnose Hawkfish       | <i>Oxycirrhites typus</i>                    | Other |
| 40407 | Sleeper                 | <i>Oxyleotris lineolatus</i>                 | Other |
| 44555 | Longnose Filefish       | <i>Oxymonacanthus longirostris</i>           | Other |
| 20759 | Smallwing Flying Fish   | <i>Oxyporhamphus micropterus micropterus</i> | Other |
| 40617 | Goby                    | <i>Oxyurichthys guibei</i>                   | Other |
| 40618 | Goby                    | <i>Oxyurichthys microlepis</i>               | Other |
| 40619 | Goby                    | <i>Oxyurichthys ophthalmonema</i>            | Other |
| 40620 | Goby                    | <i>Oxyurichthys papuensis</i>                | Other |
| 40621 | Goby                    | <i>Oxyurichthys tentacularis</i>             | Other |
| 40622 | Goby                    | <i>Padanka</i> sp.                           | Other |
| 40623 | Goby                    | <i>Palutris pruinosa</i>                     | Other |
| 40624 | Goby                    | <i>Palutris reticularis</i>                  | Other |
| 35207 | Arc-Eyed Hawkfish       | <i>Paracirrhites arcatus</i>                 | Other |
| 35208 | Freckled Hawkfish       | <i>Paracirrhites forsteri</i>                | Other |
| 35209 | Whitespot Hawkfish      | <i>Paracirrhites hemistictus</i>             | Other |
| 40625 | Goby                    | <i>Paragobiodon echinocephalus</i>           | Other |
| 40626 | Goby                    | <i>Paragobiodon lacunicolus</i>              | Other |
| 40627 | Goby                    | <i>Paragobiodon melanosoma</i>               | Other |
| 40628 | Goby                    | <i>Paragobiodon modestus</i>                 | Other |
| 40629 | Goby                    | <i>Paragobiodon xanthosoma</i>               | Other |
| 41012 | Seychelle'S Wormfish    | <i>Paragunnellichthy seychellensis</i>       | Other |
| 16900 | Barracudinas            | Paralepididae                                | Other |
| 44556 | Blacksaddle Mimic       | <i>Paraluteres prionurus</i>                 | Other |
| 44560 | Filefish                | <i>Paramonacanthus cryptodon</i>             | Other |
| 44561 | Filefish                | <i>Paramonacanthus japonicus</i>             | Other |
| 37102 | Latticed Sandperch      | <i>Parapercis clathrata</i>                  | Other |
| 37103 | Cylindrical Sandperch   | <i>Parapercis cylindrica</i>                 | Other |
| 37101 | Blk-Dotted Sandperch    | <i>Parapercis millipunctata</i>              | Other |
| 37105 | Red-Barred Sandperch    | <i>Parapercis multiplicata</i>               | Other |
| 37106 | Black-Banded Sandperch  | <i>Parapercis tetracantha</i>                | Other |
| 37104 | Blotchlip Sandperch     | <i>Parapercis xanthozona</i>                 | Other |
| 33402 | Sandperch               | <i>Parapriacanthus ransonneti</i>            | Other |
| 26433 | Meadam'S Scorpionfish   | <i>Parascorpaena mcadamsi</i>                | Other |
| 26426 | Mozambique Scorpionfish | <i>Parascorpaena mossambica</i>              | Other |

|       |                       |                                       |       |
|-------|-----------------------|---------------------------------------|-------|
| 44105 | Peacock Sole          | <i>Pardachirus pavoninus</i>          | Other |
| 39225 | Blenny                | <i>Parenchelyurus hepburni</i>        | Other |
| 20607 | Flying Fish           | <i>Parexocoetus brachypterus</i>      | Other |
| 20608 | Flying Fish           | <i>Parexocoetus mento</i>             | Other |
| 41013 | Beautiful Hover Goby  | <i>Parioglossus formosus</i>          | Other |
| 41014 | Lined Hover Goby      | <i>Parioglossus lineatus</i>          | Other |
| 41015 | Naked Hover Goby      | <i>Parioglossus nudus</i>             | Other |
| 41016 | Palustris Hover Goby  | <i>Parioglossus palustris</i>         | Other |
| 41017 | Rainford'S Hover Goby | <i>Parioglossus rainfordi</i>         | Other |
| 41018 | Rao'S Hover Goby      | <i>Parioglossus raoi</i>              | Other |
| 41019 | Taeniatus Hover Goby  | <i>Parioglossus taeniatus</i>         | Other |
| 41020 | Vertical Hover Goby   | <i>Parioglossus verticalis</i>        | Other |
| 2007  | Shortsnouted Ray      | <i>Pasinachus sephen</i>              | Other |
| 28600 | Dragonfish            | Pegasidae                             | Other |
| 33400 | Sweepers              | <i>Pempherididae</i>                  | Other |
| 33401 | Bronze Sweeper        | <i>Pempheris oualensis</i>            | Other |
| 34500 | Armourheads           | <i>Pentacerotidae</i>                 | Other |
| 32901 | Smalltooth Whiptail   | <i>Pentapodus caninus</i>             | Other |
| 32902 | 3-Striped Whiptail    | <i>Pentapodus trivittatus</i>         | Other |
| 37000 | Duckbills             | Percophidae                           | Other |
| 40630 | Goby                  | <i>Periophthalmus argentilineatus</i> | Other |
| 40631 | Goby                  | <i>Periophthalmus kalolo</i>          | Other |
| 44567 | Yelloweye Filefish    | <i>Pervagor alternans</i>             | Other |
| 44562 | Orangetail Filefish   | <i>Pervagor aspricaudatus</i>         | Other |
| 44557 | Blackbar Filefish     | <i>Pervagor janthinosoma</i>          | Other |
| 44563 | Blackheaded Filefish  | <i>Pervagor melanocephalus</i>        | Other |
| 44564 | Blacklined Filefish   | <i>Pervagor nigrolineatus</i>         | Other |
| 39260 | Blenny                | <i>Petroscirtes breviceps</i>         | Other |
| 39226 | Blenny                | <i>Petroscirtes mitratus</i>          | Other |
| 39261 | Blenny                | <i>Petroscirtes thepassi</i>          | Other |
| 39262 | Blenny                | <i>Petroscirtes variabilis</i>        | Other |
| 39227 | Blenny                | <i>Petroscirtes xestus</i>            | Other |
| 6625  | Snake Eel             | <i>Phenamomas cooperi</i>             | Other |
| 24202 | Flashlightfish        | <i>Photoblepheron palpebratus</i>     | Other |
| 25814 | Pipefish              | <i>Phoxocampus diacanthus</i>         | Other |
| 6626  | Snake Eel             | <i>Phyllophichthus xenodontus</i>     | Other |
| 18001 | Codling               | <i>Physiculus</i> sp.                 | Other |
| 37100 | Sand Perch            | Pinguipedidae                         | Other |
| 39228 | Blenny                | <i>Plagiotremus laudandus</i>         | Other |

|       |                        |  |       |
|-------|------------------------|--|-------|
| 39229 | Red Sabbertooth Blenny | <i>Plagiotremus rhynorhynchus</i>      | Other |
| 39230 | Blenny                 | <i>Plagiotremus tapienosoma</i>        | Other |
| 34001 | Batfish                | <i>Platax orbicularis</i>              | Other |
| 34002 | Pinnate Spadefish      | <i>Platax pinnatus</i>                 | Other |
| 34003 | Longfin Spadefish      | <i>Platax teira</i>                    | Other |
| 20702 | Keeled Needlefish      | <i>Platybelone argalus platyura</i>    | Other |
| 27300 | Flathead               | Platycephalidae                        | Other |
| 32710 | 2-Lined Sweetlips      | <i>Plectorhinchus albovittatus</i>     | Other |
| 32706 | Celebes Sweetlips      | <i>Plectorhinchus celebecus</i>        | Other |
| 32707 | Harlequin Sweetlips    | <i>Plectorhinchus chaetodonoides</i>   | Other |
| 32712 | Sweetlip               | <i>Plectorhinchus flavomaculatus</i>   | Other |
| 32703 | Gibbus Sweetlips       | <i>Plectorhinchus gibbosus</i>         | Other |
| 32708 | Lined Sweetlips        | <i>Plectorhinchus lessonii</i>         | Other |
| 32709 | Goldman'S Sweetlips    | <i>Plectorhinchus lineatus</i>         | Other |
| 32705 | Giant Sweetlips        | <i>Plectorhinchus obscurus</i>         | Other |
| 32704 | Spotted Sweetlips      | <i>Plectorhinchus picus</i>            | Other |
| 32713 | Sweetlip               | <i>Plectorhinchus sp</i>               | Other |
| 32702 | Oriental Sweetlips     | <i>Plectorhinchus vittatus</i>         | Other |
| 28987 | Fourmanoir'S Basslet   | <i>Plectranthias fourmanoiri</i>       | Other |
| 28968 | Basslet                | <i>Plectranthias kamii</i>             | Other |
| 28985 | Long-Finned Basslet    | <i>Plectranthias longimanus</i>        | Other |
| 28969 | Pygmy Basslet          | <i>Plectranthias nanus</i>             | Other |
| 28990 | Basslet                | <i>Plectranthias rubrifasciatus</i>    | Other |
| 28986 | Basslet                | <i>Plectranthias winniensis</i>        | Other |
| 35033 | Dick'S Damsel          | <i>Plectroglyphidodo dickii</i>        | Other |
| 35034 | Bright-Eye Damsel      | <i>Plectroglyphidodo imparipennis</i>  | Other |
| 35035 | Johnston Isle Damsel   | <i>Plectroglyphidodo johnstonianus</i> | Other |
| 35036 | Jewel Damsel           | <i>Plectroglyphidodo lacrymatus</i>    | Other |
| 35037 | White-Band Damsel      | <i>Plectroglyphidodo leucozonus</i>    | Other |
| 35038 | Phoenix Isle Damsel    | <i>Plectroglyphidodo phoenixensis</i>  | Other |
| 29400 | Longfins               | Plesiopidae                            | Other |
| 29402 | Red-Tipped Longfin     | <i>Plesiops caeruleolineatus</i>       | Other |

|       |                        |                                       |       |
|-------|------------------------|---------------------------------------|-------|
| 29403 | Bluegill Longfin       | <i>Plesiops corallicola</i>           | Other |
| 29405 | Sharp-Nosed Longfin    | <i>Plesiops oxycephalus</i>           | Other |
| 40632 | Goby                   | <i>Pleurosicya bilobatus</i>          | Other |
| 40664 | Caroline Ghost Goby    | <i>Pleurosicya carolinensis</i>       | Other |
| 40665 | Blue Coral Ghost Goby  | <i>Pleurosicya coerulea</i>           | Other |
| 40666 | Fringed Ghost Goby     | <i>Pleurosicya fringella</i>          | Other |
| 40667 | Michael'S Ghost Goby   | <i>Pleurosicya micheli</i>            | Other |
| 40668 | Common Ghost Goby      | <i>Pleurosicya mossambica</i>         | Other |
| 40633 | Goby                   | <i>Pleurosicya muscarum</i>           | Other |
| 40669 | Plicata Ghost Goby     | <i>Pleurosicya plicata</i>            | Other |
| 14900 | Eel Catfishes          | Plotosidae                            | Other |
| 14901 | Striped Eel Catfish    | <i>Plotosus lineatus</i>              | Other |
| 6207  | Barred Sand Conger     | <i>Poecilococongus fasciatus</i>      | Other |
| 29004 | Spotted Soapfish       | <i>Pogonoperca punctata</i>           | Other |
| 36101 | 6 Feeler Threadfin     | <i>Polydactylus sexfilis</i>          | Other |
| 17501 | Beardfish              | <i>Polymixia japonica</i>             | Other |
| 17500 | Beardfish              | Polymixiidae                          | Other |
| 36100 | Threadfins             | Polynemidae                           | Other |
| 34350 | Angelfishes            | Pomacanthidae                         | Other |
| 34365 | Emperor Angelfish      | <i>Pomacanthus imperator</i>          | Other |
| 34372 | Blue-Girdled Angelfish | <i>Pomacanthus navarchus</i>          | Other |
| 34375 | Semicircle Angelfish   | <i>Pomacanthus<br/>semicirculatus</i> | Other |
| 34373 | 6-Banded Angelfish     | <i>Pomacanthus sexstriatus</i>        | Other |
| 34374 | Blue-Faced Angelfish   | <i>Pomacanthus<br/>xanthometopon</i>  | Other |
| 35000 | Damselfishes           | Pomacentridae                         | Other |
| 35087 | Damselfish             | <i>Pomacentrus adelus</i>             | Other |
| 35039 | Ambon Damsel           | <i>Pomacentrus amboinensis</i>        | Other |
| 35094 | Goldbelly Damsel       | <i>Pomacentrus auriventris</i>        | Other |
| 35074 | Speckled Damsel        | <i>Pomacentrus bankanensis</i>        | Other |
| 35081 | Charcoal Damsel        | <i>Pomacentrus brachialis</i>         | Other |
| 35075 | Burrough'S Damsel      | <i>Pomacentrus burroughi</i>          | Other |
| 35084 | White-Tail Damsel      | <i>Pomacentrus chrysurus</i>          | Other |
| 35076 | Neon Damsel            | <i>Pomacentrus coelestis</i>          | Other |
| 35077 | Outer Reef Damsel      | <i>Pomacentrus emarginatus</i>        | Other |
| 35078 | Blue-Spot Damsel       | <i>Pomacentrus<br/>grammorhynchus</i> | Other |
| 35092 | Lemon Damsel           | <i>Pomacentrus moluccensis</i>        | Other |
| 35086 | Nagasaki Damsel        | <i>Pomacentrus nagasakiensis</i>      | Other |

|       |                         |                                   |       |
|-------|-------------------------|-----------------------------------|-------|
| 35093 | Black-Axil Damsel       | <i>Pomacentrus nigromanus</i>     | Other |
| 35040 | Sapphire Damsel         | <i>Pomacentrus pavo</i>           | Other |
| 35082 | Philippine Damsel       | <i>Pomacentrus philippinus</i>    | Other |
| 35083 | Reid'S Damsel           | <i>Pomacentrus reidi</i>          | Other |
| 35085 | Blueback Damsel         | <i>Pomacentrus simsiang</i>       | Other |
| 35041 | Princess Damsel         | <i>Pomacentrus vaiuli</i>         | Other |
| 35088 | Slender Reef-Damsel     | <i>Pomachromis exilis</i>         | Other |
| 35042 | Guam Damsel             | <i>Pomachromis guamensis</i>      | Other |
| 32711 | Common Javelinefish     | <i>Pomadasyus kaakan</i>          | Other |
| 26404 | Lg-Headed Scorpionfish  | <i>Pontinus macrocephalus</i>     | Other |
| 26431 | Scorpionfish            | <i>Pontinus</i> sp.               | Other |
| 26452 | Scopionfish             | <i>Pontinus tentacularis</i>      | Other |
| 39231 | Blenny                  | <i>Prealticus amboinensis</i>     | Other |
| 39232 | Blenny                  | <i>Prealticus natalis</i>         | Other |
| 30300 | Bigeyes                 | Priacanthidae                     | Other |
| 30305 | Bigeye                  | <i>Priacanthus alalaua</i>        | Other |
| 30302 | Goggle-Eye              | <i>Priacanthus hamrur</i>         | Other |
| 40634 | Goby                    | <i>Priolepis cincta</i>           | Other |
| 40635 | Goby                    | <i>Priolepis farcimen</i>         | Other |
| 40636 | Goby                    | <i>Priolepis inhaca</i>           | Other |
| 40637 | Goby                    | <i>Priolepis semidoliatus</i>     | Other |
| 30303 | Bigeye                  | <i>Pristigenys meyeri</i>         | Other |
| 20609 | Flying Fish             | <i>Prognichthys albimaculatus</i> | Other |
| 20610 | Flying Fish             | <i>Prognichthys sealei</i>        | Other |
| 42201 | Freckeled Driftfish     | <i>Psenes cyanophrys</i>          | Other |
| 44568 | Rhino Leatherjacket     | <i>Pseudalutarias nasicornis</i>  | Other |
| 30448 | Cardinalfish            | <i>Pseudamia amblyuroptera</i>    | Other |
| 30449 | Cardinalfish            | <i>Pseudamia gelatinosa</i>       | Other |
| 30450 | Cardinalfish            | <i>Pseudamia hayashii</i>         | Other |
| 30461 | Cardinalfish            | <i>Pseudamia zonata</i>           | Other |
| 30428 | Cardinalfish            | <i>Pseudamiops gracilicauda</i>   | Other |
| 28971 | Bartlet'S Fairy Basslet | <i>Pseudanthias bartlettorum</i>  | Other |
| 28972 | Bicolor Fairy Basslet   | <i>Pseudanthias bicolor</i>       | Other |
| 28961 | Red-Bar Fairy Basslet   | <i>Pseudanthias cooperi</i>       | Other |
| 28973 | Peach Fairy Basslet     | <i>Pseudanthias dispar</i>        | Other |
| 28979 | Fairy Basslet           | <i>Pseudanthias huchtii</i>       | Other |
| 28974 | Lori'S Anthias          | <i>Pseudanthias lori</i>          | Other |
| 28962 | Purple Queen            | <i>Pseudanthias pascalus</i>      | Other |
| 28963 | Sq-Spot Fairy Basslet   | <i>Pseudanthias pleurotaenia</i>  | Other |



|       |                           |                                       |       |
|-------|---------------------------|---------------------------------------|-------|
| 28975 | Randall'S Fairy Basslet   | <i>Pseudanthias randalli</i>          | Other |
| 28977 | Smithvaniz' Fairy Basslet | <i>Pseudanthias smithvanizi</i>       | Other |
| 28964 | Fairy Basslet             | <i>Pseudanthias sp</i>                | Other |
| 28980 | Fairy Basslet             | <i>Pseudanthias squammipinnis</i>     | Other |
| 28976 | Y Striped Fairy Basslet   | <i>Pseudanthias tuka</i>              | Other |
| 28978 | L-Finned Fairy Basslet    | <i>Pseudanthias ventralis</i>         | Other |
| 5637  | White Ribbon Eel          | <i>Pseudechidna brummeri</i>          | Other |
| 44508 | Ymargin Triggerfish       | <i>Pseudobalistes flavimarginatus</i> | Other |
| 44509 | Blue Triggerfish          | <i>Pseudobalistes fuscus</i>          | Other |
| 29100 | Dottybacks                | <i>Pseudochromidae</i>                | Other |
| 29101 | Surge Dottyback           | <i>Pseudochromis cyanotaenia</i>      | Other |
| 29102 | Dusky Dottyback           | <i>Pseudochromis fuscus</i>           | Other |
| 29103 | Marshall Is Dottyback     | <i>Pseudochromis marshallensis</i>    | Other |
| 29404 | Dottyback                 | <i>Pseudochromis melanotaenia</i>     | Other |
| 29105 | Long-Finned Dottyback     | <i>Pseudochromis polynemus</i>        | Other |
| 29106 | Magenta Dottyback         | <i>Pseudochromis porphyreus</i>       | Other |
| 40638 | Goby                      | <i>Pseudogobius javanicus</i>         | Other |
| 29202 | Soapfish                  | <i>Pseudogramma polyacantha</i>       | Other |
| 29203 | Soapfish                  | <i>Pseudogramma sp.</i>               | Other |
| 29200 | Soapfishes                | <i>Pseudogrammidae</i>                | Other |
| 34501 | Amourhead                 | <i>Pseudopentaceros pectoralis</i>    | Other |
| 29111 | Robust Dottyback          | <i>Pseudoplesiops multisquamatus</i>  | Other |
| 29107 | Revelle'S Basslet         | <i>Pseudoplesiops revellei</i>        | Other |
| 29108 | Rose Island Basslet       | <i>Pseudoplesiops rosae</i>           | Other |
| 29110 | Basslet                   | <i>Pseudoplesiops sp.</i>             | Other |
| 29109 | Hidden Basslet            | <i>Pseudoplesiops typus</i>           | Other |
| 41005 | Blackfin Dartfish         | <i>Ptereleotris evides</i>            | Other |
| 41021 | Filament Dartfish         | <i>Ptereleotris hanae</i>             | Other |
| 41006 | Spot-Tail Dartfish        | <i>Ptereleotris heteroptera</i>       | Other |
| 41009 | Dartfish                  | <i>Ptereleotris lineopinnis</i>       | Other |
| 41007 | Pearly Dartfish           | <i>Ptereleotris microlepis</i>        | Other |
| 41008 | Zebra Dartfish            | <i>Ptereleotris zebra</i>             | Other |
| 32357 | Yellowstreak Fusilier     | <i>Pterocaesio lativittata</i>        | Other |
| 32353 | Twinstripe Fusilier       | <i>Pterocaesio marri</i>              | Other |
| 32360 | Ruddy Fusilier            | <i>Pterocaesio pisang</i>             | Other |

|       |                        |                                       |       |
|-------|------------------------|---------------------------------------|-------|
| 32362 | Mosaic Fusilier        | <i>Pterocaesio tessellata</i>         | Other |
| 32354 | Bluestreak Fusilier    | <i>Pterocaesio tile</i>               | Other |
| 32358 | 3-Striped Fusilier     | <i>Pterocaesio trilineata</i>         | Other |
| 26405 | Spotfin Lionfish       | <i>Pterois antennata</i>              | Other |
| 26406 | Clearfin Lionfish      | <i>Pterois radiata</i>                | Other |
| 26407 | Turkeyfish             | <i>Pterois volitans</i>               | Other |
| 26602 | Ocellated Gurnard      | <i>Pterygiotrigla multiocellata</i>   | Other |
| 26601 | Gurnard                | <i>Pterygiotrigla</i> sp.             | Other |
| 31301 | Slender Suckerfish     | <i>Ptheirichthys lineatus</i>         | Other |
| 34366 | Regal Angelfish        | <i>Pygoplites diacanthus</i>          | Other |
| 28989 | Fairy Basslet          | <i>Rabaulichthys</i> sp.              | Other |
| 45003 | Trunkfish              | <i>Ranzania laevis</i>                | Other |
| 41612 | Mackerel               | <i>Rastrelliger brachysoma</i>        | Other |
| 41610 | Striped Mackerel       | <i>Rastrelliger kanagurta</i>         | Other |
| 40639 | Goby                   | <i>Redigobius bikolanus</i>           | Other |
| 40640 | Goby                   | <i>Redigobius horiae</i>              | Other |
| 40641 | Goby                   | <i>Redigobius sapangus</i>            | Other |
| 31302 | Remora                 | <i>Remora remora</i>                  | Other |
| 30451 | Cardinalfish           | <i>Rhabdamia cypselurus</i>           | Other |
| 30452 | Cardinalfish           | <i>Rhabdamia gracilis</i>             | Other |
| 39234 | Blenny                 | <i>Rhabdoblennius rhabdotrachelus</i> | Other |
| 39250 |                        | <i>Rhabdoblennius ellipes</i>         | Other |
| 39235 | Blenny                 | <i>Rhabdoblennius snowi</i>           | Other |
| 1701  | Guitarfish             | <i>Rhynchobatus djiddensis</i>        | Other |
| 44510 | Picassofish            | <i>Rhinecanthus aculeatus</i>         | Other |
| 44511 | Wedge Picassofish      | <i>Rhinecanthus rectangulus</i>       | Other |
| 44520 | Blackbelly Picassofish | <i>Rhinecanthus verrucosa</i>         | Other |
| 1700  | Guitarfish             | Rhinobatidae                          | Other |
| 5636  | Ribbon Eel             | <i>Rhinomuraena quaesita</i>          | Other |
| 26428 | Weedy Scorpionfish     | <i>Rhinopias frondosa</i>             | Other |
| 31303 | Remora                 | <i>Rhombochirus osteochir</i>         | Other |
| 44607 | Smallnose Boxfish      | <i>Rhynchostracion nasus</i>          | Other |
| 44608 | Largenose Boxfish      | <i>Rhynchostracion rhynorhynchus</i>  | Other |
| 9201  | Telescopefish          | <i>Rosaura indica</i>                 | Other |
| 44569 | Minute Filefish        | <i>Rudarius minutus</i>               | Other |
| 39253 |                        | <i>Salarius alboguttatus</i>          | Other |
| 39236 | Spotted Rock Blenny    | <i>Salarius fasciatus</i>             | Other |
| 39255 | Blenny                 | <i>Salarius luctuosus</i>             | Other |

|       |                             |                                  |       |
|-------|-----------------------------|----------------------------------|-------|
| 39254 | Blenny                      | <i>Salarius segmentatus</i>      | Other |
| 44000 | Righteye Flounders          | Samaridae                        | Other |
| 44001 | 3 Spot Flounder             | <i>Samariscus triocellatus</i>   | Other |
| 16001 | Graceful Lizardfish         | <i>Saurida gracilis</i>          | Other |
| 16002 | Nebulous Lizardfish         | <i>Saurida nebulosa</i>          | Other |
| 34100 | Scats                       | Scatophagidae                    | Other |
| 34101 | Scat                        | <i>Scatophagus argus</i>         | Other |
| 40101 | Schindleriid                | <i>Schindleria praematurus</i>   | Other |
| 40100 | Shindleriid                 | Schindleriidae                   | Other |
| 6616  | Snake Eel                   | <i>Schismorhinchus labialis</i>  | Other |
| 6617  | Snake Eel                   | <i>Schultzidia johnstonensis</i> | Other |
| 6618  | Snake Eel                   | <i>Schultzidia retropinnis</i>   | Other |
| 32404 | Spinecheek                  | <i>Scolopsis affinis</i>         | Other |
| 32402 | 2 Line Spinecheek           | <i>Scolopsis bilineatus</i>      | Other |
| 32406 | Ciliate Spinecheek          | <i>Scolopsis ciliatus</i>        | Other |
| 32401 | Bl And Wh Spinecheek        | <i>Scolopsis lineatus</i>        | Other |
| 32403 | Margarite'S Spinecheek      | <i>Scolopsis margaritifer</i>    | Other |
| 32407 | Spinecheek                  | <i>Scolopsis taeniopterus</i>    | Other |
| 32405 | 3 Line Spinecheek           | <i>Scolopsis trilineatus</i>     | Other |
| 32408 | Spinecheek                  | <i>Scolopsis xenochrous</i>      | Other |
| 41611 | Narrow-Barred King Mackerel | <i>Scomberomorus commerson</i>   | Other |
| 26400 | Scorpionfish                | Scorpaenidae                     | Other |
| 26413 | Guam Scorpionfish           | <i>Scorpaenodes guamensis</i>    | Other |
| 26429 | Hairy Scorpionfish          | <i>Scorpaenodes hirsutus</i>     | Other |
| 26414 | Kellogg'S Scorpionfish      | <i>Scorpaenodes kelloggi</i>     | Other |
| 26412 | Minor Scorpionfish          | <i>Scorpaenodes minor</i>        | Other |
| 26415 | Coral Scorpionfish          | <i>Scorpaenodes parvipinnis</i>  | Other |
| 26420 | Blotchfin Scorpionfish      | <i>Scorpaenodes varipinis</i>    | Other |
| 26417 | Devil Scorpionfish          | <i>Scorpaenopsis diabolus</i>    | Other |
| 26421 | Pygmy Scorpionfish          | <i>Scorpaenopsis fowleri</i>     | Other |
| 26422 | Flasher Scorpionfish        | <i>Scorpaenopsis macrochir</i>   | Other |
| 26416 | Tassled Scorpionfish        | <i>Scorpaenopsis oxycephala</i>  | Other |
| 26434 | Papuan Scorpionfish         | <i>Scorpaenopsis papuensis</i>   | Other |
| 26432 | Scorpionfish                | <i>Scorpaenopsis sp.</i>         | Other |
| 5654  | Tiger Snake Moray           | <i>Scuticaria tigrinis</i>       | Other |
| 26408 | Yellowspotted Scorpionfish  | <i>Sebastapistes cyanostigma</i> | Other |
| 26409 | Galactacma Scorpionfish     | <i>Sebastapistes galactacma</i>  | Other |
| 26410 | Mauritius Scorpionfish      | <i>Sebastapistes mauritiana</i>  | Other |
| 26425 | Barchin Scorpionfish        | <i>Sebastapistes strongia</i>    | Other |

|       |                       |   |       |
|-------|-----------------------|---|-------|
| 31807 | Pugnose Soapy         | <i>Secutor ruconius</i>                 | Other |
| 28970 | Basslet               | <i>Selenanthias myersi</i>              | Other |
| 28988 | Hawkfish Anthias      | <i>Serranocirrhitus latus</i>           | Other |
| 40645 | Goby                  | <i>Sicyopterus<br/>macrostetholepis</i> | Other |
| 40646 | Goby                  | <i>Sicyopterus micrurus</i>             | Other |
| 40647 | Goby                  | <i>Sicyopterus</i> sp.                  | Other |
| 40642 | Goby                  | <i>Sicyopus leprurus</i>                | Other |
| 40644 | Goby                  | <i>Sicyopus</i> sp.                     | Other |
| 40643 | Goby                  | <i>Sicyopus zosterophorum</i>           | Other |
| 5615  | Peppered Moray        | <i>Sideria picta</i>                    | Other |
| 5617  | White-Eyed Moray      | <i>Sideria prosopeion</i>               | Other |
| 40530 | Goby                  | <i>Signigobius biocellatus</i>          | Other |
| 40531 | Goby                  | <i>Silhouettea</i> sp.                  | Other |
| 30700 | Sillagos              | Sillaginidae                            | Other |
| 30701 | Cardinalfish          | <i>Sillago sihama</i>                   | Other |
| 30431 | Cardinalfish          | <i>Siphamia fistulosa</i>               | Other |
| 30459 | Cardinalfish          | <i>Siphamia fuscolineata</i>            | Other |
| 30430 | Cardinalfish          | <i>Siphamia versicolor</i>              | Other |
| 44101 | Banded Sole           | <i>Soleichthys heterohinos</i>          | Other |
| 44100 | Soles                 | Soleidae                                | Other |
| 25700 | Ghost Pipefish        | <i>Solenostomidae</i>                   | Other |
| 25701 | Ghost Pipefish        | <i>Solenostomus cyanopterus</i>         | Other |
| 25702 | Ornate Ghost Pipefish | <i>Solenostomus paradoxus</i>           | Other |
| 27305 | Flathead              | <i>Sorsogona welanderi</i>              | Other |
| 30434 | Cardinalfish          | <i>Sphaeramia nematoptera</i>           | Other |
| 30432 | Cardinalfish          | <i>Sphaeramia orbicularis</i>           | Other |
| 36004 | Sharpfin Barracuda    | <i>Sphyraena acutipinnis</i>            | Other |
| 36001 | Great Barracuda       | <i>Sphyraena barracuda</i>              | Other |
| 36008 | Yellowtail Barracuda  | <i>Sphyraena flavicauda</i>             | Other |
| 36003 | Blackspot Barracuda   | <i>Sphyraena forsteri</i>               | Other |
| 36007 | Arrow Barracuda       | <i>Sphyraena novaehollandiae</i>        | Other |
| 36002 | Pygmy Barracuda       | <i>Sphyraena obtusata</i>               | Other |
| 36006 | Slender Barracuda     | <i>Sphyraena putnamiae</i>              | Other |
| 36005 | Blackfin Barracuda    | <i>Sphyraena qenie</i>                  | Other |
| 36000 | Barracudas            | Sphyraenidae                            | Other |
| 4301  | Blue Sprat            | <i>Spratelloides delicatulus</i>        | Other |
| 4305  | Silver Sprat          | <i>Spratelloides gracilis</i>           | Other |
| 39237 | Blenny                | <i>Stanulus seychellensis</i>           | Other |
| 35043 | White-Bar Gregory     | <i>Stegastes albifasciatus</i>          | Other |

|       |                          |                                  |       |
|-------|--------------------------|----------------------------------|-------|
| 35044 | Pacific Gregory          | <i>Stegastes fasciolatus</i>     | Other |
| 35045 | Farmerfish               | <i>Stegastes lividus</i>         | Other |
| 35046 | Dusky Farmerfish         | <i>Stegastes nigricans</i>       | Other |
| 702   | Leopard Shark            | <i>Stegastoma varium</i>         | Other |
| 21809 | Panatella Silverside     | <i>Stenatherina panatella</i>    | Other |
| 40648 | Goby                     | <i>Stenogobius genivittatus</i>  | Other |
| 40649 | Goby                     | <i>Stenogobius sp.</i>           | Other |
| 8900  | Hatchetfishes            | Sternoptichidae                  | Other |
| 40650 | Goby                     | <i>Stiphodon elegans</i>         | Other |
| 40651 | Goby                     | <i>Stiphodon sp.</i>             | Other |
| 4408  | Samoa Anchovy            | <i>Stolephorus apiensis</i>      | Other |
| 4404  | Indian Anchovy           | <i>Stolephorus indicus</i>       | Other |
| 4407  | Gold Esurine Anchovy     | <i>Stolephorus insularis</i>     | Other |
| 4409  | Caroline Islands Anchovy | <i>Stolephorus multibranchus</i> | Other |
| 4403  | West Pacific Anchovy     | <i>Stolephorus pacificus</i>     | Other |
| 4499  | Anchovy                  | <i>Stolephorus sp.</i>           | Other |
| 20703 | Reef Needlefish          | <i>Strongylura incisa</i>        | Other |
| 20705 | Littoral Needlefish      | <i>Strongylura leiura leiura</i> | Other |
| 5638  | Giant Esturine Moray     | <i>Strophidon sathete</i>        | Other |
| 44512 | Scythe Triggerfish       | <i>Sufflamen bursa</i>           | Other |
| 44513 | Halfmoon Triggerfish     | <i>Sufflamen chrysoptera</i>     | Other |
| 44514 | Bridle Triggerfish       | <i>Sufflamen freanatus</i>       | Other |
| 32371 | Symphysanid              | <i>Symphysanodon typus</i>       | Other |
| 32370 | Sympysanodon             | Symphysanodontidae               | Other |
| 26418 | Stonefish                | <i>Synanceia verrucosa</i>       | Other |
| 5700  | Cutthroat Eel            | Synaphobranchidae                | Other |
| 5701  | Cutthroat Eel            | <i>Synaphobranchus sp.</i>       | Other |
| 43504 | Circlled Dragonet        | <i>Synchiropus circularis</i>    | Other |
| 43511 | Ladd'S Dragonet          | <i>Synchiropus laddi</i>         | Other |
| 45308 | Morrison'S Dragonet      | <i>Synchiropus morrisoni</i>     | Other |
| 43505 | Ocellated Dragonet       | <i>Synchiropus ocellatus</i>     | Other |
| 43510 | Dragonet                 | <i>Synchiropus sp.</i>           | Other |
| 43506 | Mandarin Fish            | <i>Synchiropus splendidus</i>    | Other |
| 43509 | Pipefish, Seahorse       | Syngnathidae                     | Other |
| 25800 | Alligator Pipefish       | <i>Syngnathoides biaculeatus</i> | Other |
| 25815 | Lizardfish               | Synodontidae                     | Other |
| 16000 | 2-Spot Lizardfish        | <i>Synodus binotatus</i>         | Other |
| 16003 | Clearfin Lizardfish      | <i>Synodus dermatogenys</i>      | Other |
| 16007 | Reef Lizardfish          | <i>Synodus englemanni</i>        | Other |

|       |                         |  |       |
|-------|-------------------------|--|-------|
| 16004 | Blackblotch Lizardfish  | <i>Synodus jaculum</i>                 | Other |
| 16005 | Variegatus Lizardfish   | <i>Synodus variegatus</i>              | Other |
| 16006 | Leaf Fish               | <i>Taenianotus triacanthus</i>         | Other |
| 26419 | Goby                    | <i>Taenioides limicola</i>             | Other |
| 40652 | Giant Reef Ray          | <i>Taeniura meyeni</i>                 | Other |
| 2002  | Crescent-Banded Grunter | <i>Terapon jarbua</i>                  | Other |
| 29901 | Thornfishes             | Teraponidae                            | Other |
| 29900 | Smooth Puffers          | Tetraodontidae                         | Other |
| 26451 | Mangrove Waspfish       | <i>Tetraroge barbata</i>               | Other |
| 26450 | Waspfishes              | Tetrarogidae                           | Other |
| 4402  | Little Priest           | <i>Thryssa baelama</i>                 | Other |
| 27302 | Broadhead Flathead      | <i>Thysanophrys arenicola</i>          | Other |
| 27303 | Longsnout Flathead      | <i>Thysanophrys chiltonae</i>          | Other |
| 27301 | Fringlip Flathead       | <i>Thysanophrys otaitensis</i>         | Other |
| 34602 | Tilapia                 | <i>Tilapia zillii</i>                  | Other |
| 33701 | Banded Archerfish       | <i>Toxotes jaculator</i>               | Other |
| 33700 | Archerfishes            | Toxotidae                              | Other |
| 25816 | Double-Ended Pipefish   | <i>Trachyrampus bicoarctata</i>        | Other |
| 44300 | Spikefishes             | Triacanthodidae                        | Other |
| 1108  | Reef Whitetip Shark     | <i>Triaenodon obesus</i>               | Other |
| 37200 | Sand Divers             | Trichonotidae                          | Other |
| 37201 | Micronesian Sand-Diver  | <i>Trichonotus sp</i>                  | Other |
| 26600 | Gurnards                | Triglidae                              | Other |
| 40653 | Goby                    | <i>Trimma caesiura</i>                 | Other |
| 40654 | Goby                    | <i>Trimma naudei</i>                   | Other |
| 40655 | Goby                    | <i>Trimma okinawae</i>                 | Other |
| 40658 | Goby                    | <i>Trimma sp. A</i>                    | Other |
| 40659 | Goby                    | <i>Trimma sp. B</i>                    | Other |
| 40656 | Goby                    | <i>Trimma taylori</i>                  | Other |
| 40657 | Goby                    | <i>Trimma tevegae</i>                  | Other |
| 40660 | Goby                    | <i>Trimmatom eviotops</i>              | Other |
| 44702 | 3 Tooth Puffer          | <i>Triodon bursarius</i>               | Other |
| 44701 | 3 Tooth Puffer          | <i>Triodon macropterus</i>             | Other |
| 44700 | Tripletooth Puffers     | Triodontidae                           | Other |
| 39000 | Triplefins              | Tripterygiidae                         | Other |
| 20706 | Keeled Houndfish        | <i>Tylosurus acus melanotus</i>        | Other |
| 20704 | Houndfish               | <i>Tylosurus crocodilis crocodilis</i> | Other |
| 39009 | Longjaw Triplefin       | <i>Ucla xenogrammus</i>                | Other |
| 37800 | Stargazers              | Uranoscopidae                          | Other |

|       |                         |                                      |                          |
|-------|-------------------------|--------------------------------------|--------------------------|
| 37801 | Stargazer               | <i>Uranoscopus</i> sp.               | Other                    |
| 2004  | Porcupine Ray           | <i>Urogymnus africanus</i>           | Other                    |
| 5639  | Unicolor Snake Moray    | <i>Uropterygius concolor</i>         | Other                    |
| 5660  | Fiji Moray Eel          | <i>Uropterygius fijiensis</i>        | Other                    |
| 5650  | Brown-Spotted Snake Eel | <i>Uropterygius fuscoguttatus</i>    | Other                    |
| 5651  | Gosline'S Snake Moray   | <i>Uropterygius goslinei</i>         | Other                    |
| 5652  | Moon Moray              | <i>Uropterygius kamar</i>            | Other                    |
| 5642  | Lg-Headed Snake Moray   | <i>Uropterygius macrocephalus</i>    | Other                    |
| 5640  | Marbled Snake Moray     | <i>Uropterygius marmoratus</i>       | Other                    |
| 5641  | Tidepool Snake Moray    | <i>Uropterygius micropterus</i>      | Other                    |
| 5653  | Lg-Spotted Snake Moray  | <i>Uropterygius polyspilus</i>       | Other                    |
| 5643  | Moray Eel               | <i>Uropterygius supraforatus</i>     | Other                    |
| 5644  | Moray Eel               | <i>Uropterygius xanthopterus</i>     | Other                    |
| 2008  | Roundray                | <i>Urotrygon daviesi</i>             | Other                    |
| 40532 | Glass Goby              | <i>Valenciennesa muralis</i>         | Other                    |
| 40663 | Parva Goby              | <i>Valenciennesa parva</i>           | Other                    |
| 40533 | Goby                    | <i>Valenciennesa puellaris</i>       | Other                    |
| 40534 | Goby                    | <i>Valenciennesa sexguttatus</i>     | Other                    |
| 40536 | Goby                    | <i>Valenciennesa</i> sp.             | Other                    |
| 40535 | Goby                    | <i>Valenciennesa strigatus</i>       | Other                    |
| 40521 | Goby                    | <i>Vanderhorstia ambanoro</i>        | Other                    |
| 40661 | Goby                    | <i>Vanderhorstia lanceolata</i>      | Other                    |
| 40522 | Goby                    | <i>Vanderhorstia ornatissima</i>     | Other                    |
| 44515 | Guided Triggerfish      | <i>Xanthichthys auromarginatus</i>   | Other                    |
| 44516 | Bluelined Triggerfish   | <i>Xanthichthys careuleolineatus</i> | Other                    |
| 44521 | Crosshatch Triggerfish  | <i>Xanthichthys mento</i>            | Other                    |
| 40713 | Wriggler                | <i>Xenisthmus</i> sp.                | Other                    |
| 40710 | Flathead Wriggler       | Xenisthmidae                         | Other                    |
| 40712 | Barred Wriggler         | <i>Xenisthmus polyzonatus</i>        | Other                    |
| 44517 | Triggerfish             | <i>Xenobalistes tumidipectoris</i>   | Other                    |
| 39238 | Blenny                  | <i>Xiphasia matsubarai</i>           | Other                    |
| 41250 | Moorish Idols           | Zanclidae                            | Other                    |
| 41251 | Moorish Idol            | <i>Zanclus cornutus</i>              | Other                    |
| 20756 | Esturine Halfbeak       | <i>Zenarchopterus dispar</i>         | Other                    |
| 49400 | ASSORTED REEF FISH      | Misc. Reeffish                       | Misc. Reeffish           |
| 49110 | SHALLOW BOTTOMFISH      | Misc. Shallow bottomfish             | Misc. Shallow bottomfish |

|       |                        |  |                     |
|-------|------------------------|--|---------------------|
| 49100 | ASSORTED<br>BOTTOMFISH | Misc. Bottomfish                       | Misc. Bottomfish    |
| 72600 | Crown-Of-Thorns        | <i>Acanthaster planci</i>              | Other Invertebrates |
| 79301 | Stonefish              | <i>Actinopyga lecanora</i>             | Other Invertebrates |
| 79302 | Blackfish              | <i>Actinopyga miliaris</i>             | Other Invertebrates |
| 79303 | Sea Cucumber           | <i>Actinopyga obesa</i>                | Other Invertebrates |
| 79304 | Sea Cucumber           | <i>Actinopyga sp</i>                   | Other Invertebrates |
| 72500 | Starfish               | Asterinidae                            | Other Invertebrates |
| 72400 | Starfish               | Asteropidae                            | Other Invertebrates |
| 72100 | Starfish               | Astropectinidae                        | Other Invertebrates |
| 79801 | Sea Cucumber           | <i>Bohadschia argus</i>                | Other Invertebrates |
| 79802 | Sea Cucumber           | <i>Bohadschia graeffei</i>             | Other Invertebrates |
| 79803 | Brown Sandfish         | <i>Bohadschia marmorata</i>            | Other Invertebrates |
| 79804 | Sea Cucumber           | <i>Bohadschia paradoxa</i>             | Other Invertebrates |
| 79805 | Sea Cucumber           | <i>Bohadschia sp.</i>                  | Other Invertebrates |
| 78900 | Irregular Urchins      | Brissidae                              | Other Invertebrates |
| 97100 | Jellyfish              | <i>Cephea sp.</i>                      | Other Invertebrates |
| 78100 | Cidarials              | Cidaridae                              | Other Invertebrates |
| 71000 | Crinoids               | Class Crinoidea                        | Other Invertebrates |
| 78000 | Sea Urchins            | Class Echinoidea                       | Other Invertebrates |
| 78800 |                        | Clypeasteridae                         | Other Invertebrates |
| 79400 | Sea Cucumbers          | Cucumariidae                           | Other Invertebrates |
| 78301 | Longspine Urchin       | <i>Diadema savignyi</i>                | Other Invertebrates |
| 78302 | Longspine Urchin       | <i>Diadema setosum</i>                 | Other Invertebrates |
| 78300 | Sea Urchins            | Diadematidae                           | Other Invertebrates |
| 78700 | Sea Urchins            | Echinoidea                             | Other Invertebrates |
| 78600 | Sea Urchins            | Echinometridae                         | Other Invertebrates |
| 72800 | Reef Starfish          | Echinosteridae                         | Other Invertebrates |
| 78304 | Longspine Urchin       | <i>Echinothrix calamaris</i>           | Other Invertebrates |
| 78303 | Longspine Urchin       | <i>Echinothrix diadema</i>             | Other Invertebrates |
| 78200 | Sea Urchins            | Echinothuriidae                        | Other Invertebrates |
| 78605 | Slate Pencil Urchin    | <i>Heterocentrotus<br/>mammillatus</i> | Other Invertebrates |
| 79201 | Lollyfish              | <i>Holothuria atra</i>                 | Other Invertebrates |
| 79202 | Pinkfish               | <i>Holothuria edulis</i>               | Other Invertebrates |
| 79203 | White Teatfish         | <i>Holothuria fuscogilva</i>           | Other Invertebrates |
| 79204 | Elephant'S Trunkfish   | <i>Holothuria fuscopunctata</i>        | Other Invertebrates |
| 79205 | Sea Cucumber           | <i>Holothuria hilla</i>                | Other Invertebrates |
| 79206 | Sea Cucumber           | <i>Holothuria impatiens</i>            | Other Invertebrates |
| 79207 | Sea Cucumber           | <i>Holothuria leucospilota</i>         | Other Invertebrates |



|       |                               |                               |                     |
|-------|-------------------------------|-------------------------------|---------------------|
| 79208 | Sea Cucumber                  | <i>Holothuria</i> sp          | Other Invertebrates |
| 79200 | Sea Cucumber                  | Holothuriidae                 | Other Invertebrates |
| 79000 | Sea Cucumbers                 | Holothuroidea                 | Other Invertebrates |
| 72700 | Spiney-Armed Starfish         | <i>Mithrodia bradleyi</i>     | Other Invertebrates |
| 72300 | Orange Starfish               | <i>Ophidiaster confertus</i>  | Other Invertebrates |
| 72200 | Starfish                      | Oreasteridae                  | Other Invertebrates |
| 79500 | Sea Cucumbers                 | Phyllophoridae                | Other Invertebrates |
| 78503 | Common Urchin                 | <i>Pseudoboletia maculata</i> | Other Invertebrates |
| 72000 | Starfish                      | Asteroidea                    | Other Invertebrates |
| 75000 | Basket, Brittle, Serpentstars | Ophiuroidea                   | Other Invertebrates |
| 72900 | Starfish                      | Sphaerasteridae               | Other Invertebrates |
| 79100 | Sea Cucumbers                 | Stichopodidae                 | Other Invertebrates |
| 79101 | Greenfish                     | <i>Stichopus chloronotus</i>  | Other Invertebrates |
| 79102 | Sea Cucumber                  | <i>Stichopus horrens</i>      | Other Invertebrates |
| 79103 | Sea Cucumber                  | <i>Stichopus noctivatus</i>   | Other Invertebrates |
| 79105 | Sea Cucumber                  | <i>Stichopus</i> sp.          | Other Invertebrates |
| 79104 | Curryfish                     | <i>Stichopus variegatus</i>   | Other Invertebrates |
| 79601 | Sea Cucumber                  | <i>Synapta maculata</i>       | Other Invertebrates |
| 79602 | Sea Cucumber                  | <i>Synapta media</i>          | Other Invertebrates |
| 79603 | Sea Cucumber                  | <i>Synapta</i> sp.            | Other Invertebrates |
| 79600 | Sea Cucumbers                 | Synaptidae                    | Other Invertebrates |
| 78400 | Sea Urchins                   | Temnopleuridae                | Other Invertebrates |
| 79901 | Prickly Redfish               | <i>Thelenota ananas</i>       | Other Invertebrates |
| 79902 | Amberfish                     | <i>Thelenotaanax</i>          | Other Invertebrates |
| 79903 | Sea Cucumber                  | <i>Thelenota</i> sp.          | Other Invertebrates |
| 78502 | Flower Urchin                 | <i>Toxopneustes pileolus</i>  | Other Invertebrates |
| 78500 | Shortspine Urchins            | Toxopneustidae                | Other Invertebrates |
| 78501 | Shortspine Urchin             | <i>Tripneustes gratilla</i>   | Other Invertebrates |

## APPENDIX B: LIST OF PROTECTED SPECIES AND DESIGNATED CRITICAL HABITAT

Table B-1. Protected species found or reasonably believed to be found near or in Hawai'i shallow-set longline waters.

| Common name             | Scientific name   | ESA listing status | MMPA status | Occurrence                   | References                                |
|-------------------------|---|--------------------|-------------|------------------------------|---|
| <b>Seabirds</b>         |   |                    |             |                              |   |
| Laysan Albatross        | <i>Phoebastria immutabilis</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Black-Footed Albatross  | <i>Phoebastria nigripes</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Short-Tailed Albatross  | <i>Phoebastria albatrus</i>   | Endangered         | N/A         | Breeding visitor in the NWHI | 35 FR 8495, 65 FR 46643, Pyle & Pyle 2009 |
| Northern Fulmar         | <i>Fulmarus glacialis</i>   | Not Listed         | N/A         | Winter resident              | Pyle & Pyle 2009                          |
| Kermadec Petrel         | <i>Pterodroma neglecta</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Herald Petrel           | <i>Pterodroma arminjoniana</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Murphy's Petrel         | <i>Pterodroma ultima</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Mottled Petrel          | <i>Pterodroma inexpectata</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Juan Fernandez Petrel   | <i>Pterodroma externa</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Hawaiian Petrel         | <i>Pterodroma sandwichensis</i><br>( <i>Pterodroma phaeopygia sandwichensis</i> ) | Endangered         | N/A         | Breeding visitor in the MHI  | 32 FR 4001, Pyle & Pyle 2009              |
| White-Necked Petrel     | <i>Pterodroma cervicalis</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Bonin Petrel            | <i>Pterodroma hypoleuca</i>   | Not Listed         | N/A         | Breeding visitor in the NWHI | Pyle & Pyle 2009                          |
| Black-Winged Petrel     | <i>Pterodroma nigripennis</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Cook Petrel             | <i>Pterodroma cookii</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Stejneger Petrel        | <i>Pterodroma longirostris</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Pycroft Petrel          | <i>Pterodroma pycrofti</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Bulwer Petrel           | <i>Bulweria bulwerii</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Flesh-Footed Shearwater | <i>Ardenna carneipes</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Wedge-Tailed Shearwater | <i>Ardenna pacifica</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Buller's Shearwater     | <i>Ardenna bulleri</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |

| Common name              | Scientific name  | ESA listing status | MMPA status | Occurrence                   | References                       |
|--------------------------|--|--------------------|-------------|------------------------------|----------------------------------|
| Sooty Shearwater         | <i>Ardenna grisea</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                 |
| Short-Tailed Shearwater  | <i>Ardenna tenuirostris</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                 |
| Christmas Shearwater     | <i>Puffinus nativitatis</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Newell's Shearwater      | <i>Puffinus newelli</i><br>( <i>Puffinus auricularis newelli</i> ) | Threatened         | N/A         | Breeding visitor             | 40 FR 44149,<br>Pyle & Pyle 2009 |
| Wilson's Storm-Petrel    | <i>Oceanites oceanicus</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                 |
| Leach's Storm-Petrel     | <i>Oceanodroma leucorhoa</i>                                       | Not Listed         | N/A         | Winter resident              | Pyle & Pyle 2009                 |
| Band-Rumped Storm-Petrel | <i>Oceanodroma castro</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Tristram Storm-Petrel    | <i>Oceanodroma tristrami</i>                                       | Not Listed         | N/A         | Breeding visitor in the NWHI | Pyle & Pyle 2009                 |
| White-Tailed Tropicbird  | <i>Phaethon lepturus</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Red-Tailed Tropicbird    | <i>Phaethon rubricauda</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Masked Booby             | <i>Sula dactylatra</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Brown Booby              | <i>Sula leucogaster</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Red-Footed Booby         | <i>Sula sula</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Great Frigatebird        | <i>Fregata minor</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Lesser Frigatebird       | <i>Fregata ariel</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Laughing Gull            | <i>Leucophaeus atricilla</i>                                       | Not Listed         | N/A         | Winter resident in the MHI   | Pyle & Pyle 2009                 |
| Franklin Gull            | <i>Leucophaeus pipixcan</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                 |
| Ring-Billed Gull         | <i>Larus delawarensis</i>  | Not Listed         | N/A         | Winter resident in the MHI   | Pyle & Pyle 2009                 |
| Herring Gull             | <i>Larus argentatus</i>  | Not Listed         | N/A         | Winter resident in the NWHI  | Pyle & Pyle 2009                 |
| Slaty-Backed Gull        | <i>Larus schistisagus</i>  | Not Listed         | N/A         | Winter resident in the NWHI  | Pyle & Pyle 2009                 |
| Glaucous-Winged Gull     | <i>Larus glaucescens</i>   | Not Listed         | N/A         | Winter resident              | Pyle & Pyle 2009                 |
| Brown Noddy              | <i>Anous stolidus</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Black Noddy              | <i>Anous minutus</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |
| Blue-Gray Noddy          | <i>Procelsterna cerulea</i>  | Not Listed         | N/A         | Breeding visitor in the NWHI | Pyle & Pyle 2009                 |
| White Tern               | <i>Gygis alba</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                 |

| Common name            | Scientific name                 | ESA listing status                     | MMPA status | Occurrence  | References  |
|------------------------|---------------------------------|--|-------------|---|---|
| Sooty Tern             | <i>Onychoprion fuscatus</i>     | Not Listed                             | N/A         | Breeding visitor  | Pyle & Pyle 2009  |
| Gray-Backed Tern       | <i>Onychoprion lunatus</i>      | Not Listed                             | N/A         | Breeding visitor  | Pyle & Pyle 2009  |
| Little Tern            | <i>Sternula albifrons</i>       | Not Listed                             | N/A         | Breeding visitor in the NWHI  | Pyle & Pyle 2009  |
| Least Tern             | <i>Sternula antillarum</i>      | Not Listed                             | N/A         | Breeding visitor in the NWHI  | Pyle & Pyle 2009  |
| Arctic Tern            | <i>Sterna paradisaea</i>        | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| South Polar Skua       | <i>Stercorarius maccormicki</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| Pomarine Jaeger        | <i>Stercorarius pomarinus</i>   | Not Listed                             | N/A         | Winter resident in the MHI  | Pyle & Pyle 2009  |
| Parasitic Jaeger       | <i>Stercorarius parasiticus</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| Long-Tailed Jaeger     | <i>Stercorarius longicaudus</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| <b>Sea turtles</b>     |                                 |  |             |   |   |
| Green Sea Turtle       | <i>Chelonia mydas</i>           | Threatened (Central North Pacific DPS) | N/A         | Most common turtle in the Hawaiian Islands, much more common in nearshore state waters (foraging grounds) than offshore federal waters. Most nesting occurs on French Frigate Shoals in the NWHI. Foraging and haul out in the MHI.       | 43 FR 32800, 81 FR 20057, Balazs et al. 1992, Kolinski et al. 2001            |
| Green Sea Turtle       | <i>Chelonia mydas</i>           | Threatened (East Pacific DPS)          | N/A         | Nest primarily in Mexico and the Galapagos Islands. Little known about their pelagic range west of 90°W, but may range as far as the Marshall Islands. Genetic testing confirmed that they are incidentally taken in the HI DSLL fishery. | 43 FR 32800, 81 FR 20057, WPRFMC 2009, Clifton et al. 1982, Karl & Bowen 1999 |
| Hawksbill Sea Turtle   | <i>Eretmochelys imbricata</i>   | Endangered <sup>a</sup>                | N/A         | Small population foraging around Hawai'i and low level nesting on Maui and Hawai'i Islands. Occur worldwide in tropical and subtropical waters.   | 35 FR 8491, NMFS & USFWS 2007, Balazs et al. 1992, Katahira et al. 1994       |
| Leatherback Sea Turtle | <i>Dermochelys coriacea</i>     | Endangered <sup>a</sup>                | N/A         | Regularly sighted in offshore waters, especially at the southeastern end of the archipelago.  | 35 FR 8491, NMFS & USFWS 1997   |

| Common name               | Scientific name                | ESA listing status  | MMPA status   | Occurrence   | References  |
|---------------------------|--------------------------------|---|---------------|--|---|
| Loggerhead Sea Turtle     | <i>Caretta caretta</i>         | Endangered (North Pacific DPS)  | N/A           | Rare in Hawai'i. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters.                        | 43 FR 32800, 76 FR 58868, Dodd 1990, Balazs 1979  |
| Olive Ridley Sea Turtle   | <i>Lepidochelys olivacea</i>   | Threatened (Entire species, except for the breeding population on the Pacific coast of Mexico, which is listed as endangered) | N/A           | Rare in Hawai'i. Occurs worldwide in tropical and warm temperate ocean waters.   | 43 FR 32800, Pitman 1990, Balacz 1982   |
| <b>Marine mammals</b>     |                                |   |               |  |   |
| Blainville's Beaked Whale | <i>Mesoplodon densirostris</i> | Not Listed  | Non-strategic | Found worldwide in tropical and temperate waters   | Mead 1989   |
| Blue Whale                | <i>Balaenoptera musculus</i>   | Endangered  | Strategic     | Acoustically recorded off of Oahu and Midway Atoll, small number of sightings around Hawai'i. Considered extremely rare, generally occur in winter and summer. | 35 FR 18319, Bradford et al. 2013, Northrop et al. 1971, Thompson & Friedl 1982, Stafford et al. 2001 |
| Bottlenose Dolphin        | <i>Tursiops truncatus</i>      | Not Listed  | Non-strategic | Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.   | Perrin et al. 2009, Martien et al. 2012   |
| Bryde's Whale             | <i>Balaenoptera edeni</i>      | Not Listed  | Unknown       | Distributed widely across tropical and warm-temperate Pacific Ocean.   | Leatherwood et al. 1982   |
| Common Dolphin            | <i>Delphinus delphis</i>       | Not Listed  | N/A           | Found worldwide in temperate and subtropical seas.   | Perrin et al. 2009  |
| Cuvier's Beaked Whale     | <i>Ziphius cavirostris</i>     | Not Listed  | Non-strategic | Occur year round in Hawaiian waters.   | McSweeney et al. 2007   |
| Dall's Porpoise           | <i>Phocoenoides dalli</i>      | Not Listed  | Non-strategic | Range across the entire north Pacific Ocean.   | Hall 1979   |
| Dwarf Sperm Whale         | <i>Kogia sima</i>              | Not Listed  | Non-strategic | Most common in waters between 500 m and 1,000 m in depth. Found worldwide in tropical and warm-temperate waters.   | Nagorsen 1985, Baird et al. 2013  |

| Common name            | Scientific name                  | ESA listing status                     | MMPA status   | Occurrence   | References   |
|------------------------|----------------------------------|--|---------------|--|--|
| False Killer Whale     | <i>Pseudorca crassidens</i>      | Not Listed                             | Non-strategic | Found worldwide in tropical and warm-temperate waters. Pelagic stock tracked to within 11 km of Hawaiian islands.  | Stacey et al. 1994, Baird et al. 2012, Bradford et al. 2015  |
| Fin Whale              | <i>Balaenoptera physalus</i>     | Endangered                             | Strategic     | Infrequent sightings in Hawai'i waters. Considered rare in Hawai'i, though may migrate into Hawaiian waters during fall/winter based on acoustic recordings. | 35 FR 18319, Hamilton et al. 2009, Thompson & Friedl 1982  |
| Fraser's Dolphin       | <i>Lagenodelphis hosei</i>       | Not Listed                             | Non-strategic | Found worldwide in tropical waters.  | Perrin et al. 2009   |
| Guadalupe Fur Seal     | <i>Arctocephalus townsendi</i>   | Threatened                             | Strategic     | Extremely rare sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.   | 50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987   |
| Hawaiian Monk Seal     | <i>Neomonachus schauinslandi</i> | Endangered <sup>a</sup>                | Strategic     | Endemic tropical seal. Occurs throughout the archipelago. MHI population spends some time foraging in federal waters during the day.                         | 41 FR 51611, Baker et al. 2011   |
| Humpback Whale         | <i>Megaptera novaeangliae</i>    | Delisted Due to Recovery (Hawai'i DPS) | Strategic     | Migrate through the archipelago and breed during the winter. Common during winter months, when they are generally found within the 100 m isobath.            | 35 FR 18319, 81 FR 62259, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978 |
| Killer Whale           | <i>Orcinus orca</i>              | Not Listed                             | Non-strategic | Rare in Hawai'i. Prefer colder waters within 800 km of continents.   | Mitchell 1975, Baird et al. 2006   |
| Longman's Beaked Whale | <i>Indopacetus pacificus</i>     | Not Listed                             | Non-strategic | Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. Rare in Hawai'i.                         | Dalebout 2003, Baird et al. 2013   |
| Melon-Headed Whale     | <i>Peponocephala electra</i>     | Not Listed                             | Non-strategic | Found in tropical and warm-temperate waters worldwide, found primarily in equatorial waters. Uncommon in Hawai'i.  | Perryman et al. 1994, Barlow 2006, Bradford et al. 2013  |

| Common name                 | Scientific name                     | ESA listing status      | MMPA status   | Occurrence   | References   |
|-----------------------------|-------------------------------------|-------------------------|---------------|--|--|
| Minke Whale                 | <i>Balaenoptera acutorostrata</i>   | Not Listed              | Non-strategic | Occur seasonally around Hawai i  | Barlow 2003, Rankin & Barlow 2005  |
| North Pacific Right Whale   | <i>Eubalaena japonica</i>           | Endangered <sup>a</sup> | Strategic     | Extremely rare in Hawai i waters   | 35 FR 18319, 73 FR 12024, Rowntree et al. 1980, Herman et al. 1980       |
| Northern Elephant Seal      | <i>Mirounga angustirostris</i>      | Not Listed              | Non-strategic | Females migrate to central North Pacific to feed on pelagic prey.  | Le Beouf et al. 2000   |
| Northern Fur Seal           | <i>Callorhinus ursinus</i>          | Not Listed              | Non-strategic | Occur throughout the North Pacific Ocean.  | Gelatt et al. 2015   |
| Pacific White-Sided Dolphin | <i>Lagenorhynchus obliquidens</i>   | Not Listed              | Non-strategic | Endemic to temperate waters of North Pacific Ocean. Occur both on the high seas and along continental margins.   | Brownell et al. 1999   |
| Pantropical Spotted Dolphin | <i>Stenella attenuata attenuata</i> | Not Listed              | Non-strategic | Common and abundant throughout the Hawaiian archipelago. Pelagic stock occurs outside of insular stock areas (20 km for Oahu and 4-island stocks, 65 km for Hawai i Island stock). | Baird et al. 2013, Oleson et al. 2013                                    |
| Pygmy Killer Whale          | <i>Feresa attenuata</i>             | Not Listed              | Non-strategic | Small resident population in Hawaiian waters. Found worldwide in tropical and subtropical waters.  | McSweeney et al. 2009, Ross & Leatherwood 1994                           |
| Pygmy Sperm Whale           | <i>Kogia breviceps</i>              | Not Listed              | Non-strategic | Found worldwide in tropical and warm-temperate waters.   | Caldwell & Caldwell 1989   |
| Risso's Dolphin             | <i>Grampus griseus</i>              | Not Listed              | Non-strategic | Found in tropical to warm-temperate waters worldwide.  | Perrin et al. 2009   |
| Rough-Toothed Dolphin       | <i>Steno bredanensis</i>            | Not Listed              | Non-strategic | Found in tropical to warm-temperate waters worldwide. Occasionally found offshore of Hawai i.  | Perrin et al. 2009, Baird et al. 2013, Barlow 2006, Bradford et al. 2013 |
| Sei Whale                   | <i>Balaenoptera borealis</i>        | Endangered              | Strategic     | Rare in Hawai i. Generally found in offshore temperate waters.   | 35 FR 18319, Barlow 2003, Bradford et al. 2013                           |

| Common name                | Scientific name                   | ESA listing status                 | MMPA status   | Occurrence  | References  |
|----------------------------|-----------------------------------|------------------------------------|---------------|---|---|
| Short-Finned Pilot Whale   | <i>Globicephala macrorhynchus</i> | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters worldwide. Commonly observed around MHI and present around NWHI.   | Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013                           |
| Sperm Whale                | <i>Physeter macrocephalus</i>     | Endangered                         | Strategic     | Found in tropical to polar waters worldwide, most abundant cetaceans in the region. Sighted off the NWHI and the MHI.   | 35 FR 18319, Rice 1960, Lee 1993, Barlow 2006, Mobley et al. 2000, Shallenberger 1981 |
| Spinner Dolphin            | <i>Stenella longirostris</i>      | Not Listed                         | Non-strategic | Found worldwide in tropical and warm-temperate waters. Pelagic stock found outside of island-associated boundaries (10 nm).   | Perrin et al. 2009  |
| Striped Dolphin            | <i>Stenella coeruleoalba</i>      | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters throughout the world.  | Perrin et al. 2009  |
| <b>Elasmobranchs</b>       |                                   |                                    |               |   |   |
| Giant manta ray            | <i>Manta birostris</i>            | Threatened                         | N/A           | Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs. | Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.                        |
| Oceanic whitetip shark     | <i>Carcharhinus longimanus</i>    | Threatened                         | N/A           | Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C  | Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984                 |
| Scalloped hammerhead shark | <i>Sphyrna lewini</i>             | Endangered (Eastern Pacific DPS)   | N/A           | Found in coastal areas from southern California to Peru.  | Compagno 1984, Baum et al. 2007, Bester 2011  |
| Scalloped hammerhead       | <i>Sphyrna lewini</i>             | Threatened (Indo-West Pacific DPS) | N/A           | Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.         | Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993               |
| <b>Corals</b>              |                                   |                                    |               |   |   |



| Common name | Scientific name              | ESA listing status | MMPA status | Occurrence   | References |
|-------------|------------------------------|--------------------|-------------|--|------------|
| N/A         | <i>Acropora globiceps</i>    | Threatened         | N/A         | Not confirmed in Hawai i waters. Occur on upper reef slopes, reef flats, and adjacent habitats in depths ranging from 0 to 8 m   | Veron 2014 |
| N/A         | <i>Acropora jacquelineae</i> | Threatened         | N/A         | Not confirmed in Hawai i waters. Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and depth range is 10 to 35 m. | Veron 2014 |
| N/A         | <i>Acropora retusa</i>       | Threatened         | N/A         | Not confirmed in Hawai i waters. Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons, and depth range is 1 to 5 m.   | Veron 2014 |
| N/A         | <i>Acropora speciosa</i>     | Threatened         | N/A         | Not confirmed in Hawai i waters. Found in protected environments with clear water and high diversity of <i>Acropora</i> and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and have been found in mesophotic habitat (40-150 m).         | Veron 2014 |
| N/A         | <i>Euphyllia paradivisa</i>  | Threatened         | N/A         | Not confirmed in Hawai i waters. Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.   | Veron 2014 |
| N/A         | <i>Isopora crateriformis</i> | Threatened         | N/A         | Not confirmed in Hawai i waters. Found in shallow, high-wave energy environments, from low tide to at least 12 meters deep, and have been reported from mesophotic depths (less than 50 m depth).  | Veron 2014 |

| Common name          | Scientific name             | ESA listing status | MMPA status | Occurrence   | References              |
|----------------------|-----------------------------|--------------------|-------------|--|-------------------------|
| N/A                  | <i>Seriatopora aculeata</i> | Threatened         | N/A         | Not confirmed in Hawai'i waters. Found in broad range of habitats including, but not limited to, upper reef slopes, mid-slope terraces, lower reef slopes, reef flats, and lagoons, and depth ranges from 3 to 40 m. | Veron 2014              |
| <b>Invertebrates</b> |                             |                    |             |  |                         |
| Chambered nautilus   | <i>Nautilus pompilius</i>   | Threatened         | N/A         | Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m.  | 83 FR 48948, CITES 2016 |

<sup>a</sup> These species have critical habitat designated under the ESA. See Table B-4.

Table B-2. Protected species found or reasonably believed to be found near or in Hawai'i deep-set longline waters.

| Common name             | Scientific name   | ESA listing status | MMPA status | Occurrence                   | References                                |
|-------------------------|---|--------------------|-------------|------------------------------|---|
| <b>Seabirds</b>         |   |                    |             |                              |   |
| Laysan Albatross        | <i>Phoebastria immutabilis</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Black-Footed Albatross  | <i>Phoebastria nigripes</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Short-Tailed Albatross  | <i>Phoebastria albatrus</i>   | Endangered         | N/A         | Breeding visitor in the NWHI | 35 FR 8495, 65 FR 46643, Pyle & Pyle 2009 |
| Northern Fulmar         | <i>Fulmarus glacialis</i>   | Not Listed         | N/A         | Winter resident              | Pyle & Pyle 2009                          |
| Kermadec Petrel         | <i>Pterodroma neglecta</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Herald Petrel           | <i>Pterodroma arminjoniana</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Murphy's Petrel         | <i>Pterodroma ultima</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Mottled Petrel          | <i>Pterodroma inexpectata</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Juan Fernandez Petrel   | <i>Pterodroma externa</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Hawaiian Petrel         | <i>Pterodroma sandwichensis</i><br>( <i>Pterodroma phaeopygia sandwichensis</i> ) | Endangered         | N/A         | Breeding visitor in the MHI  | 32 FR 4001, Pyle & Pyle 2009              |
| White-Necked Petrel     | <i>Pterodroma cervicalis</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Bonin Petrel            | <i>Pterodroma hypoleuca</i>   | Not Listed         | N/A         | Breeding visitor in the NWHI | Pyle & Pyle 2009                          |
| Black-Winged Petrel     | <i>Pterodroma nigripennis</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Cook Petrel             | <i>Pterodroma cookii</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Stejneger Petrel        | <i>Pterodroma longirostris</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Pycroft Petrel          | <i>Pterodroma pycrofti</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Bulwer Petrel           | <i>Bulweria bulwerii</i>  | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Wedge-Tailed Shearwater | <i>Ardenna pacifica</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |
| Buller's Shearwater     | <i>Ardenna bulleri</i>  | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Sooty Shearwater        | <i>Ardenna grisea</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Short-Tailed Shearwater | <i>Ardenna tenuirostris</i>   | Not Listed         | N/A         | Migrant                      | Pyle & Pyle 2009                          |
| Christmas Shearwater    | <i>Puffinus nativitatis</i>   | Not Listed         | N/A         | Breeding visitor             | Pyle & Pyle 2009                          |

| Common name              | Scientific name  | ESA listing status | MMPA status | Occurrence                      | References                          |
|--------------------------|--|--------------------|-------------|---------------------------------|-------------------------------------|
| Newell's Shearwater      | <i>Puffinus newelli</i><br>( <i>Puffinus auricularis newelli</i> ) | Threatened         | N/A         | Breeding visitor                | 40 FR 44149,<br>Pyle & Pyle<br>2009 |
| Wilson's Storm-Petrel    | <i>Oceanites oceanicus</i>   | Not Listed         | N/A         | Migrant                         | Pyle & Pyle<br>2009                 |
| Leach's Storm-Petrel     | <i>Oceanodroma leucorhoa</i>                                       | Not Listed         | N/A         | Winter resident                 | Pyle & Pyle<br>2009                 |
| Band-Rumped Storm-Petrel | <i>Oceanodroma castro</i>  | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Tristram Storm-Petrel    | <i>Oceanodroma tristrami</i>                                       | Not Listed         | N/A         | Breeding visitor in the<br>NWHI | Pyle & Pyle<br>2009                 |
| White-Tailed Tropicbird  | <i>Phaethon lepturus</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Red-Tailed Tropicbird    | <i>Phaethon rubricauda</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Masked Booby             | <i>Sula dactylatra</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Nazca Booby              | <i>Sula granti</i>   | Not Listed         | N/A         | Vagrant                         | Pyle & Pyle<br>2009                 |
| Brown Booby              | <i>Sula leucogaster</i>  | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Red-Footed Booby         | <i>Sula sula</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Great Frigatebird        | <i>Fregata minor</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Lesser Frigatebird       | <i>Fregata ariel</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Laughing Gull            | <i>Leucophaeus atricilla</i>                                       | Not Listed         | N/A         | Winter resident in the<br>MHI   | Pyle & Pyle<br>2009                 |
| Franklin Gull            | <i>Leucophaeus pipixcan</i>  | Not Listed         | N/A         | Migrant                         | Pyle & Pyle<br>2009                 |
| Ring-Billed Gull         | <i>Larus delawarensis</i>  | Not Listed         | N/A         | Winter resident in the<br>MHI   | Pyle & Pyle<br>2009                 |
| Herring Gull             | <i>Larus argentatus</i>  | Not Listed         | N/A         | Winter resident in the<br>NWHI  | Pyle & Pyle<br>2009                 |
| Slaty-Backed Gull        | <i>Larus schistisagus</i>  | Not Listed         | N/A         | Winter resident in the<br>NWHI  | Pyle & Pyle<br>2009                 |
| Glaucous-Winged Gull     | <i>Larus glaucescens</i>   | Not Listed         | N/A         | Winter resident                 | Pyle & Pyle<br>2009                 |
| Brown Noddy              | <i>Anous stolidus</i>  | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Black Noddy              | <i>Anous minutus</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Blue-Gray Noddy          | <i>Procelsterna cerulea</i>  | Not Listed         | N/A         | Breeding visitor in the<br>NWHI | Pyle & Pyle<br>2009                 |
| White Tern               | <i>Gygis alba</i>  | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Sooty Tern               | <i>Onychoprion fuscatus</i>  | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |
| Gray-Backed Tern         | <i>Onychoprion lunatus</i>   | Not Listed         | N/A         | Breeding visitor                | Pyle & Pyle<br>2009                 |

| Common name            | Scientific name                 | ESA listing status                     | MMPA status | Occurrence  | References  |
|------------------------|---------------------------------|--|-------------|---|---|
| Little Tern            | <i>Sternula albifrons</i>       | Not Listed                             | N/A         | Breeding visitor in the NWHI  | Pyle & Pyle 2009  |
| Least Tern             | <i>Sternula antillarum</i>      | Not Listed                             | N/A         | Breeding visitor in the NWHI  | Pyle & Pyle 2009  |
| Arctic Tern            | <i>Sterna paradisaea</i>        | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| South Polar Skua       | <i>Stercorarius maccormicki</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| Pomarine Jaeger        | <i>Stercorarius pomarinus</i>   | Not Listed                             | N/A         | Winter resident in the MHI  | Pyle & Pyle 2009  |
| Parasitic Jaeger       | <i>Stercorarius parasiticus</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| Long-Tailed Jaeger     | <i>Stercorarius longicaudus</i> | Not Listed                             | N/A         | Migrant   | Pyle & Pyle 2009  |
| <b>Sea turtles</b>     |                                 |  |             |   |   |
| Green Sea Turtle       | <i>Chelonia mydas</i>           | Threatened (Central North Pacific DPS) | N/A         | Most common turtle in the Hawaiian Islands, much more common in nearshore state waters (foraging grounds) than offshore federal waters. Most nesting occurs on French Frigate Shoals in the NWHI. Foraging and haulout in the MHI.        | 43 FR 32800, 81 FR 20057, Balazs et al. 1992, Kolinski et al. 2001            |
| Green Sea Turtle       | <i>Chelonia mydas</i>           | Threatened (East Pacific DPS)          | N/A         | Nest primarily in Mexico and the Galapagos Islands. Little known about their pelagic range west of 90°W, but may range as far as the Marshall Islands. Genetic testing confirmed that they are incidentally taken in the HI DSLL fishery. | 43 FR 32800, 81 FR 20057, WPRFMC 2009, Clifton et al. 1982, Karl & Bowen 1999 |
| Hawksbill Sea Turtle   | <i>Eretmochelys imbricata</i>   | Endangered <sup>a</sup>                | N/A         | Small population foraging around Hawai'i and low level nesting on Maui and Hawai'i Islands. Occur worldwide in tropical and subtropical waters.   | 35 FR 8491, NMFS & USFWS 2007, Balazs et al. 1992, Katahira et al. 1994       |
| Leatherback Sea Turtle | <i>Dermochelys coriacea</i>     | Endangered <sup>a</sup>                | N/A         | Regularly sighted in offshore waters, especially at the southeastern end of the archipelago.  | 35 FR 8491, NMFS & USFWS 1997   |

| Common name               | Scientific name                | ESA listing status  | MMPA status   | Occurrence   | References  |
|---------------------------|--------------------------------|---|---------------|--|---|
| Loggerhead Sea Turtle     | <i>Caretta caretta</i>         | Endangered (North Pacific DPS)  | N/A           | Rare in Hawai'i. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters.                        | 43 FR 32800, 76 FR 58868, Dodd 1990, Balazs 1979  |
| Olive Ridley Sea Turtle   | <i>Lepidochelys olivacea</i>   | Threatened (Entire species, except for the breeding population on the Pacific coast of Mexico, which is listed as endangered) | N/A           | Rare in Hawai'i. Occurs worldwide in tropical and warm temperate ocean waters.   | 43 FR 32800, Pitman 1990, Balacz 1982   |
| <b>Marine mammals</b>     |                                |   |               |  |   |
| Blainville's Beaked Whale | <i>Mesoplodon densirostris</i> | Not Listed  | Non-strategic | Found worldwide in tropical and temperate waters   | Mead 1989   |
| Blue Whale                | <i>Balaenoptera musculus</i>   | Endangered  | Strategic     | Acoustically recorded off of Oahu and Midway Atoll, small number of sightings around Hawai'i. Considered extremely rare, generally occur in winter and summer. | 35 FR 18319, Bradford et al. 2013, Northrop et al. 1971, Thompson & Friedl 1982, Stafford et al. 2001 |
| Bottlenose Dolphin        | <i>Tursiops truncatus</i>      | Not Listed  | Non-strategic | Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.   | Perrin et al. 2009, Martien et al. 2012   |
| Bryde's Whale             | <i>Balaenoptera edeni</i>      | Not Listed  | Unknown       | Distributed widely across tropical and warm-temperate Pacific Ocean.   | Leatherwood et al. 1982   |
| Common Dolphin            | <i>Delphinus delphis</i>       | Not Listed  | N/A           | Found worldwide in temperate and subtropical seas.   | Perrin et al. 2009  |
| Cuvier's Beaked Whale     | <i>Ziphius cavirostris</i>     | Not Listed  | Non-strategic | Occur year round in Hawaiian waters.   | McSweeney et al. 2007   |
| Dall's Porpoise           | <i>Phocoenoides dalli</i>      | Not Listed  | Non-strategic | Range across the entire north Pacific Ocean.   | Hall 1979   |
| Dwarf Sperm Whale         | <i>Kogia sima</i>              | Not Listed  | Non-strategic | Most common in waters between 500 m and 1,000 m in depth. Found worldwide in tropical and warm-temperate waters.   | Nagorsen 1985, Baird et al. 2013  |

| Common name            | Scientific name                  | ESA listing status                     | MMPA status   | Occurrence   | References   |
|------------------------|----------------------------------|--|---------------|--|--|
| False Killer Whale     | <i>Pseudorca crassidens</i>      | Not Listed                             | Non-strategic | Found worldwide in tropical and warm-temperate waters. Pelagic stock tracked to within 11 km of Hawaiian islands.  | Stacey et al. 1994, Baird et al. 2012, Bradford et al. 2015  |
| Fin Whale              | <i>Balaenoptera physalus</i>     | Endangered                             | Strategic     | Infrequent sightings in Hawai'i waters. Considered rare in Hawai'i, though may migrate into Hawaiian waters during fall/winter based on acoustic recordings. | 35 FR 18319, Hamilton et al. 2009, Thompson & Friedl 1982  |
| Fraser's Dolphin       | <i>Lagenodelphis hosei</i>       | Not Listed                             | Non-strategic | Found worldwide in tropical waters.  | Perrin et al. 2009   |
| Guadalupe Fur Seal     | <i>Arctocephalus townsendi</i>   | Threatened                             | Strategic     | Rare sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.   | 50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987   |
| Hawaiian Monk Seal     | <i>Neomonachus schauinslandi</i> | Endangered <sup>a</sup>                | Strategic     | Endemic tropical seal. Occurs throughout the archipelago. MHI population spends some time foraging in federal waters during the day.                         | 41 FR 51611, Baker et al. 2011   |
| Humpback Whale         | <i>Megaptera novaeangliae</i>    | Delisted Due to Recovery (Hawai'i DPS) | Strategic     | Migrate through the archipelago and breed during the winter. Common during winter months, when they are generally found within the 100 m isobath.            | 35 FR 18319, 81 FR 62259, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978 |
| Killer Whale           | <i>Orcinus orca</i>              | Not Listed                             | Non-strategic | Rare in Hawai'i. Prefer colder waters within 800 km of continents.   | Mitchell 1975, Baird et al. 2006   |
| Longman's Beaked Whale | <i>Indopacetus pacificus</i>     | Not Listed                             | Non-strategic | Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. Rare in Hawai'i.                         | Dalebout 2003, Baird et al. 2013   |
| Melon-Headed Whale     | <i>Peponocephala electra</i>     | Not Listed                             | Non-strategic | Found in tropical and warm-temperate waters worldwide, found primarily in equatorial waters. Uncommon in Hawai'i.  | Perryman et al. 1994, Barlow 2006, Bradford et al. 2013  |
| Minke Whale            | <i>Balaenoptera</i>              | Not Listed                             | Non-strategic | Occur seasonally around  | Barlow 2003,   |

| Common name                 | Scientific name                     | ESA listing status      | MMPA status   | Occurrence  | References   |
|-----------------------------|-------------------------------------|-------------------------|---------------|---|--|
|                             | <i>acutorostrata</i>                |                         |               | Hawai'i   | Rankin & Barlow 2005   |
| North Pacific Right Whale   | <i>Eubalaena japonica</i>           | Endangered <sup>a</sup> | Strategic     | Extremely rare in Hawai'i waters  | 35 FR 18319, 73 FR 12024, Rowntree et al. 1980, Herman et al. 1980       |
| Northern Elephant Seal      | <i>Mirounga angustirostris</i>      | Not Listed              | Non-strategic | Females migrate to central North Pacific to feed on pelagic prey  | Le Beouf et al. 2000   |
| Northern Fur Seal           | <i>Callorhinus ursinus</i>          | Not Listed              | Non-strategic | Range across the north Pacific Ocean.   | Gelatt et al. 2015   |
| Pacific White-Sided Dolphin | <i>Lagenorhynchus obliquidens</i>   | Not Listed              | Non-strategic | Endemic to temperate waters of North Pacific Ocean. Occur both on the high seas and along continental margins.  | Brownell et al. 1999   |
| Pantropical Spotted Dolphin | <i>Stenella attenuata attenuata</i> | Not Listed              | Non-strategic | Common and abundant throughout the Hawaiian archipelago. Pelagic stock occurs outside of insular stock areas (20 km for Oahu and 4-island stocks, 65 km for Hawai'i Island stock) | Baird et al. 2013, Oleson et al. 2013                                    |
| Pygmy Killer Whale          | <i>Feresa attenuata</i>             | Not Listed              | Non-strategic | Small resident population in Hawaiian waters. Found worldwide in tropical and subtropical waters.   | McSweeney et al. 2009, Ross & Leatherwood 1994                           |
| Pygmy Sperm Whale           | <i>Kogia breviceps</i>              | Not Listed              | Non-strategic | Found worldwide in tropical and warm-temperate waters.  | Caldwell & Caldwell 1989   |
| Risso's Dolphin             | <i>Grampus griseus</i>              | Not Listed              | Non-strategic | Found in tropical to warm-temperate waters worldwide.   | Perrin et al. 2009   |
| Rough-Toothed Dolphin       | <i>Steno bredanensis</i>            | Not Listed              | Non-strategic | Found in tropical to warm-temperate waters worldwide. Occasionally found offshore of Hawai'i.   | Perrin et al. 2009, Bradford et al. 2013, Barlow 2006, Baird et al. 2013 |
| Sei Whale                   | <i>Balaenoptera borealis</i>        | Endangered              | Strategic     | Rare in Hawai'i. Generally found in offshore temperate waters.  | 35 FR 18319, Barlow 2003, Bradford et al. 2013                           |
| Short-Finned Pilot Whale    | <i>Globicephala macrorhynchus</i>   | Not Listed              | Non-strategic | Found in tropical to warm-temperate waters worldwide. Commonly observed around MHI and present around NWHI.   | Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013              |



| Common name                | Scientific name                | ESA listing status                 | MMPA status   | Occurrence  | References  |
|----------------------------|--------------------------------|------------------------------------|---------------|---|---|
| Sperm Whale                | <i>Physeter macrocephalus</i>  | Endangered                         | Strategic     | Found in tropical to polar waters worldwide, most abundant cetaceans in the region. Sighted off the NWHI and the MHI.   | 35 FR 18319, Rice 1960, Lee 1993, Barlow 2006, Mobley et al. 2000, Shallenberger 1981 |
| Spinner Dolphin            | <i>Stenella longirostris</i>   | Not Listed                         | Non-strategic | Found worldwide in tropical and warm-temperate waters. Pelagic stock found outside of island-associated boundaries (10 nm)  | Perrin et al. 2009  |
| Striped Dolphin            | <i>Stenella coeruleoalba</i>   | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters throughout the world   | Perrin et al. 2009  |
| <b>Elasmobranchs</b>       |                                |                                    |               |   |   |
| Giant manta ray            | <i>Manta birostris</i>         | Threatened                         | N/A           | Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs. | Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.                        |
| Oceanic whitetip shark     | <i>Carcharhinus longimanus</i> | Threatened                         | N/A           | Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C  | Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984                 |
| Scalloped hammerhead shark | <i>Sphyrna lewini</i>          | Endangered (Eastern Pacific DPS)   | N/A           | Found in coastal areas from southern California to Peru.  | Compagno 1984, Baum et al. 2007, Bester 2011  |
| Scalloped hammerhead shark | <i>Sphyrna lewini</i>          | Threatened (Indo-West Pacific DPS) | N/A           | Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.         | Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993               |
| <b>Corals</b>              |                                |                                    |               |   |   |
| N/A                        | <i>Acropora globiceps</i>      | Threatened                         | N/A           | Occur on upper reef slopes, reef flats, and adjacent habitats in depths ranging from 0 to 8 m.  | Veron 2014  |

| Common name   | Scientific name              | ESA listing status | MMPA status | Occurrence  | References |
|---------------|------------------------------|--------------------|-------------|---|------------|
| N/A           | <i>Acropora jacquelineae</i> | Threatened         | N/A         | Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and depth range is 10 to 35 m. | Veron 2014 |
| N/A           | <i>Acropora retusa</i>       | Threatened         | N/A         | Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons, and depth range is 1 to 5 m.   | Veron 2014 |
| N/A           | <i>Acropora speciosa</i>     | Threatened         | N/A         | Found in protected environments with clear water and high diversity of <i>Acropora</i> and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and it has been found in mesophotic habitat (40-150 m).       | Veron 2014 |
| N/A           | <i>Euphyllia paradivisa</i>  | Threatened         | N/A         | Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.   | Veron 2014 |
| N/A           | <i>Isopora crateriformis</i> | Threatened         | N/A         | Found in shallow, high-wave energy environments, from low tide to at least 12 m deep, and have been reported from mesophotic depths (less than 50 m depth).   | Veron 2014 |
| N/A           | <i>Seriatopora aculeata</i>  | Threatened         | N/A         | Found in broad range of habitats including, but not limited to, upper reef slopes, mid-slope terraces, lower reef slopes, reef flats, and lagoons, and depth ranges from 3 to 40 m.   | Veron 2014 |
| Invertebrates |                              |                    |             |   |            |

| Common name        | Scientific name           | ESA listing status | MMPA status | Occurrence  | References              |
|--------------------|---------------------------|--------------------|-------------|---|-------------------------|
| Chambered nautilus | <i>Nautilus pompilius</i> | Threatened         | N/A         | Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m. | 83 FR 48948, CITES 2016 |

<sup>a</sup> These species have critical habitat designated under the ESA. See Table B-4 .

Table B-3. Protected species found or reasonably believed to be found near or in American Samoa longline waters.

| Common name          | Scientific name                     | ESA listing status | MMPA status | Occurrence | References              |
|----------------------|-------------------------------------|--------------------|-------------|------------|-------------------------|
| <b>Seabirds</b>      |                                     |                    |             |            |                         |
| Audubon's Shearwater | <i>Puffinus lherminieri</i>         | Not Listed         | N/A         | Resident   | Craig 2005              |
| Black Noddy          | <i>Anous minutus</i>                | Not Listed         | N/A         | Resident   | Craig 2005              |
| Black-Naped Tern     | <i>Sterna sumatrana</i>             | Not Listed         | N/A         | Visitor    | Craig 2005              |
| Blue-Gray Noddy      | <i>Procelsterna cerulea</i>         | Not Listed         | N/A         | Resident   | Craig 2005              |
| Bridled Tern         | <i>Onychoprion anaethetus</i>       | Not Listed         | N/A         | Visitor    | Craig 2005              |
| Brown Booby          | <i>Sula leucogaster</i>             | Not Listed         | N/A         | Resident   | Craig 2005              |
| Brown Noddy          | <i>Anous stolidus</i>               | Not Listed         | N/A         | Resident   | Craig 2005              |
| Christmas Shearwater | <i>Puffinus nativitatis</i>         | Not Listed         | N/A         | Resident?  | Craig 2005              |
| Collared Petrel      | <i>Pterodroma brevipes</i>          | Not Listed         | N/A         | Resident?  | Craig 2005              |
| White Tern           | <i>Gygis alba</i>                   | Not Listed         | N/A         | Resident   | Craig 2005              |
| Greater Crested Tern | <i>Thalasseus bergii</i>            | Not Listed         | N/A         | Visitor    | Craig 2005              |
| Gray-Backed Tern     | <i>Onychoprion lunatus</i>          | Not Listed         | N/A         | Resident   | Craig 2005              |
| Great Frigatebird    | <i>Fregata minor</i>                | Not Listed         | N/A         | Resident   | Craig 2005              |
| Herald Petrel        | <i>Pterodroma heraldica</i>         | Not Listed         | N/A         | Resident   | Craig 2005              |
| Laughing Gull        | <i>Leucophaeus atricilla</i>        | Not Listed         | N/A         | Visitor    | Craig 2005              |
| Lesser Frigatebird   | <i>Fregata ariel</i>                | Not Listed         | N/A         | Resident   | Craig 2005              |
| Masked Booby         | <i>Sula dactylatra</i>              | Not Listed         | N/A         | Resident   | Craig 2005              |
| Newell's Shearwater  | <i>Puffinus auricularis newelli</i> | Threatened         | N/A         | Visitor    | 40 FR 44149, Craig 2005 |
| Red-Footed Booby     | <i>Sula sula</i>                    | Not Listed         | N/A         | Resident   | Craig 2005              |

| Common name                 | Scientific name  | ESA listing status                     | MMPA status | Occurrence   | References   |
|-----------------------------|--|--|-------------|--|--|
| Red-Tailed Tropicbird       | <i>Phaethon rubricauda</i>   | Not Listed                             | N/A         | Resident   | Craig 2005   |
| Short-Tailed Shearwater     | <i>Ardenna tenuirostris</i>  | Not Listed                             | N/A         | Visitor  | Craig 2005   |
| Sooty Shearwater            | <i>Ardenna grisea</i>  | Not Listed                             | N/A         | Visitor  | Craig 2005   |
| Sooty Tern                  | <i>Sterna fuscata</i>  | Not Listed                             | N/A         | Resident   | Craig 2005   |
| Tahiti Petrel               | <i>Pterodroma rostrata</i>   | Not Listed                             | N/A         | Resident   | Craig 2005   |
| Wedge-Tailed Shearwater     | <i>Ardenna pacifica</i>  | Not Listed                             | N/A         | Resident?  | Craig 2005   |
| White-Necked Petrel         | <i>Pterodroma cervicalis</i>   | Not Listed                             | N/A         | Visitor  | Craig 2005   |
| White-Faced Storm-Petrel    | <i>Pelagodroma marina</i>  | Not Listed                             | N/A         | Visitor  | Craig 2005   |
| White-Tailed Tropicbird     | <i>Phaethon lepturus</i>   | Not Listed                             | N/A         | Resident   | Craig 2005   |
| White-Throated Storm-Petrel | <i>Nesofregatta fuliginosa</i>   | Not Listed                             | N/A         | Resident?  | Craig 2005   |
| Laysan Albatross            | <i>Phoebastria immutabilis</i>   | Not Listed                             | N/A         | Breed mainly in Hawai'i, and range across the North Pacific Ocean.       | Causey 2008  |
| Hawaiian Petrel             | <i>Pterodroma sandwichensis</i> ( <i>Pterodroma phaeopygia sandwichensis</i> ) | Endangered                             | N/A         | Breed in MHI, and range across the central Pacific Ocean.                | 32 FR 4001, Simons & Hodges 1998                           |
| Laysan Albatross            | <i>Phoebastria immutabilis</i>   | Not Listed                             | N/A         | Breed mainly in Hawai'i, and range across the North Pacific Ocean.       | Causey 2009  |
| Northern Fulmar             | <i>Fulmarus glacialis</i>  | Not Listed                             | N/A         | Breed and range across North Pacific Ocean.                              | Hatch & Nettleship 2012                                    |
| Short-Tailed Albatross      | <i>Phoebastria albatrus</i>  | Endangered                             | N/A         | Breed in Japan and NWHI, and range across the North Pacific Ocean.       | 35 FR 8495, 65 FR 46643, BirdLife International 2017       |
| <b>Sea turtles</b>          |  |  |             |  |  |
| Green Sea Turtle            | <i>Chelonia mydas</i>  | Endangered (Central South Pacific DPS) | N/A         | Frequently seen. Nest at Rose Atoll in small numbers.                    | 43 FR 32800, 81 FR 20057, Balacz 1994                      |
| Hawksbill Sea Turtle        | <i>Eretmochelys imbricata</i>  | Endangered <sup>a</sup>                | N/A         | Frequently seen. Nest at Rose Atoll, Swain's Island, and Tutuila.        | 35 FR 8491, NMFS & USFWS 2013, Tuato'o-Bartley et al. 1993 |
| Leatherback Sea Turtle      | <i>Dermochelys coriacea</i>  | Endangered <sup>a</sup>                | N/A         | Very rare. One juvenile recovered dead in experimental longline fishing. | 35 FR 8491, Grant 1994                                     |

| Common name               | Scientific name                | ESA listing status  | MMPA status   | Occurrence   | References   |
|---------------------------|--------------------------------|---|---------------|--|--|
| Loggerhead Sea Turtle     | <i>Caretta caretta</i>         | Endangered (South Pacific DPS)  | N/A           | No known sightings. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters. | 43 FR 32800, 76 FR 58868, Utzurrum 2002, Dodd 1990 |
| Olive Ridley Sea Turtle   | <i>Lepidochelys olivacea</i>   | Threatened (Entire species, except for the endangered breeding population on the Pacific coast of Mexico) | N/A           | Rare. Three known sightings.   | 43 FR 32800, Utzurrum 2002                         |
| <b>Marine mammals</b>     |                                |   |               |  |  |
| Blainville's Beaked Whale | <i>Mesoplodon densirostris</i> | Not Listed  | Non-strategic | Found worldwide in tropical and temperate waters   | Mead 1989  |
| Blue Whale                | <i>Balaenoptera musculus</i>   | Endangered  | Strategic     | No known sightings. Occur worldwide, and are known to be found in the western South Pacific.   | 35 FR 18319, Olson et al. 2015                     |
| Bottlenose Dolphin        | <i>Tursiops truncatus</i>      | Not Listed  | Non-strategic | Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.                         | Perrin et al. 2009, Martien et al. 2012            |
| Bryde's Whale             | <i>Balaenoptera edeni</i>      | Not Listed  | Unknown       | Distributed widely across tropical and warm-temperate Pacific Ocean.   | Leatherwood et al. 1982                            |
| Common Dolphin            | <i>Delphinus delphis</i>       | Not Listed  | N/A           | Found worldwide in temperate and subtropical seas.   | Perrin et al. 2009                                 |
| Cuvier's Beaked Whale     | <i>Ziphius cavirostris</i>     | Not Listed  | Non-strategic | Occur worldwide.   | Heyning 1989                                       |
| Dwarf Sperm Whale         | <i>Kogia sima</i>              | Not Listed  | Non-strategic | Found worldwide in tropical and warm-temperate waters.   | Nagorsen 1985                                      |
| False Killer Whale        | <i>Pseudorca crassidens</i>    | Not Listed  | Unknown       | Found in waters within the U.S. EEZ of A. Samoa  | Bradford et al. 2015                               |
| Fin Whale                 | <i>Balaenoptera physalus</i>   | Endangered  | Strategic     | No known sightings but reasonably expected to occur in A. Samoa. Found worldwide.  | 35 FR 18319, Hamilton et al. 2009                  |
| Fraser's Dolphin          | <i>Lagenodelphis hosei</i>     | Not Listed  | Non-strategic | Found worldwide in tropical waters.  | Perrin et al. 2009                                 |

| Common name                 | Scientific name                     | ESA listing status                     | MMPA status   | Occurrence  | References   |
|-----------------------------|-------------------------------------|--|---------------|---|--|
| Guadalupe Fur Seal          | <i>Arctocephalus townsendi</i>      | Threatened                             | Strategic     | No known sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.          | 50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987   |
| Humpback Whale              | <i>Megaptera novaeangliae</i>       | Delisted Due to Recovery (Oceania DPS) | Strategic     | Migrate through the archipelago and breed during the winter in American Samoan waters.                              | 35 FR 18319, 81 FR 62259,, Guarrige et al. 2007, SPWRC 2008  |
| Killer Whale                | <i>Orcinus orca</i>                 | Not Listed                             | Non-strategic | Found worldwide. Prefer colder waters within 800 km of continents.  | Leatherwood & Dalheim 1978, Mitchell 1975, Baird et al. 2006   |
| Longman's Beaked Whale      | <i>Indopacetus pacificus</i>        | Not Listed                             | Non-strategic | Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. | Dalebout 2003  |
| Melon-Headed Whale          | <i>Peponocephala electra</i>        | Not Listed                             | Non-strategic | Found in tropical and warm-temperate waters worldwide, primarily found in equatorial waters.                        | Perryman et al. 1994   |
| Minke Whale                 | <i>Balaenoptera acutorostrata</i>   | Not Listed                             | Non-strategic | Uncommon in this region, usually seen over continental shelves in the Pacific Ocean.                                | Brueggeman et al. 1990   |
| North Pacific Right Whale   | <i>Eubalaena japonica</i>           | Endangered <sup>a</sup>                | Strategic     | Extremely rare.   | 35 FR 18319, 73 FR 12024, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978 |
| Northern Elephant Seal      | <i>Mirounga angustirostris</i>      | Not Listed                             | Non-strategic | Females migrate to central North Pacific to feed on pelagic prey  | Le Beouf et al. 2000   |
| Pantropical Spotted Dolphin | <i>Stenella attenuata attenuata</i> | Not Listed                             | Non-strategic | Found in tropical and subtropical waters worldwide.   | Perrin et al. 2009   |
| Pygmy Killer Whale          | <i>Feresa attenuata</i>             | Not Listed                             | Non-strategic | Found in tropical and subtropical waters worldwide.   | Ross & Leatherwood 1994  |
| Pygmy Sperm Whale           | <i>Kogia breviceps</i>              | Not Listed                             | Non-strategic | Found worldwide in tropical and warm-temperate waters.  | Caldwell & Caldwell 1989   |

| Common name                | Scientific name                   | ESA listing status                 | MMPA status   | Occurrence  | References  |
|----------------------------|-----------------------------------|------------------------------------|---------------|---|---|
| Risso's Dolphin            | <i>Grampus griseus</i>            | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters worldwide.   | Perrin et al. 2009  |
| Rough-Toothed Dolphin      | <i>Steno bredanensis</i>          | Not Listed                         | Unknown       | Found in tropical to warm-temperate waters worldwide. Common in A. Samoa waters.  | Perrin et al. 2009, Craig 2005  |
| Sei Whale                  | <i>Balaenoptera borealis</i>      | Endangered                         | Strategic     | Generally found in offshore temperate waters.   | 35 FR 18319, Barlow 2003, Bradford et al. 2013  |
| Short-Finned Pilot Whale   | <i>Globicephala macrorhynchus</i> | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters worldwide  | Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013                           |
| Sperm Whale                | <i>Physeter macrocephalus</i>     | Endangered                         | Strategic     | Found in tropical to polar waters worldwide, most abundant cetaceans in the region.   | 35 FR 18319, Rice 1960, Barlow 2006, Lee 1993, Mobley et al. 2000, Shallenberger 1981 |
| Spinner Dolphin            | <i>Stenella longirostris</i>      | Not Listed                         | Unknown       | Common in American Samoa, found in waters with mean depth of 44 m.  | Reeves et al. 1999, Johnston et al. 2008  |
| Striped Dolphin            | <i>Stenella coeruleoalba</i>      | Not Listed                         | Non-strategic | Found in tropical to warm-temperate waters throughout the world   | Perrin et al. 2009  |
| <b>Elasmobranchs</b>       |                                   |                                    |               |   |   |
| Giant manta ray            | <i>Manta birostris</i>            | Threatened                         | N/A           | Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs. | Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.                        |
| Oceanic whitetip shark     | <i>Carcharhinus longimanus</i>    | Threatened                         | N/A           | Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C.   | Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984                 |
| Scalloped hammerhead shark | <i>Sphyrna lewini</i>             | Threatened (Indo-West Pacific DPS) | N/A           | Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.         | Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993               |

| Common name          | Scientific name              | ESA listing status | MMPA status | Occurrence  | References |
|----------------------|------------------------------|--------------------|-------------|---|------------|
| <b>Corals</b>        |                              |                    |             |   |            |
| N/A                  | <i>Acropora globiceps</i>    | Threatened         | N/A         | Occur on upper reef slopes, reef flats, and adjacent habitats in depths from 0 to 8 m   | Veron 2014 |
| N/A                  | <i>Acropora jacquelineae</i> | Threatened         | N/A         | Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and its depth range is 10 to 35 m. | Veron 2014 |
| N/A                  | <i>Acropora retusa</i>       | Threatened         | N/A         | Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons. Depth range is 1 to 5 m.   | Veron 2014 |
| N/A                  | <i>Acropora speciosa</i>     | Threatened         | N/A         | Found in protected environments with clear water and high diversity of Acropora and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and have been found in mesophotic habitat (40-150 m).                    | Veron 2014 |
| N/A                  | <i>Euphyllia paradivisa</i>  | Threatened         | N/A         | Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.   | Veron 2014 |
| N/A                  | <i>Isopora crateriformis</i> | Threatened         | N/A         | Found in shallow, high-wave energy environments, from low tide to at least 12 meters deep, and have been reported from mesophotic depths (less than 50 m depth).  | Veron 2014 |
| <b>Invertebrates</b> |                              |                    |             |   |            |



| Common name        | Scientific name           | ESA listing status | MMPA status | Occurrence  | References              |
|--------------------|---------------------------|--------------------|-------------|---|-------------------------|
| Chambered nautilus | <i>Nautilus pompilius</i> | Threatened         | N/A         | Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m. | 83 FR 48948, CITES 2016 |

<sup>a</sup> These species have critical habitat designated under the ESA. See Table B-4.

Table B-4. ESA-listed species' critical habitat in the Pacific Ocean<sup>a</sup>.

| Common Name               | Scientific Name                  | ESA Listing Status | Critical Habitat  | References                            |
|---------------------------|----------------------------------|--------------------|---|---------------------------------------|
| Hawksbill Sea Turtle      | <i>Eretmochelys imbricata</i>    | Endangered         | None in the Pacific Ocean.  | 63 FR 46693                           |
| Leatherback Sea Turtle    | <i>Dermochelys coriacea</i>      | Endangered         | Approximately 16,910 square miles (43,798 square km) stretching along the California coast from Point Arena to Point Arguello east of the 3,000 meter depth contour; and 25,004 square miles (64,760 square km) stretching from Cape Flattery, Washington to Cape Blanco, Oregon east of the 2,000 meter depth contour. | 77 FR 4170                            |
| Hawaiian Monk Seal        | <i>Neomonachus schauinslandi</i> | Endangered         | Ten areas in the Northwestern Hawaiian Islands (NWHI) and six in the main Hawaiian Islands (MHI). These areas contain one or a combination of habitat types: Preferred pupping and nursing areas, significant haul-out areas, and/or marine foraging areas, that will support conservation for the species.             | 53 FR 18988, 51 FR 16047, 80 FR 50925 |
| North Pacific Right Whale | <i>Eubalaena japonica</i>        | Endangered         | Two specific areas are designated, one in the Gulf of Alaska and another in the Bering Sea, comprising a total of approximately 95,200 square kilometers (36,750 square miles) of marine habitat.   | 73 FR 19000, 71 FR 38277              |

<sup>a</sup> For maps of critical habitat, see <https://www.fisheries.noaa.gov/national/endangered-species-conservation/critical-habitat>.

## REFERENCES

Andrews, K.R., Karczmarski, L., Au, W.W.L., Rickards, S.H., Vanderlip, C.A., Bowen, B.W., Grau, E.G., and Toonen, R.J., 2010. Rolling stones and stable homes: social structure, habitat diversity and population genetics of the Hawaiian spinner dolphin (*Stenella longirostris*). *Molecular Ecology*, 19, pp. 732-748.

- Awkerman, J.A., Anderson, D.J., and Whittow, G.C., 2009. Laysan Albatross (*Phoebastria immutabilis*). In: Rodewald, P.G. [ed.]. The Birds of North America. Ithaca: Cornell Lab of Ornithology. Retrieved from the Birds of North America at <https://birdsna.org/Species-Account/bna/species/layalb>.
- Backus, R.H., Springer, S., and Arnold, E.L., 1956. A contribution to the natural history of the white tip shark, *Pseudocarcharias kamoharui* (Poey). *Deep-Sea Research*, 3, pp. 178-188.
- Baird, R.W., McSweeney, D.J., Bane, C., Barlow, J., Salden, D.R., Antoine, L.R.K., LeDuc, R.G. and Webster, D.L., 2006. Killer Whales in Hawaiian Waters: Information on Population Identity and Feeding Habits. *Pacific Science*, 60(4), pp. 523-530.
- Baird, R.W., Gorgone, A.M., McSweeney, D.J., Ligon, A.D., Deakos, M.H., Webster, D.L., Schorr, G.S., Martien, K.K., Salden, D.R., and Mahaffy, S.D., 2009. Population structure of island-associated dolphins: Evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian Islands. *Marine Mammal Science*, 25, pp. 251-274.
- Baird, R.W., Webster, D.L., Aschettino, J.M., Schorr, G.S. and McSweeney, D.J., 2013. Odontocete cetaceans around the main Hawaiian Islands: Habitat use and relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3), 253 pp.
- Baker J.D., Harting, A.L., Wurth, T.A., and Johanos, T.C., 2011. Dramatic shifts in Hawaiian monk seal distribution predicted from divergent regional trends. *Marine Mammal Science*, 27(1), pp. 78-93.
- Balazs, G.H. 1979. Loggerhead turtle recovered from a tiger shark at Kure Atoll. *'Elepaio*, 39(12), pp. 45-47.
- Balazs, G.H. 1982. Status of sea turtles in the central Pacific Ocean. In: Bjorndal, K.A. [ed.]. Biology and Conservation of Sea Turtles. *Smithsonian Institution Scholarly Press*, Washington, D.C., 583 pp.
- Balazs, G.H., Craig, P., Winton, B.R. and Miya, R.K., 1994. Satellite telemetry of green turtles nesting at French Frigate Shoals, Hawaii, and Rose Atoll, American Samoa. In: Bjorndal, K.A., Bolten, A.B., Johnson, D.A., and Eliazar, P.J. [eds.]. Proceedings of the fourteenth annual symposium on sea turtle biology and conservation.
- Balazs, G.H., Hirth, H., Kawamoto, P., Nitta, E., Ogren, L., Wass, R., and Wetherall, J., 1992. Interim Recovery Plan for Hawaiian Sea Turtles. Honolulu Lab, Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA, Honolulu, HI 96822-2396. Southwest Fisheries Science Center Administrative Report H-92-01. 76 pp.
- Barlow, J., 2003. Preliminary Estimates of the Abundance of Cetaceans along the U.S. West Coast, 1991-2001. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.

- Barlow, J., 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science*, 22(2), pp. 446-464.
- Baum, J., Clarke, S., Domingo, A., Ducrocq, M., Lamónaca, A.F., Gaibor, N., Graham, R., Jorgensen, S., Kotas, J.E., Medina, E., Martinez-Ortiz, J., Monzini Taccone di Sitizano, J., Morales, M.R., Navarro, S.S., Pérez-Jiménez, J.C., Ruiz, C., Smith, W., Valenti, S.V. and Vooren, C.M., 2007. *Sphyrna lewini*. The IUCN Red List of Threatened Species 2007: e.T39385A10190088. Downloaded on 21 Feb 2017.
- Baum, J.K., Medina, E., Musick, J.A., and Smale, M., 2006. *Carcharhinus longimanus*. In: IUCN 2007. 2007 IUCN Red List of Threatened Species.
- Bester, C., 2011. Species Profile: Scalloped Hammerhead. Florida Museum of Natural History. <http://www.flmnh.ufl.edu/fish/Gallery/Descript/Schammer/ScallopedHammerhead.html>.
- BirdLife International, 2017. Species factsheet: *Phoebastria albatrus*. Downloaded from <http://www.birdlife.org> on 02/04/2017.
- Bonfil, R., Clarke, S., Nakano, H., Camhi, M.D., Pikitch, E.K. and Babcock, E.A., 2008. The biology and ecology of the oceanic whitetip shark, *Carcharhinus longimanus*. In: Camhi, M.D., Pikitch, E.K., and Babcock, E.A. [eds.]. *Sharks of the Open Ocean: Biology, Fisheries, and Conservation*, pp.128-139.
- Bradford, A.L., Oleson, E.M., Baird, R.W., Boggs, C.H., Forney, K.A., and Young, N.C., 2015. Revised stock boundaries for false killer whales (*Pseudorca crassidens*) in Hawaiian waters. NOAA Tech. Memo. NMFS-PIFSC-47.
- Bradford, A.L., K.A. Forney, E.M. Oleson, and J. Barlow. 2013. Line-transect abundance estimates of cetaceans in the Hawaiian EEZ. PIFSC Working Paper WP-13-004.
- Brownell, J. R. L., W. A. Walker, and K. A. Forney. 1999. Pacific white-sided dolphin, *Lagenorhynchus obliquidens* Gill, 1865. In: Ridgeway, S.H., and Harrison, R. [eds.]. *Handbook of marine mammals: the second book of dolphins and the porpoises*. Academic Press, San Diego, CA.
- Brueggeman, J.J., Green, G.A., Balcomb, K.C., Bowlby, C.E., Grotfendt, R.A., Briggs, K.T., Bonnell, M.L., Ford, R.G., Varoujean, D.H., Heinemann, D. and Chapman D.G., 1990. Oregon-Washington Marine Mammal and Seabird Survey: Information synthesis and hypothesis formulation. U.S. Department of the Interior, OCS Study MMS 89-0030.
- Caldwell, D.K. and Caldwell, M.C., 1989. Pygmy sperm whale *Kogia breviceps* (de Blainville, 1838)/dwarf sperm whale *Kogia simus* (Owen, 1866). *Handbook of marine mammals*, 4, pp. 235-260.
- Childerhouse, S., J. Jackson, C. S. Baker, N. Gales, P. J. Clapham, and R. L. Brownell, Jr., 2008. *Megaptera novaeangliae*, Oceania subpopulation. IUCN Red List of Threatened Species <http://www.iucnredlist.org/details/132832>.

- CITES (2016) Consideration of proposals for amendment of Appendices I and II: Nautilidae Convention on International Trade in Endangered Species of Wild Fauna and Flora; Seventeenth meeting of the Conference of the Parties; 24 September – 5 October 2016, Johannesburg, South Africa
- Cliffton, K., Cornejo, D.O., and Felger, R.S., 1982. Sea turtles of the Pacific coast of Mexico. *In*: Bjorndal, K.A. [ed.]. *Biology and Conservation of Sea Turtles*. Smithsonian Inst. Press, Washington, D.C., 583 pp.
- Compagno, L.J.V., 1984. *FAO Species Catalogue. Vol. 4. Sharks of the World. An Annotated and Illustrated Catalogue of Shark Species Known to Date. Carcharhiniformes. FAO Fish Synop 124, Vol. 4, Part 2.*
- Craig, P., 2005. *Natural history guide to American Samoa. National Park of American Samoa, Department of Marine and Wildlife Resources, American Samoa Community College.*
- Dalebout, M.L., Baker, C.S., Anderson, R.C., Best, P.B., Cockcroft, V.G., Hinsz, H.L., Peddemors, V. and Pitman, R.L., 2003. Appearance, distribution, and genetic distinctiveness of Longman's beaked whale, *Indopacetus pacificus*. *Marine Mammal Science*, 19(3), pp. 421-461.
- Dewar, H., Mous, P., Domeier, M., Muljadi, A., Pet, J., and Whitty, J., 2008. Movements and site fidelity of the giant manta ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology*, 155, pp. 121-133.
- Dodd, C.K., 1990. *Caretta caretta* (Linnaeus) Loggerhead Sea Turtle. *Catalogue of American Amphibians and Reptiles*, pp. 483.1-483.7.
- Eckert, K.L., B.P. Wallace, J.G. Frazier, S.A. Eckert, and P.C.H. Pritchard. 2012. Synopsis of the biological data on the leatherback sea turtle (*Dermochelys coriacea*). U.S. Department of Interior, Fish and Wildlife Service, Biological Technical Publication BTP-R4015-2012, Washington, D.C.
- Fleischer, L.A., 1987. Guadalupe fur seal, *Arctocephalus townsendi*. *In*: Croxall, J.P. and Gentry R.L. [eds.]. *Status, biology, and ecology of fur seals*. National Oceanic and Atmospheric Association, Technical Report, National Marine Fisheries Service 51, pp. 1-212.
- Gallo-Reynoso, J.P., Figueroa-Carranza, A.L. and Le Boeuf, B.J., 2008. Foraging behavior of lactating Guadalupe fur seal females. *In*: Lorenzo, C., Espinoza, E., and Ortega, J. [eds.]. *Avances en el Estudio de los Mamíferos de México. Publicaciones Especiales*, 2, pp. 595-614.
- Garrigue, C., Franklin, T., Russell, K., Burns, D., Poole, M., Paton, D., Hauser, N., Oremus, M., Constantine, R., Childerhouse, S., Mattila, D., Gibbs, N., Franklin, W., Robbins, J., Clapham, P., and Baker, C.S., 2007. First assessment of interchange of humpback whales between Oceania and the east coast of Australia. International Whaling Commission, Anchorage, Alaska. SC/59/SH15.

- Gelatt, T., Ream, R., and Johnson, D., 2015. *Callorhinus ursinus*. The IUCN Red List of Threatened Species 2015:e.T3590A45224953. Available at <http://www.iucnredlist.org/details/3590/0>.
- Grant, G.S. 1994. Juvenile leatherback turtle caught by longline fishing in American Samoa. *Marine Turtle Newsletter*, 66, pp. 3-5.
- Hall, J., 1979. A survey of cetaceans of Prince William Sound and adjacent waters - their numbers and seasonal movements. Unpubl. rep. to Alaska Outer Continental Shelf Environmental Assessment Programs. NOAA OCSEAP Juneau Project Office, Juneau, AK. 37 pp.
- Hamilton, T.A., Redfern, J.V., Barlow, J., Balance, L.T., Gerrodette, T., Holt, R.S., Forney, K.A., and Taylor, B.L., 2009. Atlas of cetacean sightings for Southwest Fisheries Science Center Cetacean and Ecosystem Surveys: 1986-2005. U.S. Dep. of Commerce, NOAA Technical Memorandum, NOAA-TM-NMFSSWFSC-440. 70 pp.
- Herman, L.M. and Antinaja, R.C., 1977. Humpback whales in the Hawaiian breeding waters: Population and pod characteristics. *Scientific Reports of the Whales Research Institute*, 29, pp. 59-85.
- Herman, L. M., C. S. Baker, P. H. Forestell, and R. C. Antinaja. 1980. Right whale, *Balaena glacialis*, sightings near Hawaii: a clue to the wintering grounds? *Marine Ecology Progress Series*, 2, pp. 271-275.
- Heyning, J.E. 1989. Cuvier's beaked whale, *Ziphius cavirostris* (Cuvier, 1823). pp. 289-308. In: S.H. Ridgway and R. Harrison (eds.) Handbook of Marine Mammals. Vol. 4. River Dolphins and the Larger Toothed Whales. *Academic Press*, London and San Diego. 442 pp.
- Hill, M.C., Oleson, E.M., and Andrews, K.R., 2010. New island-associated stocks for Hawaiian spinner dolphins (*Stenella longirostris longirostris*): Rationale and new stock boundaries. Pacific Islands Fisheries Science Center Admin Report H-10-04, 12 pp.
- Johnston, D.W., Robbins, J., Chapla, M.E., Mattila, D.K. and Andrews, K.R., 2008. Diversity, Johnston, D.W., Robbins, J., Chapla, M.E., Mattila, D.K. and Andrews, K.R., 2008. Diversity, habitat associations and stock structure of odontocete cetaceans in the waters of American Samoa, 2003-2006. *Journal of Cetacean Research and Management*, 10(1), pp. 59-66.
- Karczmarski, L., Würsig, B., Gailey, G., Larson, K.W., and Vanderlip, C., 2005. Spinner dolphins in a remote Hawaiian atoll: social grouping and population structure. *Behavioral Ecology*, 16, pp. 675-685.
- Karl, S.A. and Bowen, B.W., 1999. Evolutionary significant units versus geopolitical taxonomy: molecular systematics of an endangered sea turtle (genus *Chelonia*). *Conservation Biology*, 13, pp. 990-999.

- Katahira, L.K., C.M. Forbes, A.H. Kikuta, G.H. Balazs, and M. Bingham. 1994. Recent findings and management of hawksbill turtle nesting beaches in Hawaii. *In: Bjordal, K.A., Bolton, A.B., Johnson, D.A., and Eliazar, P.J. [eds.], Proc. of the Fourteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-351, 323 pp.*
- Klimley, A.P., 1993. Highly directional swimming by scalloped hammerhead sharks, *Sphyrna lewini*, and subsurface irradiance, temperature, bathymetry, and geomagnetic field. *Marine Biology, 117*(1), pp. 1-22.
- Kolinski, S.P., Parker, D.M., Ilo, L.I., and Ruak, J.K., 2001. An assessment of sea turtles and their marine and terrestrial habitats at Saipan, Commonwealth of the Northern Mariana Islands. *Micronesica, 34*, pp. 55-72.
- Le Boeuf, B.J., Crocker, D.E., Costa, D.P., Blackwell, S.B., Webb, P.M. and Houser, D.S. 2000. Foraging ecology of northern elephant seals. *Ecological Monographs, 70*(3), pp. 353-382.
- Lee, T. 1993. Summary of cetacean survey data collected between the years of 1974 and 1985. NOAA Tech.Mem. NMFS 181, 184 pp.
- Leatherwood, J.S. and Dahlheim, M.E., 1978. Worldwide distribution of pilot whales and killer whales. Naval Ocean System Center Technical Report 443, pp. 1-39.
- Leatherwood, S., Reeves, R.R., Perrin, W.F., and Evans, W.E. 1982. Whales, dolphins, and porpoises of the North Pacific and adjacent arctic waters. NOAA Tech. Rep. NMFS Circ. No. 444.
- Mallory M.L., Hatch, S.A., and Nettleship, D.N. 2012. Northern Fulmar (*Fulmarus glacialis*). *In: Rodewald, P.G. [ed.]. The Birds of North America. Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America at <https://birdsna.org/Species-Account/bna/species/norful>.*
- Marshall, A., L.J.V. Compagno, and M.B. Bennett. 2009. Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Kreffft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa, 2301*, pp. 1-28.
- Marshall, A., Bennett, M.B., Kodja, G., Hinojosa-Alvarez, S., Galvan-Magana, F., Harding, M., Stevens, G., and Kashiwagi, T., 2011. *Manta birostris*. The IUCN Red List of Threatened Species 2011: e.T198921A9108067.
- Martien, K.K., Baird, R.W., Hedrick, N.M., Gorgone, A.M., Thieleking, J.L., McSweeney, D.J., Robertson, K.M. and Webster, D.L. 2012. Population structure of island-associated dolphins: Evidence from mitochondrial and microsatellite markers for common bottlenose dolphins (*Tursiops truncatus*) around the main Hawaiian Islands. *Marine Mammal Science, 25*(3), pp. 208-232.

- McSweeney, D.J., Baird, R.W., and Mahaffy, S.D., 2007. Site fidelity, associations and movements of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) beaked whales off the island of Hawai'i. *Marine Mammal Science*, 23, pp. 666-687.
- McSweeney, D.J., Baird, R.W., Mahaffy, S.D., Webster, D.L., and Schorr, G.S. 2009. Site fidelity and association patterns of a rare species: Pygmy killer whales (*Feresa attenuata*) in the main Hawaiian Islands. *Marine Mammal Science*, 25, pp. 557-572.
- Mead, J.G., 1989. Beaked whales of the genus *Mesoplodon*. In: Ridgeway, S.H. & Harrison, R. [eds.]. Handbook of marine mammals. Volume 4. River dolphins and the larger toothed whales. Academic Press Ltd: London, 452 pp.
- Mitchell, E., 1975. Report on the meeting on smaller cetaceans, Montreal, April 1-11, 1974. *Journal of the Fisheries Research Board of Canada*, 32(7), pp. 914-916
- Mobley, J.R. Jr., Spitz, S.S., Forney, K.A., Grotenfendt, R., and Forestell, P.H., 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys. Southwest Fisheries Science Center Administrative Report LJ-00-14C. La Jolla, CA 92037. 26 pp.
- Nagorsen, D., 1985. *Kogia simus*. *Mammalian Species*, 239, pp. 1-6.
- NMFS and USFWS, 1998. Recovery Plan for U.S. Pacific Populations of the Leatherback Turtle (*Dermochelys coriacea*). National Marine Fisheries Service, Silver Spring, MD. 77 pp.
- NMFS and USFWS, 2007. Hawksbill Sea Turtle (*Eretmochelys imbricata*) 5-Year Review: Summary and Evaluation. National Marine Fisheries Service, Silver Spring, Maryland, and U.S. Fish and Wildlife Service Jacksonville, Florida. 90 pp.
- Norris, K.S. and Dohl, T.P., 1980. Behavior of the Hawaiian spinner dolphin, *Stenella longirostris*. *Fisheries Bulletin*, 77, pp. 821-849.
- Norris, K.S., Würsig, B., Wells, R.S., and Würsig, M., 1994. The Hawaiian Spinner Dolphin. *University of California Press*, 408 pp.
- Northrop, J., Cummings, W.C., and Morrison, M.F. 1971. Underwater 20-Hz signals recorded near Midway Island. *Journal of the Acoustical Society of America*, 49, pp. 1909-1910.
- Oleson, E.M., Boggs, C.H., Forney, K.A., Hanson, M.B., Kobayashi, D.R., Taylor, B.L., Wade, P.R., and Ylitalo, G.M., 2010. Status Review of Hawaiian Insular False Killer Whales (*Pseudorca crassidens*) under the Endangered Species Act. U.S. Dep. Commer. NOAA Tech Memo., NOAA-TM-NMFS-PIFSC-22. 140 pp.
- Oleson, E.M., Baird, R.W., Martien, K.K., and Taylor, B.L., 2013. Island-associated stocks of odontocetes in the main Hawaiian Islands: A synthesis of available information to facilitate evaluation of stock structure. PIFSC Working Paper WP-13-003. 41 pp.

- Olson, P.A., Ensor, P., Olavarria, C., Bott, N., Constantine, R., Weir, J., Childerhouse, S., van der Linde, M., Schmitt, N., Miller, B.S. and Double, M.C., 2015. New Zealand Blue Whales: Residency, Morphology, and Feeding Behavior of a Little-Known Population. *Pacific Science*, 69(4), pp. 477-485.
- Perrin, W.F., Wursig, B., and Thewissen J.G.M. [eds.], 2009. Encyclopedia of marine mammals. *Academic Press*.
- Perryman, W.L., Au, D.W.K., Leatherwood, S., and Jefferson, T.A., 1994. Melon-headed whale *Peponocephala electra* Gray, 1846. In: Ridgway, S.H. and Harrison, R. [eds.]. Handbook of Marine Mammals, Volume 5. The first book of dolphins. *Academic Press*, London, U.K.
- Pitman, R.L. 1990. Pelagic distribution and biology of sea turtles in the eastern tropical Pacific. In: Richardson, T.H., Richardson, J.I., and Donnelly, M. [eds.]. Proc. of the Tenth Annual Workshop on Sea Turtle Biology and Conservation. U.S. Dep. of Comm., NOAA Tech. Memo. NMFS-SEFC-278. 286 pp.
- Pyle, R.L., and Pyle, P., 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, Hawaii. Version 1 (31 December 2009). Accessed from <http://hbs.bishopmuseum.org/birds/rlp-monograph>.
- Rankin, S. and Barlow, J., 2005. Source of the North Pacific “boing” sound attributed to minke whales. *Journal of the Acoustical Society of America*, 118(5), pp. 3346-3335
- Reeves, R.R., Leatherwood, S., Stone, G.S., and Eldredge, L.G., 1999. Marine Mammals in the Area Served by the South Pacific Regional Environment Program. Report of the South Pacific Regional Environment Program (P.O. Box 240, Apia, Samoa). 48 pp.
- Reeves, R.R., Leatherwood, S., and Baird, R.W., 2009. Evidence of a possible decline since 1989 in false killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands. *Pacific Science*, 63(2), pp. 253–261.
- Rice, D.W., 1960. Distribution of the bottle-nosed dolphin in the leeward Hawaiian Islands. *Journal of Mammalogy*, 41(3), pp. 407-408.
- Rice, D. W., and A. A. Wolman. 1984. Humpback whale census in Hawaiian waters—February 1977. In: Norris, K.S., and Reeves, R.R. (eds.). Report on a workshop on problems related to humpback whales (*Megaptera novaeangliae*) in Hawaii. Final report to the Marine Mammal Commission, U.S. Department of Commerce, NTIS PB-280-794.
- Ross, G.J.B. and Leatherwood, S., 1994. Pygmy killer whale *Feresa attenuata* (Gray, 1874). Handbook of Marine Mammals, Volume 5, pp. 387-404.
- Rowntree, V., Darling, J., Silber, G., and M. Ferrari, M., 1980. Rare sighting of a right whale (*Eubalaena glacialis*) in Hawaii. *Canadian Journal of Zoology*, 58, pp. 308-312.



- Sanches, J.G., 1991. Catálogo dos principais peixes marinhos da República de Guiné-Bissau. Publicações avulsas do I.N.I.P. No. 16. 429 pp. as cited in Froese, R. and D. Pauly, Editors. 2000. FishBase 2000: concepts, design and data sources. ICLARM, Los Baños, Laguna, Philippines. 344 pp.
- Schorr, G.S., Baird, R.W., Hanson, M.B., Webster, D.L., McSweeney, D.J., and Andrews, R.D., 2009. Movements of satellite-tagged Blainville's beaked whales off the island of Hawai'i. *Endangered Species Research*, 10, pp. 203-213.
- Schulze-Haugen, M. and Kohler, N.E. [eds.], 2003. Guide to Sharks, Tunas, & Billfishes of the U.S. Atlantic and Gulf of Mexico. RI Sea Grant/National Marine Fisheries Service.
- Shallenberger, E.W., 1981. The status of Hawaiian cetaceans. Marine Mammal Commission Report No. MMC-77/23 (NTIS PB82-109398).
- Stafford, K.M., Nieuwirth, S.L., and Fox, C.G., 2001. Geographic and seasonal variation of blue whale calls in the North Pacific. *Journal of Cetacean Research and Management*, 3, pp. 65-76.
- Strasburg, D.W., 1958. Distribution, abundance, and habits of pelagic sharks in the central Pacific Ocean. *Fishery Bulletin*, 138(58), pp. 335-361.
- Thompson, P.O., and Friedl, W.A., 1982. A long-term study of low frequency sounds from several species of whales off Oahu, Hawaii. *Cetology*, 45, pp. 1-19.
- Tuato'o-Bartley, N., Morrell, T.E., and Craig, P., 1993. Status of sea turtles in American Samoa in 1991. *Pacific Science*, 47(3), pp. 215-221.
- Utzurum, R., 2002. Sea turtle conservation in American Samoa. In: Kinan, I. [ed.]. Proceedings of the western Pacific sea turtle cooperative research and management workshop.
- Veron, J.E.N., 2014. Results of an update of the Corals of the World Information Base for the Listing Determination of 66 Coral Species under the Endangered Species Act. Report to the Western Pacific Regional Fishery Management Council, Honolulu, Hawaii.
- Wolman, A.A. and Jurasz, C.M., 1977. Humpback whales in Hawaii: Vessel Census, 1976. *Marine Fisheries Review*, 39(7), pp. 1-5.